

Status of Shortspine Thornyhead (*Sebastolobus alascanus*) along the US West coast in 2023

by
Madison Shipley¹
Joshua Zahner¹
Sabrina Beyer¹
Adam Hayes¹
Pierre-Yves Hervann²
Andrea Odell³
Haley Oleynik⁴
Jane Y. Sullivan⁵
Matthieu Veron⁶

¹School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat Street,
Seattle, Washington 98195

²Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic
and Atmospheric Administration, National Marine Fisheries Service, 2725 Montlake
Boulevard East, Seattle, Washington 98112

³University of California Davis, One Shields Avenue, Davis, California 95616

⁴Institute for the Oceans and Fisheries, University of British Columbia, 2202 Main Mall,
Vancouver, British Columbia Canada V6T 1Z4

⁵Alaska Fisheries Science Center, U.S. Department of Commerce, National Oceanic and
Atmospheric Administration, National Marine Fisheries Service, 17109 Point Lena Loop
Road, Juneau, Alaska 99801

⁶Alaska Fisheries Science Center, U.S. Department of Commerce, National Oceanic and
Atmospheric Administration, National Marine Fisheries Service, 7600 Sand Point Way
N.E., Seattle, Washington 98115

April 2023

© Pacific Fishery Management Council, 2023

Correct citation for this publication:

Shiple, M., J. Zahner, S. Beyer, A. Hayes, P.-. Hernvann, A. Odell, H. Oleynik, J.Y. Sullivan, M. Veron. 2023. Status of Shortspine Thornyhead (*Sebastolobus alascanus*) along the US West coast in 2023. Pacific Fishery Management Council, Portland, Oregon. 11 p.

Contents

Executive summary	i
Stock	i
Catches	i
Data and assessment	i
Stock biomass and dynamics	i
Recruitment	i
Exploitation status	i
Ecosystem considerations	i
Reference points	ii
Management performance	ii
Unresolved problems and major uncertainties	ii
Decision table and projections	ii
Scientific uncertainty	ii
Research and data needs	ii
 1 Introduction	 1
1.1 Basic Information	1
1.2 Life History	1
1.3 Ecosystem Considerations	1
1.4 Historical and Current Fishery Information	1
1.5 Summary of Management History and Performance	1
1.6 Foreign Fisheries	1
 2 Data	 1
2.1 Fishery-Dependent Data	2
2.2 Fishery-Independent Data	2
2.2.1 AFSC Slope Survey	2
2.2.2 California Collaborative Fisheries Research Program	2
2.2.3 AFSC/NWFSC West Coast Triennial Shelf Survey	3
2.2.4 NWFSC West Coast Groundfish Bottom Trawl Survey	3

2.3	Biological Data	6
2.3.1	Natural Mortality	6
2.3.2	Maturation and Fecundity	6
2.3.3	Sex Ratio	6
2.3.4	Length-Weight Relationship	6
2.3.5	Growth (Length-at-Age)	6
2.3.6	Ageing Precision and Bias	6
2.4	Environmental and Ecosystem Data	6
3	Assessment Model	6
3.1	Summary of Previous Assessments and Reviews	6
3.1.1	History of Modeling Approaches (not required for an update assessment)	6
3.1.2	Most Recent STAR Panel and SSC Recommendations (not required for an update assessment)	6
3.1.3	Response to Groundfish Subcommittee Requests (not required in draft)	6
3.2	Model Structure and Assumptions	6
3.2.1	Model Changes from the Last Assessment (not required for an update assessment)	6
3.2.2	Modeling Platform and Structure	6
3.2.3	Model Parameters	7
3.2.4	Key Assumptions and Structural Choices	7
3.3	Base Model Results	7
3.3.1	Parameter Estimates	7
3.3.2	Fits to the Data	7
3.3.3	Population Trajectory	7
3.3.4	Reference Points	7
3.4	Model Diagnostics	7
3.4.1	Convergence	8
3.4.2	Sensitivity Analyses	8
3.4.3	Retrospective Analysis	8
3.4.4	Likelihood Profiles	8
3.4.5	Unresolved Problems and Major Uncertainties	8

4	Management	8
4.1	Reference Points	8
4.2	Unresolved Problems and Major Uncertainties	8
4.3	Harvest Projections and Decision Tables	8
4.4	Evaluation of Scientific Uncertainty	8
4.5	Research and Data Needs	8
5	Acknowledgments	8
6	References	9
7	Tables	10
8	Figures	11

Executive summary

Stock

This assessment reports the status of Shortspine Thornyhead (*Sebastolobus alascanus*) off the US West coast using data through xxxx.

Catches

Replace text with trends and current levels. Include Table for last 10 years. Include Figure with long-term estimates.

Data and assessment

This assessment uses the stock assessment framework Stock Synthesis (SS3).

Replace text with date of last assessment, type of assessment model, data available, new information, and information lacking.

Stock biomass and dynamics

Replace text with trends and current levels relative to virgin or historic levels and description of uncertainty. Include Table for last 10 years. Include Figure with long-term estimates.

Recruitment

Replace text with trends and current levels relative to virgin or historic levels and description of uncertainty. Include Table for last 10 years. Include Figure with long-term estimates.

Exploitation status

Replace text with total catch divided by exploitable biomass or SPR harvest rate. Include Table for last 10 years. Include Figure with trend in f relative to target vs. trend in biomass relative to the target.

Ecosystem considerations

Replace text with a summary of reviewed environmental and ecosystem factors that appear to be correlated with stock dynamics. These may include variability in they physical environment, habitat, competitors, prey, or predators that directly or indirectly affects the

stock's status, vital rates (growth, survival, productivity/recruitment) or range and distribution. Note which, if any, ecosystem factors are used in the assessment and how (e.g., as background information, in data preparations, as data inputs, in decisions about model structure).

Reference points

Replace text with management targets and definition of overfishing, including the harvest rate that brings the stock to equilibrium at $B_{40\%}$, i.e., the B_{MSY} proxy and the equilibrium stock size that results from fishing at the default harvest rate, i.e., the F_{MSY} proxy. Include Table of estimated reference points for ssb, SPR, exploitation rate, and yield based on SSB proxy for MSY, SPR proxy for MSY, and estimated MSY values.

Management performance

Include Table of most recent 10 years of catches in comparison with OFL, ABC, HG, and OY/ACL values, overfishing levels, actual catch and discard. Include OFL (encountered), OFL (retained), and OFL (dead) if different due to discard and discard mortality.

Unresolved problems and major uncertainties

Replace text with any special issues that complicate scientific assessment, questions about the best model scenario, etc.

Decision table and projections

Replace text with projected yields (OFL, ABC, and ACL), spawning biomass, and stock depletion levels for each year. OFL calculations should be based on the assumption that future catches equal ABCs and not OFLs.

Scientific uncertainty

Replace text with the sigma value and the basis for its calculation.

Research and data needs

Replace text with information gaps that seriously impede the stock assessment.

1 Introduction

1.1 Basic Information

This assessment reports the status of Shortspine Thornyhead (*Sebastolobus alascanus*) off the US West coast using data through xxxx.

1.2 Life History

Replace text.

1.3 Ecosystem Considerations

Replace text.

1.4 Historical and Current Fishery Information

Replace text.

1.5 Summary of Management History and Performance

Replace text.

1.6 Foreign Fisheries

Replace text.

2 Data

Data comprise the foundational components of stock assessment models. The decision to include or exclude particular data sources in an assessment model depends on many factors.

These factors often include, but are not limited to, the way in which data were collected (e.g., measurement method and consistency); the spatial and temporal coverage of the data; the quantity of data available per desired sampling unit; the representativeness of the data to inform the modeled processes of importance; timing of when the data were provided; limitations imposed by the Terms of Reference; and the presence of an avenue for the inclusion of the data in the assessment model. Attributes associated with a data source can change through time, as can the applicability of the data source when different modeling approaches are explored (e.g., stock structure or time-varying processes). Therefore, the specific data sources included or excluded from this assessment should not necessarily constrain the selection of data sources applicable to future stock assessments for Shortspine Thornyhead. Even if a data source is not directly used in the stock assessment they can provide valuable insights into biology, fishery behavior, or localized dynamics.

Data from a wide range of programs were available for possible inclusion in the current assessment model. Descriptions of each data source included in the model (Figure 1) and sources that were explored but not included in the base model are provided below. Data that were excluded from the base model were explicitly explored during the development of this stock assessment or have not changed since their past exploration in a previous Shortspine Thornyhead stock assessment. In some cases, the inclusion of excluded data sources were explored through sensitivity analyses (see Section 3).

2.1 Fishery-Dependent Data

2.2 Fishery-Independent Data

2.2.1 AFSC Slope Survey

The AFSC Slope Survey (Slope Survey) operated during the months of October to November aboard the R/V *Miller Freeman*. Partial survey coverage of the US west coast occurred during the years 1988-1996 and complete coverage (north of 34°30'S) during the years 1997 and 1999-2001. Typically, only these four years that are seen as complete surveys are included in assessments.

2.2.2 California Collaborative Fisheries Research Program

Since 2007, the California Collaborative Fisheries Research Program (CCFRP) has monitored several areas in California to evaluate the performance of Marine Protected Areas (MPAs) and understand nearshore fish populations (Wendt and Starr 2009; Starr et al. 2015). In 2017, the survey expanded beyond the four MPAs in central California (Año Nuevo, Point Lobos, Point Buchon, and Piedras Blancas) to include the entire California

coast. Fish are collected by volunteer anglers aboard commercial passenger fishing vessels (CPFVs) guided by one of the following academic institutions based on proximity to fishing location: Humboldt State University; Bodega Marine Laboratories; Moss Landing Marine Laboratories; Cal Poly San Luis Obispo; University of California, Santa Barbara; and Scripps Institution of Oceanography.

Surveys consist of fishing with hook-and-line gear for 30-45 minutes within randomly chosen 500 by 500 m grid cells within and outside MPAs. Prior to 2017, all fish were measured for length and release or descended to depth; since then, some were sampled for otoliths and fin clips.

2.2.3 AFSC/NWFSC West Coast Triennial Shelf Survey

The AFSC/NWFSC West Coast Triennial Shelf Survey (Triennial Survey) was first conducted by the Alaska Fisheries Science Center (AFSC) in 1977, and the survey continued until 2004 (Weinberg et al. 2002). Its basic design was a series of equally-spaced east-to-west transects across the continental shelf from which searches for tows in a specific depth range were initiated. The survey design changed slightly over time. In general, all of the surveys were conducted in the mid summer through early fall. The 1977 survey was conducted from early July through late September. The surveys from 1980 through 1989 were conducted from mid-July to late September. The 1992 survey was conducted from mid July through early October. The 1995 survey was conducted from early June through late August. The 1998 survey was conducted from early June through early August. Finally, the 2001 and 2004 surveys were conducted from May to July.

Haul depths ranged from 91-457 m during the 1977 survey with no hauls shallower than 91 m. Due to haul performance issues and truncated sampling with respect to depth, the data from 1977 were omitted from this analysis. The surveys in 1980, 1983, and 1986 covered the US West Coast south to 36.8°N latitude and a depth range of 55-366 m. The surveys in 1989 and 1992 covered the same depth range but extended the southern range to 34.5°N (near Point Conception). From 1995 through 2004, the surveys covered the depth range 55-500 m and surveyed south to 34.5°N. In 2004, the final year of the Triennial Survey series, the Northwest Fisheries Science Center (NWFSC) Fishery Resource and Monitoring Division (FRAM) conducted the survey following similar protocols to earlier years.

2.2.4 NWFSC West Coast Groundfish Bottom Trawl Survey

The NWFSC West Coast Groundfish Bottom Trawl Survey (WCGBTS) is based on a random-grid design; covering the coastal waters from a depth of 55-1,280 m (Bradburn et al. 2011). This design generally uses four industry-chartered vessels per year assigned to a roughly equal number of randomly selected grid cells and divided into two ‘passes’ of the

coast. Two vessels fish from north to south during each pass between late May to early October. This design therefore incorporates both vessel-to-vessel differences in catchability, as well as variance associated with selecting a relatively small number (approximately 700) of possible cells from a very large set of possible cells spread from the Mexican to the Canadian borders.

2.3 Biological Data

2.3.1 Natural Mortality

2.3.2 Maturation and Fecundity

2.3.3 Sex Ratio

2.3.4 Length-Weight Relationship

2.3.5 Growth (Length-at-Age)

2.3.6 Ageing Precision and Bias

2.4 Environmental and Ecosystem Data

3 Assessment Model

3.1 Summary of Previous Assessments and Reviews

3.1.1 History of Modeling Approaches (not required for an update assessment)

3.1.2 Most Recent STAR Panel and SSC Recommendations (not required for an update assessment)

3.1.3 Response to Groundfish Subcommittee Requests (not required in draft)

3.2 Model Structure and Assumptions

3.2.1 Model Changes from the Last Assessment (not required for an update assessment)

3.2.2 Modeling Platform and Structure

General model specifications (e.g., executable version, model structure, definition of fleets and areas)

3.2.3 Model Parameters

Describe estimated vs. fixed parameters, priors

3.2.4 Key Assumptions and Structural Choices

3.3 Base Model Results

3.3.1 Parameter Estimates

3.3.2 Fits to the Data

3.3.3 Population Trajectory

3.3.4 Reference Points

3.4 Model Diagnostics

Describe all diagnostics

3.4.1 Convergence

3.4.2 Sensitivity Analyses

3.4.3 Retrospective Analysis

3.4.4 Likelihood Profiles

3.4.5 Unresolved Problems and Major Uncertainties

4 Management

4.1 Reference Points

4.2 Unresolved Problems and Major Uncertainties

4.3 Harvest Projections and Decision Tables

4.4 Evaluation of Scientific Uncertainty

4.5 Research and Data Needs

5 Acknowledgments

Here are all the mad props!

6 References

- Bradburn, M.J., Keller, A.A., and Horness, B.H. 2011. The 2003 to 2008 US West Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and California: Estimates of distribution, abundance, length, and age composition. US Department of Commerce, National Oceanic; Atmospheric Administration, National Marine Fisheries Service.
- Starr, R.M., Wendt, D.E., Barnes, C.L., Marks, C.I., Malone, D., Waltz, G., Schmidt, K.T., Chiu, J., Launer, A.L., Hall, N.C., and Yochum, N. 2015. Variation in responses of fishes across multiple reserves within a network of marine protected areas in temperate waters. *PLoS One* **10**(3): p.e0118502.
- Weinberg, K.L., Wilkins, M.E., Shaw, F.R., and Zimmermann, M. 2002. The 2001 Pacific West Coast bottom trawl survey of groundfish resources: Estimates of distribution, abundance and length and age composition. {NOAA} {Technical} {Memorandum}, U.S. Department of Commerce.
- Wendt, D.E., and Starr, R.M. 2009. Collaborative research: An effective way to collect data for stock assessments and evaluate marine protected areas in California. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*. **1**: 315–324.

7 Tables

8 Figures




Figure 1: Summary of data sources used in the base model.