

Hilbert C^* -bimodules and conjugation theory

Matthew Daws

March 2025

1 Introduction

We summarise some of the ideas from [4], and clarify some of the proofs. We very closely study the [4, Section 2], giving almost complete proofs, and clarifying some points. Much of the rest of this document currently is then just a summary.¹

After a little time, I found that actually, for my purposes, the paper [5] is a somewhat more gentle introduction to similar ideas.

2 bi-Hilbertian modules

We follow [4, Section 2]. We assume the basic theory of Hilbert C^* -modules, see Lance's book [6], for example.

For $x, y \in X$ we write $\theta_{x,y}$ for the adjointable operator $z \mapsto x(y|z)$. Write $\mathcal{K}_0(X)$ for the linear span of such operators, write $\mathcal{K}(X)$ for the (operator) norm closure, and write $\mathcal{L}(X)$ for the algebra of all adjointable operators.

An A - B -bimodule is a Hilbert C^* -module X over B , with a $*$ -homomorphism $\phi: A \rightarrow \mathcal{L}(X)$ turning X into a left A -module. We caution that terminology is confused here: something this notion is termed a “correspondence”: sometimes a “bimodule” means something somewhat stronger, see [8, Section 1.5] for example. We write ${}_A X_B$ if we wish to stress which algebra is acting on the left/right.

We also have the notion of a *left* Hilbert C^* -module. An A - B -bimodule ${}_A X_B$ is *bi-Hilbertian* if it is both a (right) Hilbert C^* -module over B , a left Hilbert C^* -module over A , such that both structures lead to bimodules in the above sense, and such that the induced norms are equivalent: there are constants $\lambda, \lambda' > 0$ such that

$$\lambda' \|(x|x)_B\| \leq \|{}_A(x|x)\| \leq \lambda \|(x|x)_B\| \quad (x \in X).$$

That we get bimodules means, for example, that $(x|ay)_B = (a^*x|y)_B$ and ${}_A(x|yb) = {}_A(xb^*|y)$. We stress that no further compatibility is assumed between $(\cdot|\cdot)_B$ and ${}_A(\cdot|\cdot)$ (but see Section 2.2 below).

Proposition 2.1 (See [4, Proposition 2.4]). *Suppose that $\lambda' \|(x|x)_B\| \leq \|{}_A(x|x)\|$ for each $x \in X$. Then for any $x_1, \dots, x_n \in X$ we have*

$$\lambda' \left\| \sum_{i=1}^n \theta_{x_i, x_i} \right\| \leq \left\| \sum_{i=1}^n {}_A(x_i|x_i) \right\|.$$

Proof. I don't follow the (end of the) proof given in [4], so we give the details.

Let $T \in M_n(B)$ be the matrix with i, j entry $(x_i|x_j)_B$. As shown in [3, Lemma 2.1], we have that $\|T\| = \left\| \sum_{i=1}^n \theta_{x_i, x_i} \right\|$. Further, we have that

$$\|T\| = \sup \left\{ \left\| \sum_{i,j=1}^n b_i^*(x_i|x_j)b_j \right\| : \left\| \sum_{i=1}^n b_i^*b_i \right\| \leq 1 \right\}.$$

¹Due to lack of time and/or motivation.

For any choice of (b_i) with $\sum_i b_i^* b_i \leq 1$, set $y = \sum_i x_i b_i$, so that

$$\lambda' \|(y|y)_B\| \leq \|_A(y|y)\| = \left\| \sum_{j=1}^n {}_A(y b_j^* | x_j) \right\| \leq \|(y b_j^*)_j\| \|(x_j)_j\|. \quad (1)$$

The final inequality comes from Cauchy–Schwarz applied to the Hilbert C^* -module $X \otimes \mathbb{C}^n$ with A -valued inner product $((w_i)|(z_i)) = \sum_i {}_A(w_i|z_i)$. Thus, by definition,

$$\|(y b_j^*)_j\| = \left\| \sum_j {}_A(y b_j^* | y b_j^*) \right\|^{1/2} = \left\| \sum_j {}_A(y b_j^* b_j | y) \right\|^{1/2} \leq \|y\|_A \left\| \sum_j b_j^* b_j \right\|^{1/2} \leq \|y\|_A. \quad (2)$$

As $\|_A(y|y)\| = \|y\|_A^2$, combining (1) and (2) shows that

$$\|y\|_A^2 \leq \|y\|_A \left\| \sum_j {}_A(x_j | x_j) \right\|^{1/2} \implies \|y\|_A^2 \leq \left\| \sum_{j=1}^n {}_A(x_j | x_j) \right\|.$$

Taking the supremum over such (y_i) yields the result. \square

As shown in [4, Proposition 1.4], there is a directed set Λ such that for each $\lambda \in \Lambda$, there is a finite set $u_\lambda \subseteq X$ so that, when we set $T_\lambda = \sum_{x \in u_\lambda} \theta_{x,x}$, we have that $(T_\lambda)_{\lambda \in \Lambda}$ is an increasing bounded approximate identity for $\mathcal{K}(X)$, of norm ≤ 1 . We term (u_λ) a *generalised basis* for X .

Let ${}_A X_B$ be bi-Hilbertian. We wish to show that there is a linear map

$$F: \mathcal{K}_0(X) \rightarrow A; \quad \theta_{x,y} \mapsto {}_A(x|y).$$

The only non-trivial thing to check is that if $\sum_{i=1}^n \theta_{x_i, y_i} = 0$ then $\sum_{i=1}^n {}_A(x_i | y_i) = 0$. This seems not to be obvious; here is one way to show this. Let (u_λ) be a generalised basis, and for each $\lambda \in \Lambda$, set

$$F_\lambda: \mathcal{L}(X_B) \rightarrow A; \quad T \mapsto \sum_{y \in u_\lambda} {}_A(Ty | y).$$

As $\lim_\lambda \sum_{y \in u_\lambda} \theta_{y,y}(z) = \lim_\lambda \sum_{y \in u_\lambda} y(y|z)_B = y$ for $\|\cdot\|_B$, and hence also for $\|\cdot\|_A$ as these norms are equivalent, we see that

$$F_\lambda(\theta_{x,z}) = \sum_{y \in u_\lambda} {}_A(x(z|y)_B | y) = \sum_{y \in u_\lambda} {}_A(x|y(y|z)_B) \rightarrow {}_A(x|z).$$

In fact, this only uses that $\|_A(x|x)\| \leq \lambda \|(x|x)_B\|$ for all $x \in X$. We can now conclude that

$$\sum_{i=1}^n \theta_{x_i, y_i} = 0 \implies \sum_{i=1}^n {}_A(x_i | y_i) = \sum_{i=1}^n \lim_\lambda F_\lambda(\theta_{x_i, y_i}) = \lim_\lambda F_\lambda\left(\sum_{i=1}^n \theta_{x_i, y_i}\right) = 0.$$

So F exists.

We can now give some properties of F ; see Lemma A.1 below for context, especially for (d).

Proposition 2.2 ([4, Lemma 2.6]). *The map $F: \mathcal{K}(X_B) \rightarrow A; \theta_{x,y} \mapsto {}_A(x|y)$ satisfies:*

- (a) $F(T^*T) \geq 0$ for $T \in \mathcal{K}_0(X_B)$;
- (b) $F(T)^* = F(T^*)$ for $T \in \mathcal{K}_0(X_B)$;
- (c) letting $\phi: A \rightarrow \mathcal{L}(X_B)$ be the left-module action, we have $F(\phi(a)T) = aF(T)$ and $F(T\phi(a)) = F(T)a$ for $a \in A, T \in \mathcal{K}_0(X_B)$;
- (d) supposing also the lower inequality $\lambda' \|(x|x)_B\| \leq \|_A(x|x)\|$, we have $\|F(T)\| \geq \lambda' \|T\|$ for any T of the form $\sum_{i=1}^n \theta_{x_i, x_i}$.

Proof. Part (a) follows from Lemma A.1. Part (b) is immediate. Part (c) follows from the identities $\phi(a)\theta_{x,y} = \theta_{a \cdot x, y}$ and $\theta_{x,y}\phi(a) = \theta_{x, a^* \cdot y}$. Proposition 2.1 gives (d). \square

When do we have the analogue of Proposition 2.1 for the right-hand inequality? This doesn't always hold, but the following gives some equivalent conditions.

Proposition 2.3 ([4, Proposition 2.7]). *The following are equivalent:*

1. *there is $\lambda > 0$ so that for all $x_1, \dots, x_n \in X$, we have*

$$\left\| \sum_{j=1}^n {}_A(x_i|x_i) \right\| \leq \lambda \left\| \sum_{j=1}^n \theta_{x_i, x_i} \right\|;$$

2. *there is $\lambda > 0$ so that for all $x_1, \dots, x_n, y_1, \dots, y_n \in X$, we have*

$$\left\| \sum_{j=1}^n {}_A(x_i|y_i) \right\| \leq \lambda \left\| \sum_{j=1}^n \theta_{x_i, y_i} \right\|;$$

3. *$F(T) \geq 0$ for each $T \in \mathcal{K}_0(X_B) \cap \mathcal{K}(X_B)^+$, and $\lambda = \sup_{\lambda} \|F(\sum_{y \in u_\lambda} \theta_{y,y})\| < \infty$ for some generalised basis (u_λ) .*

When these conditions hold, the smallest constants $\lambda > 0$ for which 1 and 2 hold agree, and these agree with the λ in 3. In particular, condition 3 does not depend on the choice of generalised basis.

The smallest such $\lambda > 0$ is the *right numerical index* of X , denoted $r\text{-I}[X]$. We define the contragradient module \bar{X} , Appendix A.1, which is a B - A -bimodule, and then the *left numerical index* is $l\text{-I}[X] = r\text{-I}[\bar{X}]$.

Notice that condition 2 is simply the statement that $\|F\| \leq \lambda$.

Proposition 2.4 ([4, Corollary 2.11]). *Let X have finite right numerical index. Let λ' be the constant in our assumed inequality $\lambda' \|(x|x)_B\| \leq \|{}_A(x|x)\|$ for each $x \in X$, and let $\phi: A \rightarrow \mathcal{L}(X)$ be the left action. We have that $\phi(F(T)) \geq \lambda' T$ for $T \in \mathcal{K}(X)^+$.*

Proof. By Proposition 2.2(d) combined with Lemma A.1, we have that $\|F(T)\| \geq \lambda' \|T\|$ for any $T = S^*S$ with $S \in \mathcal{K}_0(X)$. By continuity, the same holds for $T = S^*S$ with $S \in \mathcal{K}(X)$, that is, for any $T \in \mathcal{K}(X)^+$.

We now adapt an argument from [2, Pages 90–91]. For $\epsilon > 0$ we have that $\epsilon + F(T)$ is invertible (as $F(T) \geq 0$) and positive, and so we can set $S = T^{1/2} \phi((\epsilon + F(T))^{-1/2}) = T^{1/2} (\epsilon + \phi(F(T)))^{-1/2} \in \mathcal{K}(X_B)$. Then

$$F(S^*S) = F((\epsilon + \phi(F(T)))^{-1/2} T (\epsilon + \phi(F(T)))^{-1/2}) = (\epsilon + F(T))^{-1/2} F(T) (\epsilon + F(T))^{-1/2},$$

in the last step using Proposition 2.2(c). So $F(S^*S) \leq 1$, and hence $\lambda' \|S^*S\| \leq \|F(S^*S)\| \leq 1$. Hence $\lambda' S^*S \leq 1$ and so $\lambda' (\epsilon + \phi(F(T)))^{-1/2} T (\epsilon + \phi(F(T)))^{-1/2} \leq 1$. Multiplying both sides by $(\epsilon + \phi(F(T)))^{1/2}$ we obtain $\lambda' T \leq \epsilon + \phi(F(T))$ and letting $\epsilon \rightarrow 0$, we conclude that $\lambda' T \leq \phi(F(T))$ as claimed. \square

2.1 Examples

2.1.1 Conditional expectations

Let $B \subseteq A$ be an inclusion of C^* -algebras, and let $E: A \rightarrow B$ be a conditional expectation. This means that E is contractive and $E(b) = b$ for $b \in B$; then automatically E is (completely) positive and a bimodule map over A , see [10, Theorem III.3.4]. Assume there is a constant $\lambda > 0$ so that $\|E(a)\| \geq \lambda \|a\|$ for each $a \in A^+$.

We turn ${}_A X_B = A$ into a A - B -bimodule for the obvious actions, B -valued inner-product $(x|y)_B = E(x^*y)$ and A -valued inner-product ${}_A(x|y) = xy^*$. As E is contractive, and by our assumption, we have for each $x \in X$,

$$\|(x|x)_A\| = \|E(x^*x)\| \leq \|x^*x\| = \|x\|^2 = \|x^*\|^2 = \|xx^*\| = \|{}_B(x|x)\| \leq \lambda^{-1} \|E(x^*x)\| = \lambda^{-1} \|(x|x)_A\|,$$

and so the two inner-products give equivalent norms.

Consider the contragradient module ${}_B\overline{X}_A$, see Appendix A.1, so the right inner-product is $(\overline{x}|\overline{y})_A = {}_A(x|y) = xy^*$. The map $u: \overline{X} \rightarrow A; \overline{x} \mapsto x^*$ is hence linear, bijective, and an isometry for the right-norm. Using (6) we see that

$$u\theta_{\overline{x},\overline{y}}u^{-1}(z) = u\theta_{\overline{x},\overline{y}}(\overline{z^*}) = u(\overline{{}_A(z^*|y) \cdot x}) = u(\overline{z^*y^*x}) = x^*yz, \quad (3)$$

and so $\theta_{\overline{x},\overline{y}}$ is identified with x^*y . Hence [3, Lemma 2.1] (also used in the proof of Proposition 2.1) shows that

$$\left\| \sum_{i=1}^n x_i^* x_i \right\|_A = \left\| \sum_{i=1}^n \theta_{\overline{x_i}, \overline{x_i}} \right\| = \left\| ((\overline{x_i}|\overline{x_i})_A)_{i,j} \right\|_{M_n(A)} = \left\| ({}_A(x_i|x_i))_{i,j} \right\|_{M_n(A)} = \left\| (x_i x_i^*)_{i,j} \right\|_{M_n(A)}.$$

Analogously, [3, Lemma 2.1] applied to X shows that

$$\left\| (E(x_i^* x_j))_{i,j} \right\|_{M_n(A)} = \left\| ((x_i|x_j)_B)_{i,j} \right\|_{M_n(A)} = \left\| \sum_{i=1}^n \theta_{x_i, x_i} \right\|. \quad (4)$$

As the two inner-products give equivalent norms, as ${}_A X = A$ is complete, also X_B is complete, and so [2, Theorem 1] implies that the map $(E - \lambda \text{id}_A)$ is positive (this is essentially the argument we used in the proof of Proposition 2.4 above). In fact, [2, Theorem 1] gives us more: there is $\lambda' > 0$ so that $(E - \lambda' \text{id}_A)$ is completely positive. Together with (3) and (4) we obtain

$$\left\| \sum_{i=1}^n \theta_{x_i, x_i} \right\| = \left\| (E(x_i^* x_j))_{i,j} \right\|_{M_n(A)} \geq \lambda' \left\| (x_i^* x_j)_{i,j} \right\|_{M_n(A)} = \left\| \sum_{i=1}^n x_i x_i^* \right\|_A = \left\| \sum_{i=1}^n {}_A(x_i|x_i) \right\|_A.$$

Hence we've verified the first condition in Proposition 2.3, and so X has finite right numerical index, with constant at most $1/\lambda'$. [4, Proposition 2.12] shows we have equality, $r\text{-I}[X] = 1/\lambda'$.

Furthermore, to verify [4, Proposition 2.12] for \overline{X} , for constant $\lambda'' > 0$, we wish to show that

$$\left\| \sum_{i=1}^n {}_B(\overline{x_i}|\overline{x_i}) \right\| \leq \lambda'' \left\| \sum_{i=1}^n \theta_{\overline{x_i}, \overline{x_i}} \right\|.$$

Using the map u again, this is equivalent to

$$\left\| E\left(\sum_{i=1}^n x_i^* x_i\right) \right\| = \left\| \sum_{i=1}^n (x_i|x_i)_B \right\| \leq \lambda'' \left\| \sum_{i=1}^n x_i^* x_i \right\|.$$

Clearly this holds with $\lambda'' = 1$ and this is the best constant. So $l\text{-I}[X] = 1$.

Following Watatani, [11, Definition 1.2.2], a *quasi-basis* for E is a finite family $(u_1, v_1), \dots, (u_n, v_n)$ in $A \times A$ with

$$\sum_{i=1}^n u_i E(v_i x) = \sum_{i=1}^n E(x u_i) v_i = x \quad (x \in A). \quad (5)$$

The *index* of E is $\text{Index}(E) = \sum_i u_i v_i$. Then [11, Proposition 2.6.2] shows that

$$E(a) \geq \|\text{Index}(E)\|^{-1} a \quad (a \in A^+).$$

Hence such a conditional expectation fits into this framework (note that in [11] we always assume that E is faithful).

2.1.2 Finite basis

Remember that X has a finite basis if there are $(u_i)_{i=1}^n$ in X with $\sum_i \theta_{u_i, u_i} = \text{id}_X$. By comparison, notice that (5) can be re-written as

$$\sum_i \theta_{u_i, v_i^*}(x) = \sum_i u_i \cdot (v_i^*|x) = x, \quad \sum_i \theta_{v_i^*, u_i}(x^*) = \sum_i v_i^* E(u_i^* x^*) = x^* \quad (x \in X),$$

and so a quasi-basis has $\sum_i \theta_{u_i, v_i^*} = \text{id}_X$ (as then taking the adjoint shows that $\sum_i \theta_{v_i^*, u_i} = \text{id}_X$). In fact, [11, Lemma 2.1.6] shows that we can always suppose that $v_i = u_i^*$ and so we have a finite basis.

If we have a finite basis $(u_i)_{i=1}^n$ then $\mathcal{K}_0(X) = \mathcal{K}(X) = \mathcal{L}(X)$, and so F obviously extends to $\mathcal{K}(X)$. Hence $r\text{-I}[X] = \|F(1)\|$ by Proposition 2.3.

It is hence natural to also want \overline{X} to have a finite basis. See [5] for much more in this setting.

2.2 Imprimitivity bimodules

We depart from Rieffel's original definition in [9] and instead follow the very readable [1]. Of course, this material is by now well-covered in many sources. A bi-Hilbertian module is a *Hilbert A-B-bimodule* when we have the stronger compatibility between the two inner-products:

$${}_A(x|y) \cdot z = a \cdot (y|z)_B.$$

This automatically implies that the left A-action (respectively, right B-action) is adjointable, see [1, Remark 1.9].

Let $I_A \subseteq A$ be the closed ideal generated by $\{{}_A(x|y) : x, y \in X\}$, and $I_B \subseteq B$ the closed ideal generated by $\{(x|y)_B : x, y \in X\}$. This, contrary to Rieffel, we do not assume $I_A = A$ and $I_B = B$. By [1, Proposition 1.10] we have that $\mathcal{K}(X) \cong I_A$ and $\mathcal{K}(\bar{X}) \cong I_B$. Furthermore, [1, Corollary 1.11] shows that $\|{}_A(x|x)\| = \|(x|x)_B\|$ for each $x \in X$. Thus in our main assumed inequality we can take $\lambda = \lambda' = 1$.

See [5, Corollary 1.28] for a characterisation of Imprimitivity bimodules in the main setting of this note.

3 Further constructions

We recall from e.g. [6, Proposition 2.5] the notion of a $*$ -homomorphism being *non-degenerate*. As argued on [6, Page 5] when (u_i) is an approximate identity for B , we have that $x \cdot u_i \rightarrow x$ for each $x \in X_B$.

Lemma 3.1 ([4, Proposition 2.16]). *When ${}_A X_B$ is bi-Hilbertian, the $*$ -homomorphism $\phi : A \rightarrow \mathcal{L}(X_B)$ is non-degenerate.*

Proof. By the observation about approximate identities, $A \cdot X$ is dense in X for the norm from the A -valued inner-product. Hence also for the norm from the B -valued inner product, which is exactly the definition of ϕ begin non-degenerate. \square

We can always extend ϕ by weak*-weak*-continuity to a $*$ -homomorphism $\phi^{**} : A^{**} \rightarrow \mathcal{K}(X)^{**}$. To do this, we first use that $\mathcal{L}(X)$ is the multiplier algebra of $\mathcal{K}(X)$, and that $M(C) \subseteq C^{**}$ for any C^* -algebra C . Thus ϕ can be considered as a $*$ -homomorphism $A \rightarrow \mathcal{K}(X)^{**}$, and then we take the normal extension. Then ϕ^{**} will be unital, as ϕ is non-degenerate.

Let ${}_A X_B$ have finite right numerical index, so that $F : \mathcal{K}(X_B) \rightarrow A$ is continuous. Hence F extends to $F^{**} : \mathcal{K}(X_B)^{**} \rightarrow A^{**}$. The *right index element* is $F^{**}(1) \in A^{**}$ denoted $r\text{-Ind}[X]$. Letting (u_λ) be a generalised basis for X_B , we have that $T_\lambda = \sum_{y \in u_\lambda} \theta_{y,y}$ is an increasing approximate identity for $\mathcal{K}(X_B)$ and so increases to 1 in $\mathcal{K}(X_B)^{**}$. Hence $r\text{-Ind}[X] = \lim_\lambda F(T_\lambda)$ and so by Proposition 2.33, $r\text{-I}[X] = \|r\text{-Ind}[X]\|$.

4 Categories of bimodules

We follow [4, Section 4]. Let \mathcal{A} be a class of C^* -algebras. Objects of the category $\mathcal{H}_{\mathcal{A}}$ are (right) Hilbert C^* -modules over elements $B \in \mathcal{A}$, endowed with a non-degenerate left action of $A \in \mathcal{A}$, say ${}_A X_B$. The hom space is either empty, or $\text{hom}({}_A X_B, {}_A Y_B) = \mathcal{L}(X_B, Y_B)$ the adjointable linear maps, when X, Y are over the same $A, B \in \mathcal{A}$. We use the inner tensor product, so ${}_A X_B \otimes_B {}_B Y_C$. Given $T \in \mathcal{L}(X_B, X'_B)$ we can form $T \otimes \text{id}_Y$ which is adjointable.

Any $B \in \mathcal{A}$ gives $\iota_B = {}_B B_B \in \mathcal{H}_{\mathcal{A}}$ in the obvious way, and ${}_A X_B \otimes \iota_B \cong X$ canonically. We also have $\iota_A \otimes {}_A {}_A X_B \cong X$ as the left action is assumed non-degenerate.

Notice that $\mathcal{H}_{\mathcal{A}}$ is not quite a tensor $2\text{-}C^*$ -category, because $\text{id}_X \otimes T$ need not exist. However, if we look at elements $T \in \mathcal{L}(X_B, X'_B)$ which commute with the left A action, then $\text{id}_X \otimes T$ will exist, see Proposition A.2. Write ${}_{\mathcal{A}}\mathcal{H}_{\mathcal{A}}$ for this category with morphism bimodule maps.

We can also introduce a “weak-closure” of this: we have the same objects, but take the weak closure of $\mathcal{K}(X_B, Y_B)$ in a suitable bidual-like construction. See [4, Proposition 4.2] for a clearer statement.²

²But be aware that the proof of this proposition seems to end midway through the given proof: I think the last two paragraphs should not be in the proof, but rather the main body of the text.

We can now introduce the notion of a *conjugate* following [7]. We state this just for ${}_{\mathcal{A}}\mathcal{H}_{\mathcal{A}}$. Given ${}_AX_B$, a conjugate is ${}_BY_A$ if there are $R \in {}_A\mathcal{L}_B(B, Y \otimes_A X)$ and $\bar{R} \in {}_B\mathcal{L}_A(A, X \otimes_B Y)$ which satisfy the conjugate equations

$$(\bar{R}^* \otimes I_X)(I_X \otimes R) = I_X, \quad (R^* \otimes I_Y)(I_Y \otimes \bar{R}) = I_Y.$$

To interpret these, we repeatedly use that $X \otimes_B B \cong X$, and so forth.

To avoid the weak completions, we shall just state results under slightly stronger hypotheses. Following [4, Definition 2.23] we say that a bi-Hilbertian X is of *finite right index* when X is of finite right numerical index, and $r\text{-Ind}[X] \in M(A)$. (Recall that by definition, this element is only in A^{**} in general.) Similarly *finite left index* and *finite index* if both.

When X is of finite index, we find that \bar{X} is a conjugate of X , with intertwiners

$$\bar{R}^*(x \otimes \bar{y}) = {}_A(x|y), \quad R^*(\bar{x} \otimes y) = (x|y)_B.$$

See [4, Theorem 4.4].

There is a converse, [4, Theorem 4.13]. Let X_B be a right Hilbert C^* -module over B , and suppose there is a non-degenerate $\phi: A \rightarrow {}_AX_B$, so X is an A - B -bimodule (but not assumed bi-Hilbertian). Suppose that Y is a conjugate object, with intertwiners R, \bar{R} . We can then define

$${}_A(x|a \cdot y) = \bar{R}^*(\theta_{x,y} \otimes I_Y)\bar{R}(a^*) \quad (a \in A, x, y \in X).$$

This gives an A -valued inner-product making X bi-Hilbertian, of finite index. Once we have this structure, Y is bi-unitarily equivalent to \bar{X} , with the intertwiners taking the above form.

So a characterisation of Imprimitivity bimodules (aka ‘‘Strong Morita Equivalence’’) see [4, Corollary 4.14].

A Results on Hilbert modules

I don’t know a reference for the following.

Lemma A.1. *Let X_B be a Hilbert C^* -module over B . For $T \in \mathcal{K}_0(X_B)$ we have that $T^*T = \sum_{i=1}^n \theta_{z_i, z_i}$ for some z_i in X .*

Proof. Compare <https://mathoverflow.net/questions/488971>. Let $T = \sum_{i=1}^n \theta_{x_i, y_i}$, so that $T^*T = \sum_{i,j} \theta_{y_i \cdot (x_i|x_j), y_j}$.

Consider the matrix $((x_i|x_j))_{i,j} \in M_n(B)$. Given $(a_i)_{i=1}^n$ in B ,

$$\sum_{i,j} a_i^*(x_i|x_j)a_j = \left(\sum_i x_i \cdot a_i \middle| \sum_j x_j \cdot a_j \right) \geq 0,$$

so by [10, Lemma IV.3.2], the matrix is positive in $M_n(B)$. Let $R \in M_n(B)$ be the positive square-root, so $(x_i|x_j) = (R^*R)_{i,j} = \sum_k (R_{i,k})^* R_{k,j}$. Thus

$$T^*T = \sum_{i,j,k} \theta_{y_i \cdot R_{i,k}^* R_{k,j}, y_j} = \sum_{i,j,k} \theta_{y_i \cdot R_{i,k}^* R_{k,j} \cdot R_{k,j}^*} = \sum_k \theta_{z_k, z_k},$$

here setting $z_k = \sum_i y_i \cdot R_{k,i}^*$. □

Again, I am not aware of explicit reference for the following. The idea of this proof is motivated by Lance’s book [6].

Proposition A.2. *Let $T \in \mathcal{L}(Y_C, Y'_C)$ commute with the left action of B on ${}_BY_C$ and ${}_BY'_C$. Then $\text{id}_X \otimes T \in \mathcal{L}({}_AX \otimes_B Y_C, {}_AX \otimes_B Y'_C)$ with adjoint $\text{id}_X \otimes T^*$.*

Proof. Let $x_1, \dots, x_n \in X$ and $y_1, \dots, y_n \in Y$, and let $x = ((x_i|x_j))_{i,j} \in M_n(B)$. By [6, Lemma 4.2], x is positive; let $b \in M_n(B)$ be a square-root. Consider $u = \sum_i x_i \otimes y_i \in X \otimes_B Y$, so

$$\begin{aligned} \|(\text{id}_X \otimes T)u\|^2 &= \left\| \sum_{i,j} (x_i \otimes T y_i | x_j \otimes T y_j) \right\| = \left\| \sum_{i,j} (T y_i | (x_i|x_j) \cdot T y_j) \right\| \\ &= \left\| \sum_{i,j,k} (T y_i | b_{k,i}^* b_{k,j} \cdot T y_j) \right\| = \left\| \sum_{i,j,k} (T(b_{k,i} \cdot y_i) | T(b_{k,j} \cdot y_j)) \right\| \end{aligned}$$

here using that T commutes with the right B -action. This is of the form $(Tv|Tv)$ and by [6, Proposition 1.2] we have that $(Tv|Tv) \leq \|T\|^2(v|v)$. Hence

$$\|(\text{id}_X \otimes T)u\|^2 \leq \|T\|^2 \left\| \sum_{i,j,k} (b_{k,i} \cdot y_i | b_{k,j} \cdot y_j) \right\| = \|T\|^2 \|u\|^2,$$

by reversing the previous calculation. Hence $\text{id}_X \otimes T$ is bounded, and a routine calculation shows that it is adjointable, with adjoint $\text{id}_X \otimes T^*$. \square

A.1 Contragradient modules

Given ${}_A X_B$ we let \bar{X} be the contragradient vector space, which becomes a B - A -bimodule for actions

$$b \cdot \bar{x} \cdot a = \overline{a^* \cdot x \cdot b^*} \quad (a \in A, b \in B, x \in X).$$

We define the inner-products by

$${}_B(\bar{x}|\bar{y}) = (x|y)_B, \quad (\bar{x}|\bar{y})_A = {}_A(x|y) \quad (x, y \in A).$$

Notice that this definition correctly makes, for example, the right B -valued inner-product conjugate linear in the 1st variable.

Clearly if ${}_A X_B$ is bi-Hilbertian then so is ${}_B \bar{X}_A$, as the norms will still be equivalent.

We consider finite-rank operators on \bar{X} :

$$\theta_{\bar{x}, \bar{y}}: \bar{z} \mapsto \bar{x} \cdot (\bar{y}|\bar{z})_A = \bar{x} \cdot {}_A(y|z) = \overline{{}_A(y|z)^* \cdot x} = \overline{{}_A(z|y)} \cdot x. \quad (6)$$

References

- [1] Lawrence G. Brown, James A. Mingo, and Nien-Tsu Shen. Quasi-multipliers and embeddings of Hilbert C^* -bimodules. *Canad. J. Math.*, 46(6):1150–1174, 1994.
- [2] Michael Frank and Eberhard Kirchberg. On conditional expectations of finite index. *J. Operator Theory*, 40(1):87–111, 1998.
- [3] Tsuyoshi Kajiwara, Claudia Pinzari, and Yasuo Watatani. Ideal structure and simplicity of the C^* -algebras generated by Hilbert bimodules. *J. Funct. Anal.*, 159(2):295–322, 1998.
- [4] Tsuyoshi Kajiwara, Claudia Pinzari, and Yasuo Watatani. Jones index theory for Hilbert C^* -bimodules and its equivalence with conjugation theory. *J. Funct. Anal.*, 215(1):1–49, 2004.
- [5] Tsuyoshi Kajiwara and Yasuo Watatani. Jones index theory by Hilbert C^* -bimodules and K -theory. *Trans. Amer. Math. Soc.*, 352(8):3429–3472, 2000.
- [6] E. C. Lance. *Hilbert C^* -modules*, volume 210 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1995. A toolkit for operator algebraists.
- [7] R. Longo and J. E. Roberts. A theory of dimension. *K-Theory*, 11(2):103–159, 1997.
- [8] V. M. Manuilov and E. V. Troitsky. *Hilbert C^* -modules*, volume 226 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 2005. Translated from the 2001 Russian original by the authors.
- [9] Marc A. Rieffel. Induced representations of C^* -algebras. *Advances in Math.*, 13:176–257, 1974.
- [10] M. Takesaki. *Theory of operator algebras. I*, volume 124 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2002. Reprint of the first (1979) edition, Operator Algebras and Non-commutative Geometry, 5.
- [11] Yasuo Watatani. Index for C^* -subalgebras. *Mem. Amer. Math. Soc.*, 83(424):vi+117, 1990.