

COMP11120 Notes

Todd Davies

December 28, 2013

1 Discrete Structures

1.1 Terminology

A *structure* consists of certain *sets*. It also contains *elements* of these sets, *operations* on these sets and *relations* on these sets.

1.2 Number systems to learn

The following number must be learnt:

- \mathbb{N} The set of natural numbers (all whole numbers from 0 to ∞)
- \mathbb{Z} The set of integers (all whole numbers from $-\infty$ to ∞)
- \mathbb{Q} The set of rational numbers (any integer divided by any other integer e.g. $\frac{5}{4} = 1.25$)
- \mathbb{R} The set of real numbers (all finite and infinite decimal numbers)

1.2.1 Operations

Each number system has a set of valid operations that can be performed on elements in that system. Number systems only contain operations that will produce an output that is still within the number system.

For example, the number system \mathbb{N} contains the operations of addition and multiplication. This is because the summation of any two positive integers will *always*

be a member of \mathbb{N} , and the same goes for multiplication.

However, you may be wondering why subtraction and division aren't included in this number system. This is because for some numbers, the result of subtraction or division won't be inside the set \mathbb{N} . An example of this would be subtracting 4 from 2. Even though both of the operands are inside \mathbb{N} , the answer isn't.

Different sets may have different operations available. For example, you can concentrate any two members of the set *String* and end up with another *String*.

Types of operation Operations that have an operand either side of them are called *infix* operations. An example of an infix operation is addition. Infix operations are also referred to as *binary* operations since they have *two* operands.

Commutativity If an operation is commutative, the order of the operands doesn't matter. For example, addition is commutative since:

$$a + b = b + a \quad (1)$$

Subtraction however, isn't commutative:

$$a - b \neq b - a \quad (2)$$

Associativity An operation is associative if inserting or changing brackets doesn't change the outcome of the operation. For example, multiplication is associative since:

$$(a \times b) \times c = a \times (b \times c) \quad (3)$$

An operation may only be commutative or associative if it is commutative or associative for *all* elements of the set that the operation supports.

Distinguished elements A set may contain distinguished elements that have strange effects on certain operations in the set. An example is the number 1.

If we multiply *something* by 1, then the result will always be the same *something*. The same goes for 0 with addition. Because of this, we refer to 1 and 0 as distinguished elements of the set \mathbb{Z} .

1.2.2 Relations

Each of the sets $\mathbb{N}, \mathbb{Q}, \mathbb{Q}, \mathbb{R}$ carries binary comparison relations \leq and $<$. Different sets (such as *String*) may have other relations, such as:

- Is a section of
- Is an initial section of
- Occurs in

All relations return values in the set *Bool*.

1.3 Bases

Conventionally, we count using base 10. Base 10 includes, you guessed it, ten different symbols from 0 through to 9.

Sometimes however, it is convenient to count using different bases. Popular bases include:

Base n	Member symbols	Name
$n = 2$	$\mathbb{Z}_2 = \{0, 1\}$	Binary
$n = 8$	$\mathbb{Z}_8 = \{0, 1, 2, 3, 4, 5, 6, 7\}$	Octal
$n = 10$	$\mathbb{Z}_{10} = \{0, 1, \dots, 9\}$	Decimal
$n = 16$	$\mathbb{Z}_{16} = \{0 - 9, A - F\}$	Hexadecimal

1.3.1 How to read numbers in any given base

The formula for reading a number in a given base is as follows:

$$\sum_{i=0}^k a_i b^i \quad (4)$$

Where the number you're trying to read takes the form $a_k, a_{k-1}, \dots, a_2, a_1, a_0$ and b is the base you're using.

Example 1 Lets apply the formula to the base 10 number 27385:

$$\begin{aligned} 27385 &= (5 \times 10^0) + (8 \times 10^1) + (3 \times 10^2) \\ &\quad + (7 \times 10^3) + (2 \times 10^4) \\ &= (5 \times 1) + (8 \times 10) + (3 \times 100) \\ &\quad + (7 \times 1000) + (2 \times 10000) \\ &= 5 + 80 + 300 + 7000 + 20000 \\ &= 27385 \end{aligned} \tag{5}$$

Example 2 Lets apply the formula to the base 16 number $F00BA4$:

$$\begin{aligned} F00BA4 &= (4 \times 16^0) + (A \times 16^1) + (B \times 16^2) \\ &\quad + (0 \times 16^3) + (0 \times 16^4) + (F \times 16^5) \\ &= (4 \times 16^0) + (10 \times 16) + (11 \times 256) \\ &\quad + (0 \times 4096) + (0 \times 65536) + (15 \times 1048576) \\ &= 4 + 160 + 2816 + 0 + 0 + 15728640 \\ &= 15731620 \end{aligned} \tag{6}$$

1.3.2 Changing from base 10 to base n

In order to change into base n from base 10, we just repeatedly divide by n and use the remainder as the value for base n . Here are a few examples:

Example 1 Convert 893 into base 2.

$$\begin{array}{rcl} 893 \div 2 & = & 446 \text{ r}1 \\ 446 \div 2 & = & 223 \text{ r}0 \\ 223 \div 2 & = & 111 \text{ r}1 \\ 111 \div 2 & = & 55 \text{ r}1 \\ 55 \div 2 & = & 27 \text{ r}1 \\ 27 \div 2 & = & 13 \text{ r}1 \\ 13 \div 2 & = & 6 \text{ r}1 \\ 6 \div 2 & = & 3 \text{ r}0 \\ 3 \div 2 & = & 1 \text{ r}1 \\ 1 \div 2 & = & 0 \text{ r}1 \end{array}$$

Reading up from the bottom, we can see that the binary (base 2) representation is 1101111101.

Example 2 Convert 893 into base 9.

$$\begin{array}{rcll}
893 \div 9 & = & 99 & \text{r2} \\
99 \div 9 & = & 11 & \text{r0} \\
11 \div 9 & = & 1 & \text{r2} \\
1 \div 9 & = & 0 & \text{r1}
\end{array}$$

Reading up from the bottom, we can see that the nonal (base 9) representation is 1202.

Example 2 Convert 893 into base 16.

$$\begin{array}{rcll}
893 \div 16 & = & 55 & \text{r13} \\
55 \div 16 & = & 3 & \text{r7} \\
3 \div 16 & = & 0 & \text{r3}
\end{array}$$

Reading up from the bottom, we can see that the hexadecimal (base 16) representation is 3, 7, 13 or 37D.

1.4 A structure for the integers

The set \mathbb{Z} of all integers $\dots, -3, -2, -1, 0, 1, 2, 3, \dots$ includes the subset \mathbb{N} of all natural numbers together with the negative integers. We will be using three basic binary operations on the carrier set \mathbb{Z} : addition, multiplication and subtraction.

The standard notation for these operations is $+$, \times and $-$ and they are used as infix operations.

Sometimes different notations are used, for example, in programming $*$ is usually used, or there is a convention to write xy as an abbreviation for $x \times y$.

Both $+$ and \times are **commutative** but $-$ is not.

1. For all integers x, y both, $x + y = y + x$ and $xy = yx$
2. There are integers x, y with $x - y \neq y - x$.

Commutativity ensures that for most purposes the order in which the two arguments are consumed is irrelevant.

Both $+$ and \times are **associative** but $-$ is not.

1. For all integers x, y, z both $(x + y) + z = x + (y + z)$, $(xy)z = x(yz)$.
2. There are integers x, y, z with $(x - y) - z \neq x - (y - z)$.

Associativity ensures that for most purposes brackets are not needed to punctuate an expression. For in-

stance, we can make sense of $x + y + z$ and xyz since it doesn't matter where the brackets are, the resulting values are the same.

Below are the definitions of commutativity and associativity.

1. \otimes is **commutative** if and only if

$$a1 \otimes a2 = a2 \otimes a1 \quad (7)$$

for all $a1, a2, \in, A$

2. \otimes is **associative** if and only if

$$(a1 \otimes a2) \otimes a3 = a1 \otimes (a2 \otimes a3) \quad (8)$$

for all $a1, a2, a3, \in, A$

Some numbers have special effects on operations in a set. Some examples are 0 and 1. When 0 is added to a number, the result is still that number, and when multiplying a number by 1, the result is still that number. This means for the set \mathbb{Z} , 0 and 1 are neutral elements for the operations addition and multiplication respectively.

1.5 The formal language

We use the formal language to talk about \mathbb{Z} . Expressions in the formal language are built up from atoms in a recursive fashion, so an expression may be $(x \circ y)$ where x and y are expressions and \circ is one of the three operations $(+, -, \times)$. The brackets are important in the formal language, since they ensure that strings can only be parsed in one way.

Strings that contain literals that aren't neutral elements or have invalid bracketing are not valid in the formal language. Examples may include:

$$(x + y + z) \tag{9}$$

$$(x + 2) \tag{10}$$

$$x((y \tag{11}$$

We can use a parse tree to show how an expression is built up from its identifiers ($a, b, c \dots$), constants ($0, 1 \dots$) and operations ($+, -, \times \dots$).

For example, the parse tree of $(x \times (y + z))$ is:

$$\frac{x \quad \frac{y \quad z}{(y + z)} (+)}{(x \times (y + z))} (\times) \quad (12)$$

Often, especially with large parse trees, it's a pain to write so many identifiers. Because of this, it is a convention to replace identifiers with dots after they've been used once, like so:

$$\frac{x \quad \frac{y \quad z}{\cdot} (+)}{\cdot} (\times) \quad (13)$$

In order to parse a parse tree, we must assign an appropriate value to each of the 'leaves' of the tree (i.e. all the identifiers) and let the values trickle down towards the root node of the tree where the evaluated answer will appear.

Lets use the above example again. Let $x = 4$, $y = 6$ and $z = 2$:

$$\frac{4}{32} \frac{\frac{6}{8} 2}{(\times)} (+) \quad (14)$$

At this point, parse trees may seem very pointless, but this is because we're doing very simple arithmetic. However, when we start to define other operators that do unfamiliar things or don't use the *infix* notation, then using parse trees can be a big help!

1.6 The properties of sets

All sets have some properties in common that we can use to manipulate them. Examples include membership, equality, inclusion etc.

1.6.1 Set membership

To indicate that an element is a member of a set, we use the \in symbol. For example, to say that the element

true is a member of the set \mathbb{Bool} , we would write:

$$true \in \mathbb{Bool} \tag{15}$$

Interestingly, we can parse this into English in many ways though. It could mean any of the following things:

- *true* is an element of the set \mathbb{Bool}
- *true* is an member of the set \mathbb{Bool}
- *true* is contained in \mathbb{Bool}
- \mathbb{Bool} contains *true*

Conversely, to indicate that an element is not a member of a set, we use the symbol \notin :

$$sheep \notin \mathbb{Bool} \tag{16}$$

Note that there is no concept of *order* or *repetition* in sets. This means that the following sets are all equal:

$$\{1, 2, 3\} \tag{17}$$

$$\{2, 3, 1\} \tag{18}$$

$$\{2, 3, 1, 2\} \tag{19}$$

$$\{3, 3, 3, 3, 3, 2, 1\} \tag{20}$$

1.6.2 Set equality

Sets are equal if they have exactly the same members. Note that as far as sets are concerned, duplicate members are treated as just one member.

The notation for set equality is very easy. To say the set X is equal to the set Y , we write:

$$X = Y \tag{21}$$

However, you must understand that this is only true if for each element a in X that element will also be contained in Y and for each element a in Y , that element will also be contained within X :

$$\begin{aligned} \text{For each } a, a \\ \in X \\ \leftrightarrow a \\ \in Y \end{aligned} \tag{22}$$

1.6.3 Set inclusion

If one set is a subset of another set, all the members of the first set are also found within the second set. In mathematical terms:

$$\text{For each } a, a \in X \rightarrow a \in Y \tag{23}$$

The notation for inclusion is \subseteq , so in the above example, we would write:

$$X \subseteq Y \tag{24}$$

1.6.4 The empty set

The empty set is a set that contains no members at all. It's symbol is \emptyset .

Because the empty set has no members, it is a subset of all other sets:

$$\emptyset \in A \tag{25}$$

1.6.5 Singleton sets

For any entity a , we can form a set consisting only of a :

$$\{a\} \tag{26}$$

Be aware, a singleton set is not the same as the element contained within the set:

$$a \neq \{a\} \tag{27}$$

1.6.6 Set union

There are several ways of combining multiple sets together to create another set. One such method is set union, the symbol of which is \cup . The union of two sets is a set containing all the members of *both* sets. For example:

$$A = \{1, 3, 5, 7, 9\} \quad (28)$$

$$B = \{1, 1, 2, 3, 5, 8, 13\} \quad (29)$$

$$A \cup B = \{1, 2, 3, 5, 7, 8, 9, 13\} \quad (30)$$

We could also define the union of two sets mathematically, like so:

$$x \in A \cup B \leftrightarrow x \in A \text{ or } x \in B \quad (31)$$

1.6.7 Set intersection

Another way of combining sets is intersection. Intersecting two sets will produce a set of elements that belong to both of the sets. The symbol for intersection is \cap .

If we defined intersection mathematically, we would do so like this:

$$x \in A \cap B \leftrightarrow x \in A \text{ and } x \in B \quad (32)$$

1.6.8 Relative complement

The relative complement of two sets is all the members in the first set that aren't members of the second set. The symbol for the relative complement is $-$. Defined mathematically, we get:

$$x \in A - B \leftrightarrow x \in A \text{ and } x \notin B \quad (33)$$

1.6.9 The universal set

The universal set contains *all of the possible elements*. The notation we use to describe the universal set is S . We can define the De Morgan's laws using the universal set:

$$X' = S - X \quad (34)$$

1.7 Boolean algebra of sets

If we have a universal set, S , and consider only subsets of S . Then these are all the things that we have:

- Two distinguished subsets \emptyset and S
- A unary operation $(.)'$ on such subsets
- Two binary operations; \cap and \cup on such subsets

This structure of operations and sets is known as **Boolean algebra**. Boolean algebra has many basic properties that we can exploit in order to manipulate expression into other forms. Find them on the flashcards that go with these notes.

2 Propositional Logic

2.1 The logical connectives

Most of these are very similar to the properties of sets that we came across in the boolean algebra section. Learn these:

Name	Symbol	Meaning
Negation	\neg	not
Conjunction	\wedge	and
Disjunction	\vee	or
Implication	\implies	implies
Bi-implication	\iff	iff

All of these symbols are used as infixes, and so go in between two parameters. An exception to this however, is negation, which is a unary symbol, and is placed as a prefix before it's argument.

2.1.1 The truth tables of the logical connectives

Using a truth table, we can fully describe the behaviour of the connectives we have just described in the previous section:

p_1	p_2	$\neg p_1$	$p_1 \wedge p_2$	$p_1 \vee p_2$
T	T	F	T	T
T	F	F	F	T
F	T	T	F	T
F	F	T	F	F

p_1	p_2	$p_1 \implies p_2$	$p_1 \iff p_2$
T	T	T	T
T	F	F	F
F	T	T	F
F	F	T	T

A note about implication: In order to understand why the truth table for implication is as stated above, consider this example:

$$x > 3 \text{ implies } x > 1 \quad (35)$$

Now take a look at the truth table for this expression:

x	$x > 3$	$x > 1$	$x > 3 \implies x > 1$
4	T	T	T
2	F	T	T
0	F	F	T

As you can see, there is no way to make the first expression ($x > 3$) true, but the second expression ($x > 1$) false. Henceforth, the first expression implies the second expression.

2.2 The formulae of propositional logic

In order to build up a piece of propositional logic, we must construct an expression from *atomic formulae* and connectives.

There are three rules we can use to generate a formulae in PL:

1. Every atomic formula is a formula of PL
2. If A is a formula, then so is $\neg A$

3. If A, B are formulae, then so are $(A \wedge B)$, $(A \vee B)$,
 $(A \implies B)$, $(A \iff B)$

Syntactic conventions are of course in widespread use.

For example, when you have written an expression, it is usual to leave off the outermost parentheses, so $(p \wedge (r \vee q))$ would become $p \wedge (r \vee q)$.

It is also common to leave out parentheses from repeated uses of conjunction or disjunction, so $p \wedge (q \wedge r)$ would become $p \wedge q \wedge r$.

2.3 Truth valuations

A truth valuation is a list of allocations that define the values of variables in an expression, where:

$$\{p_1 = x_1, \dots, p_n = x_n\} \tag{36}$$

In order to determine the end value of the PL formula, you must first construct a truth valuation of it's

component atoms, and only then can the formula be evaluated.

2.4 Tautologies and contradictions

If the outcome of a formula is always T for every possible truth valuation, then the formula is said to be a **tautology**. An example of such an expression is $p \vee \neg p$.

In contrast, a formula that will yield an outcome of F for every possible truth valuation is named a **contradiction**. Such an example may be $p \wedge \neg p$.

If an expression is not a tautology and isn't a contradiction either, then it is named **satisfiable**.

3 Probability

3.1 Sample spaces and events

A **Sample Space** is a set of all the possible outcomes of an *experiment*. In this sense, an experiment is taken to be some occurrence that will produce one of many outcomes.

The notation we use for the sample space is Ω .

$$\Omega = \{\omega_1, \omega_2, \dots, \omega_n\} \quad (37)$$

An **event** is a set containing one or more outcomes of an experiment. Since all outcomes are included in the set Ω , all events are also subsets of the sample space. This works both ways, any subset of Ω is an event. We use ω to represent an event.

An event is said to have occurred if the outcome of an experiment contains an element of that event.

Since all events are sets, it makes sense that we would have an empty set and a universal set. As with dis-

crete logic (see page), we also use the empty set in probability. Here, we use it to describe an event that *never* occurs. The notation for the empty set is \emptyset . We have already discussed the universal set (see page) of probability, it's the sample space, Ω .

To sum up events an sample spaces, here's an example. If toss a coin, the following events are generated:

Event	Description
\emptyset	Nothing happens. This will never occur.
$\{H\}$	The coin shows heads.
$\{T\}$	The coin shows tails.
Ω	The coin shows either heads or tails. This event always occurs.

3.2 More set properties

All of the set operations we discussed in the Discrete Structures section of the notes still apply here, these can be found in section 1.6 on page .

However, there are three additional properties that will

be useful in this section.

3.2.1 The number of items in a set

We can use the operator $\#$ to find the number of elements in a set. If we have an event A that contains the elements $\{\omega_1, \omega_2, \omega_3\}$ then the number of outcomes it contains is three, therefore $\#A = 3$.

3.2.2 Set difference

If we have two sets A and B , and B is a subset of A (i.e. $B \subset A$), then we can find the set difference between A and B .

The set difference is defined as **all the elements in A that are not in B** . The symbol for set difference is “ \setminus ”.

Mathematically, this is:

$$A \setminus B = \{\omega \in \Omega \mid \omega \in A \text{ and } \omega \notin B\} \quad (38)$$

3.2.3 Disjoint sets

If two sets are disjoint, then they share no common elements, that is to say that:

$$A \cap B = \emptyset \quad (39)$$

This is important in probability, since if two events are disjoint, then they will never occur at the same time.

4 Probability measures

A probability measure is a mapping between an event and the probability that the event will occur.

$$\begin{aligned} \mathbb{P} : \{ \textit{Collection of all events} \} & \quad (40) \\ & \rightarrow [0, 1] \end{aligned}$$

This defines \mathbb{P} as a number between 0 and 1 (i.e. a probability). Since a probability measure is defined

for *all* events that can occur, then the following are true:

- $\mathbb{P}(\Omega) = 1$
- If A and B are disjoint events then $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$
- If $\mathbb{P}(A) = 0$ then A will never occur.
- If $\mathbb{P}(A) = 1$ then A will always occur.

In order to verify whether any given mapping is a probability measure, you must check it against the first two of the above bullets; that the sum of all the mappings is 1 and the mapping of the union of any two disjoint events is the same as the sum of the separate mappings of the disjoint events.

4.1 Indicator functions

An indicator function will yield a value of 1 if an element ω is inside a set, and a value of 0 if the element is not inside the set. It can be defined as follows:

$$1_A(\omega_i) = \begin{cases} 1 & \text{if } \omega_i \in A \\ 0 & \text{if } \omega_i \notin A \end{cases} \quad (41)$$

We can therefore define the probability of an event A to be the sum of all the probabilities of the events inside A :

$$\mathbb{P}(A) = \sum_{i=1}^n 1_A(\omega_i) p_i \quad (42)$$

5 Computing the probability of equally likely functions

If we have a sample space $\Omega = \{\omega_1, \omega_2, \dots, \omega_n\}$ where each element is equally likely to occur, then we can say that $p_i = \mathbb{P}(\{\omega_i\})$. Since all the elements are equally likely to occur, we can say that $p_1 = p_2 = \dots = p_i$ and also that $p_1 + p_2 + \dots + p_i = 1$.

Using ideas from the previous section, we can find the probability of any event A inside a sample space where

all elements are equally likely. To do this, all we need to do is find the sum of all the elements in the sample space, but multiply the elements by the indicator function of A before we add them onto the sum:

$$\mathbb{P}(A) = \sum_{i=1}^n 1_A(\omega_i) p_i \quad (43)$$

However, since all the elements inside Ω are equally likely, that must mean that for any event ω_i , it's probability must be $\frac{1}{n}$ where n is the number of elements inside the sample space. If we take this into account, we can formulate the following formula:

$$\mathbb{P}(A) = \frac{1}{n} \sum_{i=1}^n 1_A(\omega_i) \quad (44)$$

However, we can make this even more simple; if we recognise that any element in A has the same probability of occurring as any element in Ω , then it's easy to realise that the probability of A occurring will be

equal to the relationship between sizes of the sets A and Ω :

$$\mathbb{P}(A) = \frac{\#A}{\#\Omega} \quad (45)$$

5.1 Finding the binomial coefficient

The binomial coefficient is represented as $\binom{n}{k}$, both n and k are integers. It is pronounced *n choose k*.

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} \quad (46)$$

5.2 Choosing items

If we have a set of three items a, b and c , then how many different ways can we choose two of those three items from the set?

This depends on two things. Does the order of the items matter, and can we choose an item twice?

If the order is important and we can choose an item many times, then we can choose the following items:

$$(a, a), (a, b), (a, c), (b, a), (b, b), (b, c), (c, a), (c, b), (c, c) \quad (47)$$

If the order is not important, then we have:

$$(a, a), (a, b), (a, c), (b, b), (b, c), (c, c) \quad (48)$$

If the order is important, but we can't choose an item twice, we get:

$$(a, b), (a, c), (b, a), (b, c), (c, a), (c, b) \quad (49)$$

Finally, if the order is not important and the items cannot be selected twice, then:

$$(a, b), (a, c), (b, c) \quad (50)$$

In order to calculate the number of possible different selections we can make in these different cases, we can use this table:

	Order important?	
	Yes	No
Without replacement	$\frac{n!}{(n-k)!}$	$\binom{n}{k}$
With replacement	n^k	Complicated

It's easy to verify that these are true; since:

- With replacement, order important:

$$\begin{aligned} n^k &= 3^2 \\ &= 9 \end{aligned} \tag{51}$$

- Without replacement, order important:

$$\begin{aligned} \frac{n!}{(n-k)!} &= \frac{3!}{(3-2)!} \\ &= 6 \end{aligned} \tag{52}$$

- Without replacement and order is not important:

$$\begin{aligned}
 \binom{n}{k} &= \frac{n!}{k(n-k)!} \\
 &= \frac{3!}{2(3-2)!} \\
 &= \frac{6}{2} \\
 &= 3
 \end{aligned}
 \tag{53}$$

6 Conditioning

If an experiment is performed where we already know that a certain event will occur, then we can factor in an additional factor into our probabilistic theories. This factor could be called a condition.

An example is throwing a dice. If we define the event $B = \{5, 6\}$ then there is a probability of $\frac{1}{3}$ that the event B will occur. If we know in advance that B will occur, then the probabilities that the events outside B will occur suddenly drop to zero:

$$\mathbb{P}(\{1\}) = \dots = \mathbb{P}(\{4\}) = 0 \quad (54)$$

The probability that B will occur is 1, therefore the probability that either event in B will occur is $\frac{1}{2}$.

To clarify the following two things have just happened:

- All outcomes not in the event B have a probability 0.
- All outcomes in the event B have their probability *scaled* by a factor of $\frac{1}{\mathbb{P}(B)}$.

In a more general case, the probability that any event A occurs, given that the event B occurs is:

$$\frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} \quad (55)$$

The notation we use for conditional probability (i.e. the probability an event A will occur given that the event definitely will B occur) is:

$$\mathbb{P}(A|B) \quad (56)$$

6.1 Independent events

If the probability of the event A does not change when we add the conditional that an event B will certainly occur and $\mathbb{P}(B) \geq 0$, then the events A and B are independent. This could be described as:

$$\mathbb{P}(A) = \mathbb{P}(A|B) \tag{57}$$

We can manipulate this mathematical definition of independence using the equations we described in the at the start of the conditionals section:

$$\begin{aligned} \mathbb{P}(A) = \mathbb{P}(A|B) &\iff \mathbb{P}(A) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} \\ &\iff \mathbb{P}(A)\mathbb{P}(B) = \mathbb{P}(A \cap B) \end{aligned} \tag{58}$$

Henceforth, we often take the last of these equations as our definition for independence.

Of course, it's easy to do the following too:

$$\begin{aligned}
\mathbb{P}(A)\mathbb{P}(B) &= \mathbb{P}(A \cap B) \\
&\iff \mathbb{P}(B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)} \\
&\iff \frac{\mathbb{P}(B \cap A)}{\mathbb{P}(A)} \\
&\iff \mathbb{P}(B|A)
\end{aligned} \tag{59}$$

$$\mathbb{P}(A|B) \iff \mathbb{P}(B|A) \tag{60}$$

So, an event is independent of another event if knowledge that one event occurs won't give you any extra information about how likely it is that the other event will occur. This doesn't mean that the events are disjoint (unless the probability of either was 0).

6.2 Partitions

Partitions are sections of a sample space that:

- Don't overlap.
- Include all elements inside the sample space between them.

So, suppose I had partitions $E_0 \dots E_n$, the following would hold:

- $E_i \cap E_j = \emptyset$ for and i and j
- $E_0 \cup E_1 \cup \dots \cup E_n = \Omega$

This means that all partitions are mutually disjoint.

Note that the probability of the union of any two partitions will be the sum of the probability of the separate partitions, since all partitions are disjoint.

6.3 The law of total probability

The law of total probability splits any event A up in terms of the partitions in the sample space:

$$A = (A \cap E_0) \cup (A \cap E_1) \cup \dots \cup (A \cap E_n) \quad (61)$$

We can write this in terms of probabilities:

$$\mathbb{P}(A) = \mathbb{P}(A|E_1)\mathbb{P}(E_1) + \cdots + \mathbb{P}(A|E_n)\mathbb{P}(E_n) \quad (62)$$

6.4 Bayes Theorem

The derivation of Bayes theorem is quite easy; start with an event partitioned into m partitions, then take $\mathbb{P}(E_i|A)$ and use first the definition of conditional probability, then the multiplicative law, then the law of total probability. The process is described mathematically below:

$$\begin{aligned} \mathbb{P}(E_i|A) &= \frac{\mathbb{P}(E_i \cap A)}{\mathbb{P}(A)} \\ &= \frac{\mathbb{P}(A|E_i)\mathbb{P}(E_i)}{\mathbb{P}(A)} \\ &= \frac{\mathbb{P}(A|E_i)\mathbb{P}(E_i)}{\mathbb{P}(A|E_1)\mathbb{P}(E_1) + \cdots + \mathbb{P}(A|E_m)\mathbb{P}(E_m)} \end{aligned} \quad (63)$$

Bayes theorem is good if you have the probability of some event A that depends on the condition of the event B happening (i.e. you know $\mathbb{P}(A|B)$, but you instead want to know $\mathbb{P}(B|A)$).

6.4.1 The complementary law for conditional probabilities

The complementary law states that:

$$\mathbb{P}(A|B) = 1 - \mathbb{P}(A^c|B) \quad (64)$$

7 Random variables

A random variable is a mapping between the sample space Ω to real numbers (\mathbb{R}). A random variable is usually denoted by X .

$$\begin{aligned} X : \Omega \\ \rightarrow \mathbb{R} \end{aligned} \quad (65)$$

This means that every outcome in the sample space corresponds to a real number. The range of the random variable is very intuitive; if we were throwing a dice, then the range would be $\{1, 2, \dots, 6\}$.

The sum of the probabilities of the random variable is always equal to 1:

$$\sum_{i=1}^m \mathbb{P}(X = r_i) = 1 \quad (66)$$

7.1 Constructing new random variables from existing ones

Creating a random variable from another random variable is fairly easy; it's basically a composite function. If we have a random variable X and a function f then:

$$Y(\omega) = f(X(\omega)) \text{ for all } \omega \in \Omega \quad (67)$$

Note that the range of the new function (Y) may be different from the range of the old one (X).

7.2 Probability mass functions

Usually abbreviated to *pmf* a probability mass function is a function (denoted by p) that contains information regarding the probabilities of X in given ranges.

Since the output of a *pmf* is a probability, the range of a *pmf* is always $[0, 1]$.

Mathematically, the probability mass function of a random variable X is defined as:

$$\begin{aligned} p(r_i) : & \\ &= \mathbb{P}(X \\ &= r_i) \text{ for all } i > 1 \end{aligned} \tag{68}$$

7.3 Cumulative distribution functions

A cumulative distribution function *cdf* is similar to a *pmf* except that:

- The value of the *cdf* is the sum of the *pmf* up to that point.

- A *cdf* can be generated for any function, not just discrete ones like a *pmf* can.

However, both the *cdf* and the *pmf* share the same range. The definition of *cdf* is:

$$F(x) = \mathbb{P}(X \leq x) \text{ for all } x \in \mathbb{R} \quad (69)$$

A graph of a *cdf* function contains all the information about the random variable it has mapped. We can deduce it's range, the *pmf* and where the graph jumps. Note that black dots on the graph indicate where a value is, and white dots at the same x coordinate show where the value isn't.