

Simulating Quantum Drude Oscillators on a photonic quantum computer

Matthieu Sarkis*

*Department of Physics and Materials Science
University of Luxembourg, L-1511,
Luxembourg City, Luxembourg.*

†

(Dated: March 9, 2023)

I. TO DO

- scatter plot (or linearly interpolated plot) with three different angles (optimal angle, one below, one above without binding)
- plot with the morse fit for the optimal angle
- plot of the quality of the morse fit as a function of the angle
- plot with the joint wigner function at different distances (from large distance to short distance)
- plot of the quadratures joint density at different distances
- plot of the entropy?
- the best angle is θ_{opt} [7]
- do the r^2 statistical test

II. INTRODUCTION

We work in atomic units, for which $m_e = q_e = \hbar = \frac{1}{4\pi\epsilon_0} = 1$.

Dispersion forces, also known as van der Waals (vdW) forces originate from the electromagnetic interaction between electrically neutral atoms or molecules which do not have permanent electric moments [1–4]. They are ever-present long-range forces between atoms or molecules arising from the zero-point fluctuations of the quantum electromagnetic field [5–9]. Their importance can be appreciated on both macro- and micro-scales; for instance, the first macroscopic signature of dispersion forces is the well-known correction to the equation of state of an ideal gas that led to the van der Waals equation [10]. Moreover, dispersion forces also influence the structure of liquids and solids such as the anomalies of water [11] as well as the macroscopic properties

of macromolecules such as their structure [12], stability [13, 14], dynamics [15–17], and electric [18] and optical [19] responses. The most natural framework for the investigation of the response of matter subjected to such forces is quantum electrodynamics [20–24]. However, as widely shown in the literature, the inclusion of vdW dispersion interactions can be done by means of many-body methods [25–32]. Dispersion vdW interactions are often represented within the Lennard-Jones approach, namely, through a pairwise two-body interatomic potential [33–38] of the form C_6/R^6 (where R is the interatomic distance and C_6 a system-dependent constant). Among all the existing models in the literature, the many-body dispersion (MBD) framework has been undoubtedly proved to be an accurate approach [37, 39]. In the MBD framework, the electronic response of valence electrons in atoms and molecules is supposed to be linear and this can be formally done through the introduction of the quantum Drude oscillator (QDO). A single QDO is coarse-grained quantum-mechanical model in which the properties of an atom are encompassed in a small number of parameters. The model consists in assimilating the atom to a point particle of mass m and electric charge $-q$ attached to a fixed (infinite mass) center of charge $+q$ by a harmonic spring characterized by a frequency ω . Molecules are then defined as a collection of QDOs in dipole-dipole interaction. For some specific choices of the matter system geometry, the quantum Hamiltonian can be exactly diagonalized. For instance, in a closed linear chain of molecules, one can analytically solve for the spectrum of the Hamiltonian [40]. In the case of a general geometry, instead, one can solve for the London-van der Waals interaction energy through a perturbative approach [41]. This simple model has been extensively used in various contexts, for instance in order to tackle the electronic structure problem for isolated molecules, in particular long range interactions, as well as to study the impact of an ambient bath or of an external electric field on molecular properties [42, 43]. Though simple, through a numerical treatment this system was shown to capture long-range phenomena in large biomolecular systems [44]. By construction, the MBD framework relies on the dipole approximation of the electron–electron Coulomb interaction leaving aside any contribution coming from high-order terms. In the literature, dispersion forces have been addressed mostly for atomic dimers and small

* matthieu.sarkis@uni.lu

†

systems, via multipolar generalizations of the pairwise second-order perturbative approaches [33, 34, 38, 45]. In this Letter ...

III. DEFINITION OF THE MODEL

A. Three-dimensional model

The Hamiltonian describing a system of N QDOs in 3d is given by:

$$H_{(3,3)} = \sum_{i=1}^N \left[\frac{\mathbf{p}_i^2}{2m_i} + \frac{1}{2} m_i \omega_i^2 \mathbf{x}_i^2 \right] + \sum_{i<j} V_{\text{Coul}}(\mathbf{x}_i, \mathbf{x}_j), \quad (1)$$

with the Coulomb interaction receiving contributions from every pair of constituents (centers and point particles):

$$\frac{V_{\text{Coul}}(\mathbf{x}_i, \mathbf{x}_j)}{q_i q_j} = \frac{1}{r_{ij}} - \frac{1}{|\mathbf{r}_{ij} + \mathbf{x}_i|} - \frac{1}{|\mathbf{r}_{ij} - \mathbf{x}_j|} + \frac{1}{|\mathbf{r}_{ij} - \mathbf{x}_j + \mathbf{x}_i|}$$

The subscript (3,3) in the above definition will become clear soon. In terms of components, the full Coulomb potential reads:

$$\begin{aligned} \frac{V_{\text{Coul}}(\mathbf{x}_i, \mathbf{x}_j)}{q_i q_j} &= \frac{1}{r_{ij}} - \frac{1}{\sqrt{r_{ij}^2 + x_i^2 + y_i^2 + z_i^2 + 2r_{iz}}} \\ &- \frac{1}{\sqrt{r_{ij}^2 + x_j^2 + y_j^2 + z_j^2 - 2r_{ij}z_j}} \\ &+ \frac{1}{\sqrt{r_{ij}^2 + (x_j - x_i)^2 + (y_j - y_i)^2 + (z_j - z_i)^2 - 2r_{ij}(z_j - z_i)}} \end{aligned} \quad (2)$$

In the multipolar expansion, this can be expressed as a power series in the inverse distance separating the two centers:

$$V_{\text{Coul}}(\mathbf{x}_i, \mathbf{x}_j) = \sum_{n \geq 0} V_n(\mathbf{x}_i, \mathbf{x}_j), \quad (3)$$

with the following scaling behavior in terms of the distance between the centers:

$$V_n(\mathbf{x}_i, \mathbf{x}_j) \propto r_{ij}^{-n-3}. \quad (4)$$

The potential V_0 corresponds then to the dipole-dipole interaction, V_1 to the dipole-quadrupole interaction, and V_2 to the quadrupole-quadrupole and dipole-octupole interaction. Let us give here the expression for the multi-

polar potential up to quartic order:

$$V_0(\mathbf{x}_i, \mathbf{x}_j) = q_i q_j \frac{x_i x_j + y_i y_j - z_i z_j}{r_{ij}^3} \quad (5)$$

$$\begin{aligned} V_1(\mathbf{x}_i, \mathbf{x}_j) &= \frac{q_i q_j}{2r_{ij}^4} \left(-3x_i^2 z_j - 6x_i x_j z_i + 6x_i x_j z_j + 3x_j^2 z_i \right. \\ &\quad \left. - 3y_i^2 z_j - 6y_i y_j z_i + 6y_i y_j z_j + 3y_j^2 z_i \right. \\ &\quad \left. + 6z_i^2 z_j - 6z_i z_j^2 \right) \end{aligned} \quad (6)$$

$$\begin{aligned} V_2(\mathbf{x}_i, \mathbf{x}_j) &= \frac{q_i q_j}{2r_{ij}^4} \left(-6x_i^3 x_j + 9x_i^2 x_j^2 - 6x_i^2 y_i y_j + 3x_i^2 y_j^2 \right. \\ &\quad \left. + 24x_i^2 z_i z_j - 12x_i^2 z_j^2 - 6x_i x_j^3 - 6x_i x_j z_i^2 \right. \\ &\quad \left. + 12x_i x_j y_i y_j - 6x_i x_j y_j^2 + 24x_i x_j z_i^2 \right. \\ &\quad \left. - 48x_i x_j z_i z_j + 24x_i x_j z_j^2 + 3x_j^2 y_i^2 - 6x_j^2 y_i y_j \right. \\ &\quad \left. - 12x_j^2 z_i^2 + 24x_j^2 z_i z_j - 6y_i^3 y_j + 9y_i^2 y_j^2 \right. \\ &\quad \left. + 24y_i^2 z_i z_j - 12y_i^2 z_j^2 - 6y_i y_j^3 + 24y_i y_j z_i^2 \right. \\ &\quad \left. - 48y_i y_j z_i z_j + 24y_i y_j z_j^2 - 12y_j^2 z_i^2 + 24y_j^2 z_i z_j \right. \\ &\quad \left. - 16z_i^3 z_j + 24z_i^2 z_j^2 - 16z_i z_j^3 \right) \end{aligned} \quad (7)$$

We define the dimensionless position and momenta associated to QDO i :

$$\mathbf{X}_i := \sqrt{\frac{m_i \omega_i}{\hbar}} \mathbf{x}_i, \quad \mathbf{P}_i := \frac{\mathbf{p}_i}{\sqrt{2\hbar m_i \omega_i}}, \quad (8)$$

in terms of which the $3N$ creation and annihilation operators read ($\alpha = 1, 2, 3$)

$$\mathbf{a}_i = \frac{\mathbf{X}_i + i\mathbf{P}_i}{\sqrt{2}}, \quad \mathbf{a}_i^\dagger = \frac{\mathbf{X}_i - i\mathbf{P}_i}{\sqrt{2}}. \quad (9)$$

In terms of the dimensionless canonical variables, the Hamiltonian reads

$$\begin{aligned} H_{(3,3)} &= \sum_{i=1}^N \frac{\hbar \omega_i}{2} (\mathbf{X}_i^2 + \mathbf{P}_i^2) \\ &+ \sum_{i<j} V_{\text{Coul}} \left(\sqrt{\frac{\hbar}{m_i \omega_i}} \mathbf{X}_i, \sqrt{\frac{\hbar}{m_j \omega_j}} \mathbf{X}_j \right) \end{aligned} \quad (10)$$

and can be rewritten

$$\begin{aligned} H_{(3,3)} &= \sum_{i=1}^N \frac{\hbar \omega_i}{2} \left(\mathbf{a}_i^\dagger \cdot \mathbf{a}_i + \frac{3}{2} \right) \\ &+ \sum_{i<j} V_{\text{Coul}} \left(\sqrt{\frac{\hbar}{m_i \omega_i}} \frac{\mathbf{a}_i + \mathbf{a}_i^\dagger}{\sqrt{2}}, \sqrt{\frac{\hbar}{m_j \omega_j}} \frac{\mathbf{a}_j + \mathbf{a}_j^\dagger}{\sqrt{2}} \right) \end{aligned} \quad (11)$$

B. One-dimensional case

Let us define one-dimensional instances of the QDO model as follows: we restrict the movement of the two electrons to be along a common axis (directed by a unit

vector \hat{e}_θ) which form an angle θ with respect to the vector \mathbf{r}_{ij} connecting the two nuclei. As particular cases, we obtain models in which the electrons are constrained to move either in the direction parallel to the axis separating the two nuclei ($\theta = 0$), or perpendicular to the latter ($\theta = \pi/2$). We therefore have a family of one-dimensional models which can be obtained from the full-fledged 3d model simply by setting to zero the contribution from the oscillator modes belonging to the plane perpendicular to \hat{e}_θ . Let us denote by x the remaining degree of freedom. As will be explained in the results section, one particular model will prove relevant, the so-called *space regularized model*, in which the angle θ is chosen to be infinitesimally small $\theta = \epsilon \ll 1$.

$$V_0(x_i, x_j) = q_i q_j \frac{x_i x_j}{r_{ij}^3} \times \begin{cases} 1 - 3 \cos^2 \theta, & \text{generic case} \\ -2, & \text{parallel case} \\ 1, & \text{perpendicular case} \\ -\frac{1}{2}, & \text{oblique case} \\ -2 - 6\epsilon, & \text{regularized case} \end{cases} \quad (12)$$

The next terms in the multipolar expansion are:

$$V_1(x_i, x_j) = q_i q_j \frac{x_i x_j (x_i - x_j)}{r_{ij}^4} \times \begin{cases} \frac{3 \cos \theta (-3 + 5 \cos^2 \theta)}{2}, & \text{generic case} \\ 3, & \text{parallel case} \\ 0, & \text{perpendicular case} \\ -\frac{3}{4\sqrt{2}}, & \text{oblique case} \\ 3 + 18\epsilon, & \text{regularized case} \end{cases} \quad (13)$$

$$V_2(x_i, x_j) = q_i q_j \frac{x_i x_j (2x_i^2 - 3x_i x_j + 2x_j^2)}{r_{ij}^5} \times \begin{cases} -\frac{3 - 30 \cos^2 \theta + 35 \cos^4 \theta}{4}, & \text{generic case} \\ -2, & \text{parallel case} \\ -\frac{3}{4}, & \text{perpendicular case} \\ \frac{13}{16}, & \text{oblique case} \\ -2 - 20\epsilon, & \text{regularized case} \end{cases} \quad (14)$$

For the full Coulomb potential, assuming that the electrons are constrained to move along an axis, we get the following expressions:

$$\frac{V_{\text{Coul}}^\perp(x_i, x_j)}{q_i q_j} = \frac{1}{r_{ij}} - \frac{1}{\sqrt{r_{ij}^2 + x_i^2}} - \frac{1}{\sqrt{r_{ij}^2 + x_j^2}} + \frac{1}{\sqrt{r_{ij}^2 + (x_j - x_i)^2}} \quad (15)$$

in the case where the electrons move perpendicular to the axis joining the two nuclei, and

$$\frac{V_{\text{Coul}}^\parallel(x_i, x_j)}{q_i q_j} = \frac{1}{r_{ij}} - \frac{1}{|r_{ij} + x_i|} - \frac{1}{|r_{ij} - x_j|} + \frac{1}{|r_{ij} + x_i - x_j|}$$

in the case where they move parallel to the latter. In the case of a generic angle θ between \mathbf{r}_{ij} and $\mathbf{x}_i/\mathbf{x}_j$, one has

$$\frac{V_{\text{Coul}}^\theta(x_i, x_j)}{q_i q_j} = \frac{1}{r_{ij}} - \frac{1}{\sqrt{r_{ij}^2 + 2r_{ij}(\cos \theta)x_i + x_i^2}} - \frac{1}{\sqrt{r_{ij}^2 - 2r_{ij}(\cos \theta)x_j + x_j^2}} + \frac{1}{\sqrt{r_{ij}^2 - 2r_{ij}(\cos \theta)(x_j - x_i) + (x_j - x_i)^2}}. \quad (16)$$

The space regularized model Coulomb potential reads

$$\frac{V_{\text{Coul}}^\epsilon(x_i, x_j)}{q_i q_j} = \frac{1}{r_{ij}} - \frac{1}{\sqrt{r_{ij}^2 + 2r_{ij}(1 + \epsilon)x_i + x_i^2}} - \frac{1}{\sqrt{r_{ij}^2 - 2r_{ij}(1 + \epsilon)x_j + x_j^2}} + \frac{1}{\sqrt{r_{ij}^2 - 2r_{ij}(1 + \epsilon)(x_j - x_i) + (x_j - x_i)^2}}. \quad (17)$$

C. 12 different models

For the simulation, we will restrict the system to two QDOs. Depending on the space dimensionality (1d oblique, 1d parallel, 1d perpendicular, 3d) and the choice of potential (quadratic, quartic, Coulomb), we therefore have 12 models to try and study, gathered in table (I).

TABLE I. The twelve models

POT \ DIM	1d oblique	1d parallel	1d perpendicular	3d
QUAD	$H_{(1,0)}$	$H_{(1,1)}$	$H_{(1,2)}$	$H_{(1,3)}$
QUART	$H_{(2,0)}$	$H_{(2,1)}$	$H_{(2,2)}$	$H_{(2,3)}$
COULOMB	$H_{(3,0)}$	$H_{(3,1)}$	$H_{(3,2)}$	$H_{(3,3)}$

IV. PHOTONIC CIRCUIT

The circuit implements a unitary $U(\theta)$ acting on an input reference state (the Fock vacuum for instance) that we simply take to be the vacuum state $|0\rangle$. The state prepared by the circuit is therefore given by

$$|\psi(\theta)\rangle = U(\theta)|0\rangle. \quad (18)$$

In the dipolar approximation, namely for the Hamiltonians $H_{(1,1)}$, $H_{(1,2)}$ and $H_{(1,3)}$, we expect that using a Gaussian state would be enough. The circuit is therefore composed of at most quadratic optical components (squeezing operations for our ansatz). However for the other models, non-Gaussian operation should be added in the end of each layer in the ansatz circuit.

Once the ansatz state $|\psi(\theta)\rangle$ has been produced, one should extract the value of the energy in that state, namely compute the value of

$$\langle\psi(\theta)|H|\psi(\theta)\rangle \quad (19)$$

To be specific, let us take the model $H_{(3,3)}$, and let us denote by angular brackets the expectation in state $|\psi(\theta)\rangle$. One has

$$\begin{aligned} \langle H_{(3,3)} \rangle &= \sum_{i=1}^N \frac{\hbar\omega_i}{2} \left(\langle n_{i,x} \rangle + \langle n_{i,y} \rangle + \langle n_{i,z} \rangle + \frac{3}{2} \right) \\ &+ \sum_{i < j} \left\langle V_{\text{Coul}} \left(\sqrt{\frac{\hbar}{m_i\omega_i}} \mathbf{X}_i, \sqrt{\frac{\hbar}{m_j\omega_j}} \mathbf{X}_j \right) \right\rangle \end{aligned} \quad (20)$$

by linearity of the expectation. On the second line one has to compute something of the form $\langle f(X_{i,\alpha}) \rangle$, where $(X_{i,\alpha})$ denotes collectively the position quadrature of all the photon channels (including QDO and spatial index). One therefore needs to extract the statistics of quadratures by preparing and measuring the state $|\psi(\theta)\rangle$ in the quadrature basis. Once the joint density ρ of $(X_{i,\alpha})$ in the state $|\psi(\theta)\rangle$ is known, one can compute

$$\langle f(X_{i,\alpha}) \rangle = \int_{\mathbb{R}^{3N}} f(x_{i,\alpha}) \rho(x_{i,\alpha}) \prod_{i,\alpha} dx_{i,\alpha}. \quad (21)$$

There is a complication related to the fact that strawberryfields does not allow to access the joint law of the position quadratures. We therefore have to extract it by ourselves. Let us first suppose that there are K photon modes. The statevector is represented in the Fock basis as follows:

$$|\psi\rangle = \sum_{n_1, \dots, n_{2K}=0}^{\infty} \alpha_{n_1 \dots n_{2K}} |n_1\rangle \otimes \dots \otimes |n_{2K}\rangle. \quad (22)$$

The modes labeled by (n_1, \dots, n_K) correspond to QDO₁, while the modes (n_{K+1}, \dots, n_{2K}) are attached to QDO₂. The amplitude of a specific tuple of the quadratures (X_1, \dots, X_K) is therefore given by:

$$\langle X_1, \dots, X_{2K} | \psi \rangle = \sum_{n_1, \dots, n_{2K}=0}^{\infty} \alpha_{n_1 \dots n_{2K}} \prod_{i=1}^{2K} \frac{e^{-\sum_{i=1}^{2K} \frac{X_i^2}{2}} H_{n_i}(X_i)}{\sqrt{\pi^{1/2} 2^{n_i} n_i!}},$$

in terms of the Hermite polynomials. The joint law of the quadratures in the state $|\psi\rangle$ is therefore given by

$$\begin{aligned} \rho(X_1, \dots, X_{2K}) &= \sum_{\substack{n_1, \dots, n_{2K} \\ m_1, \dots, m_{2K}}} \alpha_{n_1 \dots n_{2K}} \alpha_{m_1 \dots m_{2K}}^* \\ &\times \prod_{i=1}^{2K} \frac{e^{-X_i^2} H_{n_i}(X_i) H_{m_i}(X_i)}{\sqrt{\pi^{1/2} 2^{n_i} n_i!} \sqrt{\pi^{1/2} 2^{m_i} m_i!}} \end{aligned} \quad (23)$$

Notice that the tensor $\alpha_{n_1, \dots, n_{2K}}$ is precisely the output of ‘state.ket()’ in strawberryfields. Also the Hermite polynomials are implemented in ‘scipy.special.hermite’:

```
from scipy import special
p = special.hermite(3, monic=False)
```

After extracting as well the mean photon numbers $\langle n_{i,\alpha} \rangle$, one obtains $\langle H_{(3,3)} \rangle$.

V. VARIATIONAL ALGORITHM

Depending on the model $H_{(\mu,\nu)}$ of interest, we define the following loss function:

$$\mathcal{C}_{(\mu,\nu)}(\theta) := \langle\psi(\theta)|H_{(\mu,\nu)}|\psi(\theta)\rangle \quad (24)$$

In order to compute this loss, one therefore has to measure both the photon number operator on each channel, as well as the position quadrature operator on each channel, as described in the previous section.

Algorithm 1: Extract distribution of observable

Parameters: statevector $|\psi\rangle$, observable \mathcal{O} , shots $M \in \mathbb{N}$

Result: Probability distribution of \mathcal{O} in state $|\psi\rangle$

```

for  $m = 1$  to  $\dim(\text{spec}(\mathcal{O}))$  do
  | Initialize  $N_m \leftarrow 0$ ;
for  $j = 1$  to  $M$  do
  | Measure the state  $|\psi\rangle$  in the basis  $\mathcal{O}$ , obtain the
  | eigenvalue  $o_m$ , set  $N_m \leftarrow N_m + 1$ ;
for  $m = 1$  to  $\dim(\text{spec}(\mathcal{O}))$  do
  | Normalize  $N_m \leftarrow N_m/M$ ;
return  $(N_m)_m$ .

```

Algorithm 2: Computation of the loss

Parameters: $M \in \mathbb{N}$

Result: Value of the loss \mathcal{C}

```

Initialize  $\mathcal{C} \leftarrow 0$ ;
Get the position quadratures distribution with alg.
(1);
Get the photon numbers distribution with alg. (1);
Compute the loss  $\mathcal{C}$  using eq. (20);
return  $\mathcal{C}$ .

```

Algorithm 3: Training of the parameterized photonic circuit

Parameters: $N_{\text{steps}} \in \mathbb{N}$, initial parameters $\theta_0 \in \mathbb{R}^K$, learning rate $\eta \in \mathbb{R}_+$

Result: Optimized hyperparameters $\theta \in \mathbb{R}^K$

```

Initialize hyperparameters  $\theta \leftarrow \theta_0$ ;
for  $i = 1$  to  $N_{\text{steps}}$  do
  | Compute the loss  $\mathcal{C}$  with alg. (2);
  | Compute the gradient  $\nabla_{\theta}\mathcal{C}$  with shift rule and alg.
  | (2);
  | Update the parameters  $\theta \leftarrow \theta - \eta\nabla_{\theta}\mathcal{C}$ ;
end for
return  $\theta$ .

```

VI. RESULTS

We gather here the results of the simulations. We focus on the case of 2 QDOs. In particular we plot the profile of the binding energy as a function of the distance between the two nuclei. The binding energy is simply defined as the ground state energy of the interacting system to which one subtract the ground state energy of

the uninteracting system, namely that of a pair of free harmonic oscillators in this Drude model.

VII. QUANTUM MUTUAL INFORMATION

Let us give the expression of the partial density matrix associated to QDO₁. The total state of the system, we recall, is expressed in the Fock basis as

$$|\psi\rangle = \sum_{n_1, \dots, n_{2K}=0}^{\infty} \alpha_{n_1 \dots n_{2K}} |n_1\rangle \otimes \dots \otimes |n_{2K}\rangle, \quad (25)$$

leading to the following expression for the density matrix:

$$\rho = \sum_{\substack{n_1, \dots, n_{2K} \\ m_1, \dots, m_{2K}}} \alpha_{m_1 \dots m_{2K}}^* \alpha_{n_1 \dots n_{2K}} |n_1\rangle \langle m_1| \otimes \dots \otimes |n_{2K}\rangle \langle m_{2K}|.$$

The partial trace associated to QDO₁ is therefore given by:

$$\rho_1 = \sum_{\substack{n_1, \dots, n_K \\ m_1, \dots, m_K \\ l_1, \dots, l_K}} \alpha_{m_1 \dots m_K l_1, \dots, l_K}^* \alpha_{n_1 \dots n_K l_1, \dots, l_K} |n_1\rangle \langle m_1| \otimes \dots \otimes |n_K\rangle \langle m_K|. \quad (26)$$

Since the state of the total system is pure, the von Neumann entropy of the total density matrix is zero. QDO number 2 can then be interpreted as purifying the system composed solely of QDO number 1. The two QDO therefore have identical von Neumann entropy $S(\rho_1)$, the entanglement entropy. The quantum mutual information of the system is therefore given by

$$I(1:2) = S(\rho_1) + S(\rho_2) - S(\rho) = 2S(\rho_1), \quad (27)$$

with the von Neumann entropy being defined as

$$S(\rho) = -\text{Tr} \rho \log \rho. \quad (28)$$

The profile of the quantum mutual information as a function of the interatomic distance is provided in fig. (...)

VIII. CONCLUSION**ACKNOWLEDGMENTS****CODE AVAILABILITY**

The reader will find an open source python code accompanying this paper at github repository.

[1] H. Margenau and N. R. Kestner, *Theory of Intermolecular Forces: International Series of Monographs in Natural*

Philosophy. Elsevier, 2013, vol. 18.

- [2] I. G. Kaplan, *Intermolecular interactions: physical picture, computational methods and model potentials*. John Wiley & Sons, 2006.
- [3] A. Stone, *The theory of intermolecular forces*. oUP Oxford, 2013.
- [4] J. O. Hirschfelder, *Intermolecular Forces, Volume 12*. John Wiley & Sons, 2009, vol. 12.
- [5] H. B. Casimir and D. Polder, “The influence of retardation on the london-van der waals forces,” *Physical Review*, vol. 73, no. 4, p. 360, 1948.
- [6] S. Y. Buhmann, *Dispersion Forces I: Macroscopic quantum electrodynamics and ground-state Casimir, Casimir-Polder and van der Waals forces*. Springer, 2013, vol. 247.
- [7] S. Y. Buhmann and D.-G. Welsch, “Dispersion forces in macroscopic quantum electrodynamics,” *Progress in quantum electronics*, vol. 31, no. 2, pp. 51–130, 2007.
- [8] G. Compagno, R. Passante, and F. Persico, *Atom-field interactions and dressed atoms*, 1995.
- [9] R. Passante, “Dispersion interactions between neutral atoms and the quantum electrodynamical vacuum,” *Symmetry*, vol. 10, no. 12, p. 735, 2018.
- [10] P. W. Milonni, *The quantum vacuum: an introduction to quantum electrodynamics*. Academic press, 2013.
- [11] R. Schmid, “Recent advances in the description of the structure of water, the hydrophobic effect, and the like-dissolves-like rule,” *Monatshefte für Chemie/Chemical Monthly*, vol. 132, no. 11, pp. 1295–1326, 2001.
- [12] J. Hoja, H.-Y. Ko, M. A. Neumann, R. Car, R. A. DiStasio Jr, and A. Tkatchenko, “Reliable and practical computational description of molecular crystal polymorphs,” *Science Advances*, vol. 5, no. 1, p. eaau3338, 2019.
- [13] J. Hoja and A. Tkatchenko, “First-principles stability ranking of molecular crystal polymorphs with the dft+mbd approach,” *Faraday Discussions*, vol. 211, pp. 253–274, 2018.
- [14] M. Mortazavi, J. G. Brandenburg, R. J. Maurer, and A. Tkatchenko, “Structure and stability of molecular crystals with many-body dispersion-inclusive density functional tight binding,” *The journal of physical chemistry letters*, vol. 9, no. 2, pp. 399–405, 2018.
- [15] M. Stöhr and A. Tkatchenko, “Quantum mechanics of proteins in explicit water: The role of plasmon-like solute-solvent interactions,” *Science advances*, vol. 5, no. 12, p. eaax0024, 2019.
- [16] A. M. Reilly and A. Tkatchenko, “Role of dispersion interactions in the polymorphism and entropic stabilization of the aspirin crystal,” *Physical review letters*, vol. 113, no. 5, p. 055701, 2014.
- [17] M. Galante and A. Tkatchenko, “Anisotropic van der waals dispersion forces in polymers: Structural symmetry breaking leads to enhanced conformational search,” *arXiv preprint arXiv:2110.06646*, 2021.
- [18] A. Kleshchonok and A. Tkatchenko, “Tailoring van der waals dispersion interactions with external electric charges,” *Nature communications*, vol. 9, no. 1, pp. 1–9, 2018.
- [19] A. Ambrosetti, P. Umari, P. L. Silvestrelli, J. Elliott, and A. Tkatchenko, “Optical van-der-waals forces in molecules: from electronic bethe-salpeter calculations to the many-body dispersion model,” *Nature communications*, vol. 13, no. 1, pp. 1–8, 2022.
- [20] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Photons and Atoms-Introduction to Quantum Electrodynamics*, 1997.
- [21] —, *Atom-photon interactions: basic processes and applications*, 1998.
- [22] G. Preparata, *QED Coherence in matter*. World Scientific, 1995.
- [23] A. Salam, *Molecular quantum electrodynamics: long-range intermolecular interactions*. John Wiley & Sons, 2009.
- [24] D. Craig, “T. thirunamachandran in molecular quantum electrodynamics,” 1998.
- [25] D. Richardson, “Dispersion contribution of two-atom interaction energy: Multipole interactions,” *Journal of Physics A: Mathematical and General*, vol. 8, no. 11, p. 1828, 1975.
- [26] J. Mahanty and B. Ninham, “Dispersion contributions to surface energy,” *The Journal of Chemical Physics*, vol. 59, no. 11, pp. 6157–6162, 1973.
- [27] L. Woods, D. A. R. Dalvit, A. Tkatchenko, P. Rodriguez-Lopez, A. W. Rodriguez, and R. Podgornik, “Materials perspective on casimir and van der waals interactions,” *Reviews of Modern Physics*, vol. 88, no. 4, p. 045003, 2016.
- [28] A. Tkatchenko, “Current understanding of van der waals effects in realistic materials,” *Advanced Functional Materials*, vol. 25, no. 13, pp. 2054–2061, 2015.
- [29] X. Ren, P. Rinke, C. Joas, and M. Scheffler, “Random-phase approximation and its applications in computational chemistry and materials science,” *Journal of Materials Science*, vol. 47, no. 21, pp. 7447–7471, 2012.
- [30] J. Harl and G. Kresse, “Accurate bulk properties from approximate many-body techniques,” *Physical review letters*, vol. 103, no. 5, p. 056401, 2009.
- [31] J. F. Dobson and T. Gould, “Calculation of dispersion energies,” *Journal of Physics: Condensed Matter*, vol. 24, no. 7, p. 073201, 2012.
- [32] V. A. Parsegian, *Van der Waals forces: a handbook for biologists, chemists, engineers, and physicists*. Cambridge university press, 2005.
- [33] A. D. Becke and E. R. Johnson, “A simple effective potential for exchange,” *The Journal of chemical physics*, vol. 124, no. 22, p. 221101, 2006.
- [34] —, “Exchange-hole dipole moment and the dispersion interaction: High-order dispersion coefficients,” *The Journal of chemical physics*, vol. 124, no. 1, p. 014104, 2006.
- [35] S. Grimme, J. Antony, S. Ehrlich, and H. Krieg, “A consistent and accurate ab initio parametrization of density functional dispersion correction (dft-d) for the 94 elements h-pu,” *The Journal of chemical physics*, vol. 132, no. 15, p. 154104, 2010.
- [36] S. Grimme, “Semiempirical gga-type density functional constructed with a long-range dispersion correction,” *Journal of computational chemistry*, vol. 27, no. 15, pp. 1787–1799, 2006.
- [37] A. Tkatchenko, R. A. DiStasio Jr, R. Car, and M. Scheffler, “Accurate and efficient method for many-body van der waals interactions,” *Physical review letters*, vol. 108, no. 23, p. 236402, 2012.
- [38] D. Massa, A. Ambrosetti, and P. L. Silvestrelli, “Many-body van der waals interactions beyond the dipole approximation,” *The Journal of Chemical Physics*, vol. 154, no. 22, p. 224115, 2021.
- [39] A. Ambrosetti, A. M. Reilly, R. A. DiStasio Jr, and A. Tkatchenko, “Long-range correlation energy calcu-

- lated from coupled atomic response functions,” *The Journal of chemical physics*, vol. 140, no. 18, p. 18A508, 2014.
- [40] W. L. Bade and J. G. Kirkwood, “Drude-model calculation of dispersion forces. ii. the linear lattice,” *The Journal of Chemical Physics*, vol. 27, no. 6, pp. 1284–1288, 1957. [Online]. Available: <https://doi.org/10.1063/1.1743992>
- [41] W. L. Bade, “Drude-model calculation of dispersion forces. i. general theory,” *The Journal of Chemical Physics*, vol. 27, no. 6, pp. 1280–1284, 1957. [Online]. Available: <https://doi.org/10.1063/1.1743991>
- [42] M. R. Karimpour, D. V. Fedorov, and A. Tkatchenko, “Quantum framework for describing retarded and nonretarded molecular interactions in external electric fields,” *Physical Review Research*, vol. 4, no. 1, jan 2022. [Online]. Available: <https://doi.org/10.1103/2Fphysrevresearch.4.013011>
- [43] —, “Comprehensive Quantum Framework for Describing Retarded and Non-Retarded Molecular Interactions in External Electric Fields,” 2021.
- [44] M. Gori, P. Kurian, and A. Tkatchenko, “Second quantization approach to many-body dispersion interactions,” 2022. [Online]. Available: <https://arxiv.org/abs/2205.11549>
- [45] D. Massa, A. Ambrosetti, and P. L. Silvestrelli, “Beyond-dipole van der waals contributions within the many-body dispersion framework,” *Electronic Structure*, vol. 3, no. 4, p. 044002, 2021.
- [46] B. L. Blaney, B. BL, and E. GE, “Van der waals molecules.” 1976.
- [47] A. Jones, “Quantum drude oscillators for accurate many-body intermolecular forces,” 2010.
- [48] A. P. Jones, J. Crain, V. P. Sokhan, T. W. Whitfield, and G. J. Martyna, “Quantum drude oscillator model of atoms and molecules: Many-body polarization and dispersion interactions for atomistic simulation,” *Phys. Rev. B*, vol. 87, p. 144103, Apr 2013. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevB.87.144103>
- [49] M. Sadhukhan and F. R. Manby, “Quantum mechanics of drude oscillators with full coulomb interaction,” *Physical Review B*, vol. 94, no. 11, p. 115106, 2016.
- [50] K. Tang and J. P. Toennies, “An improved simple model for the van der waals potential based on universal damping functions for the dispersion coefficients,” *The Journal of chemical physics*, vol. 80, no. 8, pp. 3726–3741, 1984.
- [51] N. M. Tubman and J. McMinis, “Renyi entanglement entropy of molecules: Interaction effects and signatures of bonding,” *arXiv preprint arXiv:1204.4731*, 2012.
- [52] S.-H. Zhang and Z.-Y. Yan, “Frequency dependence of the entanglement entropy production in a system of coupled driven nonlinear oscillators,” *Entropy*, vol. 21, no. 9, p. 889, 2019.
- [53] S. Lloyd and S. L. Braunstein, “Quantum computation over continuous variables,” *Physical Review Letters*, vol. 82, no. 8, p. 1784, 1999.
- [54] P. Van Loock, W. Munro, K. Nemoto, T. Spiller, T. Ladd, S. L. Braunstein, and G. Milburn, “Hybrid quantum computation in quantum optics,” *Physical Review A*, vol. 78, no. 2, p. 022303, 2008.
- [55] N. Killoran, J. Izaac, N. Quesada, V. Bergholm, M. Amy, and C. Weedbrook, “Strawberry fields: A software platform for photonic quantum computing,” *Quantum*, vol. 3, p. 129, 2019.
- [56] M. A. Nielsen and I. Chuang, “Quantum computation and quantum information,” 2002.
- [57] A. K. Pati, S. L. Braunstein, and S. Lloyd, “Quantum searching with continuous variables,” *arXiv preprint quant-ph/0002082*, 2000.
- [58] A. K. Pati and S. L. Braunstein, “Deutsch-jozsa algorithm for continuous variables,” *Quantum Information with Continuous Variables*, pp. 31–36, 2003.
- [59] S. L. Braunstein and P. Van Loock, “Quantum information with continuous variables,” *Reviews of modern physics*, vol. 77, no. 2, p. 513, 2005.
- [60] U. L. Andersen, G. Leuchs, and C. Silberhorn, “Continuous-variable quantum information processing,” *Laser & Photonics Reviews*, vol. 4, no. 3, pp. 337–354, 2010.
- [61] S. J. Lomonaco Jr and L. H. Kauffman, “A continuous variable shor algorithm,” *arXiv preprint quant-ph/0210141*, 2002.
- [62] S. L. Braunstein, “Error correction for continuous quantum variables,” *Physical Review Letters*, vol. 80, no. 18, p. 4084, 1998.
- [63] J. M. Arrazola, V. Bergholm, K. Brádler, T. R. Bromley, M. J. Collins, I. Dhand, A. Fumagalli, T. Gerrits, A. Goussev, L. G. Helt *et al.*, “Quantum circuits with many photons on a programmable nanophotonic chip,” *Nature*, vol. 591, no. 7848, pp. 54–60, 2021.
- [64] M. Tillmann, B. Dakić, R. Heilmann, S. Nolte, A. Szameit, and P. Walther, “Experimental boson sampling,” *Nature photonics*, vol. 7, no. 7, pp. 540–544, 2013.
- [65] H.-S. Zhong, H. Wang, Y.-H. Deng, M.-C. Chen, L.-C. Peng, Y.-H. Luo, J. Qin, D. Wu, X. Ding, Y. Hu *et al.*, “Quantum computational advantage using photons,” *Science*, vol. 370, no. 6523, pp. 1460–1463, 2020.
- [66] T. R. Bromley, J. M. Arrazola, S. Jahangiri, J. Izaac, N. Quesada, A. D. Gran, M. Schuld, J. Swinerton, Z. Zabaneh, and N. Killoran, “Applications of near-term photonic quantum computers: software and algorithms,” *Quantum Science and Technology*, vol. 5, no. 3, p. 034010, 2020.
- [67] M. Schuld and F. Petruccione, *Machine learning with quantum computers*. Springer, 2021.
- [68] N. Killoran, T. R. Bromley, J. M. Arrazola, M. Schuld, N. Quesada, and S. Lloyd, “Continuous-variable quantum neural networks,” *Physical Review Research*, vol. 1, no. 3, p. 033063, 2019.
- [69] J. M. Arrazola, T. R. Bromley, J. Izaac, C. R. Myers, K. Brádler, and N. Killoran, “Machine learning method for state preparation and gate synthesis on photonic quantum computers,” *Quantum Science and Technology*, vol. 4, no. 2, p. 024004, 2019.
- [70] Y. Enomoto, K. Anai, K. Udagawa, and S. Takeda, “Continuous-variable quantum approximate optimization on a programmable photonic quantum processor,” *arXiv preprint arXiv:2206.07214*, 2022.
- [71] M. Cerezo, A. Arrasmith, R. Babbush, S. C. Benjamin, S. Endo, K. Fujii, J. R. McClean, K. Mitarai, X. Yuan, L. Cincio *et al.*, “Variational quantum algorithms,” *Nature Reviews Physics*, vol. 3, no. 9, pp. 625–644, 2021.
- [72] A. Peruzzo, J. McClean, P. Shadbolt, M.-H. Yung, X.-Q. Zhou, P. J. Love, A. Aspuru-Guzik, and J. L. O’Brien, “A variational eigenvalue solver on a photonic quantum processor,” *Nature communications*, vol. 5, no. 1, p. 4213, 2014.

- [73] K. Mitarai, M. Negoro, M. Kitagawa, and K. Fujii, “Quantum circuit learning,” *Physical Review A*, vol. 98, no. 3, p. 032309, 2018.
- [74] K. Marshall, R. Pooser, G. Siopsis, and C. Weedbrook, “Quantum simulation of quantum field theory using continuous variables,” *Physical Review A*, vol. 92, no. 6, p. 063825, 2015.
- [75] K. Yeter-Aydeniz, E. Moschandreou, and G. Siopsis, “Quantum imaginary-time evolution algorithm for quantum field theories with continuous variables,” *Physical Review A*, vol. 105, no. 1, p. 012412, 2022.
- [76] O. Kiss, F. Tacchino, S. Vallecorsa, and I. Tavernelli, “Quantum neural networks force fields generation,” *Machine Learning: Science and Technology*, vol. 3, no. 3, p. 035004, 2022.
- [77] J. Apanavicius, Y. Feng, Y. Flores, M. Hassan, and M. McGuigan, “Morse potential on a quantum computer for molecules and supersymmetric quantum mechanics,” *arXiv preprint arXiv:2102.05102*, 2021.
- [78] L. W. Anderson, M. Kiffner, P. K. Barkoutsos, I. Tavernelli, J. Crain, and D. Jaksch, “Coarse-grained intermolecular interactions on quantum processors,” *Physical Review A*, vol. 105, no. 6, p. 062409, 2022.
- [79] H. Wang, S. Ashhab, and F. Nori, “Efficient quantum algorithm for preparing molecular-system-like states on a quantum computer,” *Physical Review A*, vol. 79, no. 4, p. 042335, 2009.
- [80] Y. Nam, J.-S. Chen, N. C. Pienti, K. Wright, C. Delaney, D. Maslov, K. R. Brown, S. Allen, J. M. Amini, J. Apisdorf *et al.*, “Ground-state energy estimation of the water molecule on a trapped-ion quantum computer,” *npj Quantum Information*, vol. 6, no. 1, p. 33, 2020.
- [81] J. Romero, R. Babbush, J. R. McClean, C. Hempel, P. J. Love, and A. Aspuru-Guzik, “Strategies for quantum computing molecular energies using the unitary coupled cluster ansatz,” *Quantum Science and Technology*, vol. 4, no. 1, p. 014008, 2018.
- [82] J. I. Colless, V. V. Ramasesh, D. Dahlen, M. S. Blok, M. E. Kimchi-Schwartz, J. R. McClean, J. Carter, W. A. de Jong, and I. Siddiqi, “Computation of molecular spectra on a quantum processor with an error-resilient algorithm,” *Physical Review X*, vol. 8, no. 1, p. 011021, 2018.