

Regression discontinuity design

Matthieu Stigler
Matthieu.Stigler@gmail.com

NIPFP, 3 June 2014

Outline

Introduction

The econometrics
of RDD

The statistics of
RDD

Sensitivity tests

Conclusion

- ① Introduction
- ② The econometrics of RDD
- ③ The statistics of RDD
- ④ Sensitivity tests
- ⑤ Conclusion

The idea

Introduction

The econometrics
of RDD

The statistics of
RDD

Sensitivity tests

Conclusion

The goal: Estimate a treatment effect in a case where there might be self-selection/endogeneity.

The framework: Selection is based on a simple known (administrative) rule, based on a cutoff

The idea: treated and untreated are in general different, but those just around the cutoff should be similar.

The solution: make a local comparison, comparing just above with just below

Two cases

Introduction

The econometrics
of RDD

The statistics of
RDD

Sensitivity tests

Conclusion



Two (three) cases:

Sharp RDD: Assignment is strict (i.e. all above are treated, all below untreated)

Fuzzy RDD: Assignment does not lead to full compliance

Mixed: One sided-compliance

Figure 1: Assignment probabilities in sharp versus fuzzy



Examples

Age: Retirement and Medicare (Card 2004). Alcohol in US and mortality (Carpenter et al 2007)

School score: Angrist et al. 2011, 2012...

Class size: If above 40 students, split 1 class into 2. (Angrist Lavy 1999)

Geo. location: Geo-wise treatment (for state, country): compare individuals at the border

Majority rule: For elections, Lee (2008)

Height: Military service ?

Poverty rate: Indian BPL? Super fuzzy...

Introduction

The econometrics of RDD

The statistics of RDD

Sensitivity tests

Conclusion

The history

Introduction

The econometrics
of RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Majour contributions in eco:

- Thistlewaite and Campbell (1960)
- Angrist (1999) Fuzzy RDD, early application in economics
- Hahn et al (2001) Identification of ATE by RDD
- Porter (2003): Estimation of RDD
- Imbens and Lemieux (2008) State of the art in RDD
- Lee and Lemieux (2010) State of the art in RDD

cf Cook (2008) *Waiting for Life to Arrive: A History of the RDD*

Causality in econometrics

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Granger: X causes Y if X helps predict Y

Counterfactual: X causes Y if there is a difference between the Y observed when X occurred and the un-observed counterfactual Y *if X had not occurred*.

Refs: Granger (1969), Holland (1980)

Causality in econometrics

Definitions:

Def: treatment variable

Individual i has treatment status:

$$D_i = \begin{cases} 1 & \text{if treated} \\ 0 & \text{if not treated} \end{cases}$$

Def: potential outcome

Individual i has outcome:

$Y_i(D_i)$: observed outcome (treated or not)
 $Y_i(1 - D_i)$: countefactual/hypothetical outcome

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Term counter-factual

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Beware that counterfactual is used in two ways:

Hypothetical: The unobserved outcome: $Y_i(0)|D_i = 1$ or
 $Y_i(1)|D_i = 0$

Approximation: The closest factual: $Y_i(0)|D_i = 0$ for $Y_i(0)|D_i = 1$

Term has changed from: we use **as** counter-factual to we use **the** counter-factual

Term counter-factual II

An example of the similarity of counter-factual notions:

Everyday life

Do you feel warmer with the jacket?

- Than you were before? \Rightarrow **closest factual**
- Than you would be without? \Rightarrow **unobserved outcome**

Counterfactual thinking is not new:

- Used every day, studied extensively in psychology
- Lewis (1973) "Causation", *Journal of Philosophy*
- Used in law, forensic medicine...

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Term counter-factual III

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Another confusion in economics about Counterfactual analysis

CGE: What if policy A (country A reduces tariff by 3%)

Impact evaluation: Answers factual questions by asking
counter-factual questions.

But IE answers counter-factual questions: treatment effect on the
untreated.

Causal inference

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Def: Effects of the causes

The cause D causes the effect $Y_i(D = 1) - Y_i(D = 0)$

Fundamental Problem of Causal Inference (Holland 1980)

It is impossible to observe simultaneously the value of $Y_i(D = 1)$ and $Y_i(D = 0)$ on the same unit and, therefore, it is impossible to observe the effect of D for i .

Causal inference: the statistical approach

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

In statistics, given the fundamental problem, we do inference for population, not individuals:

Definition: average treatment effect

$$ATE \equiv E[Y_i(1) - Y_i(0)]$$

Back to causalities

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

We see now more fundamental differences between Granger and Counterfactual causality:

Granger: Defines causes of effects: Among X_1, X_2, \dots, X_k which cause(s) Y ?

Counterfactual: Measures effects of causes: what is effect of X_1 on Y ?

But is it really effects of causes? Most usually, is single effect of single cause.

Causal inference: the statistical approach

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Two more definitions

Definition

ATE on treated: $ATT \equiv E[Y_i(1) - Y_i(0)|D = 1]$

ATE on untreated: $ATU \equiv E[Y_i(1) - Y_i(0)|D = 0]$

Estimating the ATE

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

One could think that ATE is estimated simply:

$$\begin{aligned}ATE &\equiv E[Y_i(1) - Y_i(0)] \\ &= E[Y_i(1)] - E[Y_i(0)]\end{aligned}$$

Unfortunately, we do observe $E[Y_i(1)]$ only for the $\{j|D_j = 1\}$ i.e. we observe $E[Y_i(1)|D_i = 1]$!

The frameworks in impact evaluation:

Sample selection: Old-fashioned

RCT: allocate D randomly, so
 $E[Y_i(1)|D_i = 1] = E[Y_i(1)]$ and hence just
compare means!

Matching: Assume D is based on observables: ATT and ATU,
ATE

Diff-diff Assume D is based only on time-invariant
unobservables: ATT

IV: Find exogenous moves in D (estimating
LATE/CATE)

RDD: Assume D is random close to threshold: Local ATE

Identification in the sharp case

In the sharp RDD, treatment is assigned based on whether the forcing variable x is below/above a threshold c .

$$D_i = 1(x_i \geq c) = \begin{cases} 1 & \text{if } x_i \geq c \\ 0 & \text{if } x_i < c \end{cases}$$

We will identify the ATE at a point:

Definition: point ATE

$$ATE(c) \equiv E[Y_i(1) - Y_i(0) | x_i = c]$$

Note here we have a local ATE in a well-defined sense (local in space of x)!

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Figure 2: Assignment probabilities in sharp versus fuzzy






Figure 3: Assignment probabilities in sharp versus fuzzy



The RDD estimand

To estimate:

$$ATE(c) \equiv E[Y_i(1) - Y_i(0)|x_i = c] = \overbrace{E[Y_i(1)|c]}^{\text{Factual}} - \overbrace{E[Y_i(0)|c]}^{\text{Counter-factual}}$$

we approximate the *counter-factual* untreated at c by the *factual* untreated just below c

$$\lim_{x \uparrow c} E[Y_i(0)|x_i = c] \approx E[Y_i(0)|x_i = c]$$

So end up using:

$$\overbrace{\lim_{x \downarrow c} E[Y_i(1)|x_i = c]}^{\text{Just treated}} - \overbrace{\lim_{x \uparrow c} E[Y_i(0)|x_i = c]}^{\text{Just untreated}}$$

Assumptions

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Main assumptions

- Identifying assumption: $E[Y(1)|c]$ and $E[Y(0)|c]$ are continuous in c .
- No manipulation of x around c

Assumption 2

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests


Conclusion

Test 2: Manipulation of forcing variable

Check if individuals can manipulate their value around x ?

- Age? (if *declared* age...)
- Declared income?
- Number of employees?
- Test score: can retake exam?

Figure 4: Assignment probabilities in sharp versus fuzzy



Cases

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Sharp case: $D_i = 1(x_i \geq c)$

Fuzzy case: $\lim_{x \downarrow c} Pr(D_i = 1|x) \neq \lim_{x \uparrow c} Pr(D_i = 1|x)$

Sharp new: $\overbrace{\lim_{x \downarrow c} Pr(D_i = 1|x)}^{=1} - \overbrace{\lim_{x \uparrow c} Pr(D_i = 1|x)}^{=0} = 1$

Mixed case: One-sided compliance only

Cases

In the fuzzy RDD, we use:

$$ATE \equiv \frac{\lim_{x \downarrow c} E[Y|X = x] - \lim_{x \uparrow c} E[Y|X = x]}{\lim_{x \downarrow c} E[D|X = x] - \lim_{x \uparrow c} E[D|X = x]} \underbrace{\frac{\lim_{x \downarrow c} E[D|X = x] - \lim_{x \uparrow c} E[D|X = x]}{\lim_{x \downarrow c} E[D|X = x] - \lim_{x \uparrow c} E[D|X = x]}}_{=1 \text{ for sharp!}}$$

This is just the form the IV takes for binary D and binary Z: Wald estimator! (see MHE 4.1.2)

$$IV_{\text{Wald}} \equiv \frac{E[Y|z_i = 1] - E[Y|z_i = 0]}{E[D|z_i = 1] - E[D|z_i = 0]}$$

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Identification in the fuzzy case

Introduction

The econometrics
of RDD

Causal inference with
potential outcomes

Sharp RDD

Fuzzy RDD

The statistics of
RDD

Sensitivity tests

Conclusion

But in the fuzzy case, like with standard IV, we estimate only the complier-average effect (CATE/LATE)

And here, we only get a local version of that! \Rightarrow local-local-average!

On the IV interpretation of RDD

Remember the two conditions for IV with Y, X, Z :


Relevant: Should be relevant in first stage ($\beta \neq 0$) :

$$X_i = \alpha + \beta Z_i + \varepsilon_i^X$$

Exclusion: Should be excluded of structural form: ($\theta = 0$) :

$$Y_i = \gamma + \delta X_i + \theta Z_i + \varepsilon_i^Y$$

RDD recognises $\theta = 0$ only locally!



Parametric model

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

We want to estimate: $\lim_{x \downarrow c} E[Y_i(1)|c] - \lim_{x \uparrow c} E[Y_i(0)|c]$

So can estimate separately on each side of each side of c :

Left: $(1 - D)Y = \alpha_l + \beta_l(1 - D)(X - c) + \epsilon_l$

Right: $DY = \alpha_r + \beta_r D(X - c) + \epsilon_r$

with $D = I(X \geq c)$, and test $H_0 : \alpha_r = \alpha_l$

Assuming homoskedasticity ($\sigma_l = \sigma_r$, can estimate instead:

$$Y = \alpha + \tau D + \beta_1(X - c) + \beta_2 D(X - c) + \epsilon$$

Now we have $\tau = \alpha_r - \alpha_l$

Parametric model: extension II

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

Eventually, we could assume the same slope on each side:

$$Y = \alpha + \tau D + \beta(X - c) + \epsilon$$

Parametric model: extension II

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

Eventually, add higher-order polynomials:

$$Y = \alpha + \tau D + \beta_1^1(X - c) + \beta_2^1 D(X - c) + \dots + \beta_1^p(X - c)^p + \beta_2^p D(X - c)^p + \epsilon$$

What if use polynomial of order zero? **Group means test (=RCT!)**

What is parametric?

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

What is parametric?

- Functional form
- Distributional assumptions (cf MLE)

What is non-parametric here?

- Local/pointwise estimation **This is exactly what we want!**

Local/pointwise estimation

With kernel regression/smoother, want to estimate:

$$E(Y|X) = m(X)$$

The idea is to estimate $\hat{y}(x_i)$ averaging over local neighbours.

$$\hat{m}_h(x) = \sum_{i=1}^n w_i Y_i$$

Two approaches for w_i :

- use k nearest neighbours.
- use all neighbours within a window h .

Local/pointwise estimation

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

The Nadaraya-Watson (1964) estimator use a window with weights:

$$\hat{m}_h(x) = \frac{\sum_{i=1}^n \mathcal{K}_h(x - X_i) Y_i}{\sum_{i=1}^n \mathcal{K}_h(x - X_i)}$$

where \mathcal{K} is a kernel with a bandwidth h .

Kernel and bandwidth choice

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion


Parameters in non-parametric estimation:

Kernel \mathcal{K} : Rectangular, Gaussian...

Bandwidth h :

- $h \searrow$: bias \searrow , variance \nearrow
- $h \nearrow$: bias \nearrow , variance \searrow

Figure 5: various kernels



Problems

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

Problems with Nadaraya-Watson:

- Biased
- Higher bias at boundary point **This is however what we want :-)**

Solution: Use local-linear regression

Local-linear regression

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

Local-linear regression (LLR): fit a local weighted OLS at every point!
So we get $\hat{m}(x_i)$, and also for every point get a new $\hat{\alpha}(x), \hat{\beta}(x)$

It is a simple weighted regression, with weights obtained from kernel around x .

$$\hat{\alpha}(c), \hat{\beta}(c), \hat{\tau}(c) = \arg \min_{\alpha, \beta, \tau} \sum_{i=1}^n (Y_i - \alpha - \tau D - \beta(X_i - c))^2 \mathcal{K} \left(\frac{X_i - c}{h} \right)$$

Local-linear regression

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

Links with LLR:

OLS: If use LLR with $h \rightarrow \infty$ and rectangular \mathcal{K} :
LLR \rightarrow OLS!

Nadaraya-Watson: NW is a LLR with only a constant!

Choice of bandwidth

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

Two procedures to choose bandwidth:

Plug-in: minimise (asymptotic) $\text{MSE}(h) = \text{bias}(h)^2 + \text{var}(h)$
plugging-in estimates of bias and variance

Cross-validation: Minimise empirical MSE: $\frac{1}{N} \sum_{i=1}^N (Y_i - \hat{\mu}(X_i))^2$

Inference

Introduction

The econometrics
of RDD

The statistics of
RDD

Parametric approach

Nonparametric
approaches

Sensitivity tests

Conclusion

Distribution: Normal!

Two types of variance estimators for non-parametric:

- Parametric: $\sigma (X' X)^{-1}$
- Non-parametric: Messy!!

Rate of convergence: slower than parametric!

Sensitivity tests

Introduction

The econometrics
of RDD

The statistics of
RDD

Sensitivity tests

Conclusion

- Bandwidth sensitivity
- Manipulation of forcing variable
- Placebo tests
- Discontinuity also for covariates?

Sensitivity tests: sensitivity to bandwidth

Introduction

The econometrics
of RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Test 1: sensitivity to bandwidth

Compute estimate at various bandwidths, compare

Sensitivity tests: Manipulation of forcing variable 1

Introduction

The econometrics
of RDD


The statistics of
RDD

Sensitivity tests

Conclusion

Test: look for discontinuity of density around threshold

McCrary (2008): look for discontinuity of the density of X at point c :



Sensitivity tests: Manipulation of forcing variable 2

Introduction

The econometrics
of RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Actually, this approach helps also to check how a constraint is binding, in general.

Cf paper of Card and DellaVigna (2012) *How Do Authors Respond to Page Limits*

Figure 6: AER limits (Card DellaVigna 2012)





Figure 7: JEEA limits (Card DellaVigna 2012)



Sensitivity tests: Placebo tests

Introduction

The econometrics
of RDD

The statistics of
RDD

Sensitivity tests

Conclusion

Test 3: Placebo tests

Test for effect at a point known for known effect: estimate RDD at non-cutoff point.

Test 4: Discontinuity also for covariates

Like placebo, run RDD using as new y a covariate.

But passing test is not necessary! Just reduces credibility.

RDD in a nutshell

- Estimating ATE just around cutoff
- So identify effect only around the cutoff, low external validity
- Step 1: Visual representation important
- Step 2: Estimation by parametric or non-parametric
- Step 3: Few sensitivity tests