

Are galaxies in compact groups special?

Matthieu Tricottet¹, Gary A. Mamon², and Eugenia Díaz-Giménez^{3,4}

¹ 80, rue d'Alésia, 75014 Paris, France

e-mail: matthieu.tricottet@gmail.com

² Institut d'Astrophysique de Paris (UMR 7095: CNRS & Sorbonne Université), 98 bis boulevard Arago, 75014 Paris, France

³ CONICET. Instituto de Astronomía Teórica y Experimental (IATE), Laprida 854, X5000BGR, Córdoba, Argentina

⁴ Universidad Nacional de Córdoba (UNC). Observatorio Astronómico de Córdoba (OAC), Laprida 854, X5000BGR, Córdoba, Argentina

ABSTRACT

We investigate the properties of galaxies in compact groups (CGs) and compare them to a control sample of galaxies.

Key words. galaxies: groups: general – galaxies: evolution – galaxies: interactions – galaxies: star formation – galaxies: statistics

1. Introduction

2. Data

2.1. Samples

We use the compact-group and control samples constructed in our previous study [Tricottet et al. \(2025\)](#). We briefly recall their construction here. Galaxy properties are taken from the SDSS DR12 catalogue of [Tempel et al. \(2017\)](#), while regular groups are drawn from the SDSS DR7 group catalogue of [Lim et al. \(2017\)](#); the two catalogues were matched through SDSS object identifiers to define the Lim–Tempel working catalogue. Compact groups are selected from the modified Hickson compact-group catalogue of [Díaz-Giménez et al. \(2018\)](#). Starting from the reliable HMCs with exactly four members, we retain only systems in which all galaxies are present in the Lim–Tempel catalogue and impose the volume- and luminosity-complete limits $0.005 < z_{\text{BGG}} < 0.0452$ and $M_{r,\text{BGG}} \leq -21.81$, with all 4 galaxies lying within 3 magnitudes of the brightest group galaxy. This defines the parent CG₄ sample of 78 compact groups. The regular-group comparison samples are built from Lim–Tempel groups satisfying the same redshift and BGG-luminosity limits, with at least 4 members and at least 3 satellites within 3 magnitudes of the BGG. From this parent control catalogue, we define RG₄ groups as regular groups with exactly four members, excluding systems identical to a CG₄, and we additionally use the Control_{4B} and Control_{4C} samples, formed respectively from the 4 brightest galaxies and from the BGG plus its 3 closest projected companions in each parent regular group, after removing groups that share galaxies with CG₄s. In contrast to our previous study, we remove CGs classified as *split* following [Zheng & Shen \(2021\)](#). This reduces the compact-group sample from 78 to 62 groups. We also remove the systems associated with Lim group 3688 from our Control_{4B} and Control_{4C} samples. This group contains 6 galaxies with redshifts between 0.032 and 0.034, and 1 with a redshift of

0.048. This group appears to be spurious, and the outlying galaxy, although not the BGG, is selected in both our original Control_{4B} and Control_{4C} samples, significantly affecting the group velocity dispersion (1979 and 1981 km s^{−1}, respectively).

We use SDSS DR 16 to retrieve stellar masses, star formation rates (SFRs), and Galaxy Zoo morphologies ([Lintott et al. 2011](#)). We specifically extract the columns `sfr_tot_p50`, `specsfr_tot_p50`, and `lgm_tot_p50` from the `galSpecExtra` table, and `p_el_debiased` and `p_cs_debiased` from the `zooSpec` table.

2.2. sSFR

To build a general star formation classification that could be uniformly applied to all our samples, we extracted galaxies from the SDSS DR 16 using the query shown in Listing 1.

```
SELECT
    s.specObjID,
    s.z,
    p.petroMag_r - p.extinction_r AS
        r_obs,
    p.petroMag_u - p.extinction_u AS
        u_obs,
    p.petroMag_g - p.extinction_g AS
        g_obs,
    p.petroMag_i - p.extinction_i AS
        i_obs,
    p.petroMag_z - p.extinction_z AS
        z_obs,
    p.objID,
    g.sfr_tot_p50, g.specsfr_tot_p50, g.
        lgm_tot_p50,
    l.h_alpha_eqw, l.h_beta_eqw, l.
        oiii_5007_eqw, l.nii_6584_eqw,
    l.h_alpha_flux, l.h_beta_flux, l.
        oiii_5007_flux, l.nii_6584_flux,
    z.p_el_debiased AS p_E,
    z.p_cs_debiased AS p_S
```




Fig. 1. BPT diagram showing our separation between non-AGN and AGN galaxies.

```

16 FROM SpecObj AS s
17 JOIN PhotoObj AS p ON s.bestObjID = p
   .objID
18 JOIN galSpecExtra as g ON s.specObjID
80 = g.specObjID
19 JOIN galSpecLine as l ON s.specObjID
   = l.specObjID
20 JOIN zooSpec AS z ON s.specObjID = z.
   specObjID
21 WHERE s.z BETWEEN 0.005 AND 0.0452
22 AND (p.petroMag_r - p.extinction_r <=
23 17.77)
24 AND s.class = 'GALAXY'
90 AND g.lgm_tot_p50 > -1000

```

Listing 1. Query used for selecting spectral & photometric data from the SDSS

We select non-AGN galaxies by first placing them on the classical BPT diagnostic of [Veilleux & Osterbrock \(1987\)](#) and requiring measured values of both emission-line ratios

$$\log_{10}([\text{N II}]\lambda 6584/\text{H}\alpha) \text{ and } \log_{10}([\text{O III}]\lambda 5007/\text{H}\beta).$$

A galaxy is flagged as an AGN if it satisfies either

$$\log_{10} \frac{[\text{N II}]}{\text{H}\alpha} > 0,$$

or if it lies above the empirical demarcation of [Kauffmann et al. \(2003\)](#):

$$\log_{10} \frac{[\text{O III}]}{\text{H}\beta} > \frac{0.61}{\log_{10}([\text{N II}]/\text{H}\alpha) - 0.05} + 1.3,$$

100 in which case it is removed from the non-AGN sample. All remaining galaxies—including those with missing line ratios—are retained as non-AGN. The non-AGN selection process is shown in Fig. 1.

Galaxies with `specsfr_tot_p50` equal to -9999 in the SDSS are classified as quenched (for display purposes, they are assigned the arbitrary low value $10^{-14.0} \text{ Gyr}^{-1}$). The remaining galaxies are separated into star-forming and passive populations using a two-component Gaussian Mixture Model (GMM) in the $\log M_*$ – $\log \text{sSFR}$ plane. Its parameters are determined by minimizing the Kullback–Leibler divergence between the empirical and model




Fig. 2. Observed non-quenched SDSS distribution in the $\log M_*$ – $\log \text{sSFR}$ plane and best-fitting two-component GMM density.

densities ([Kullback & Leibler \(1951\)](#); [McLachlan & Peel \(2000\)](#)). For any trial parameter vector θ , we reconstruct the mixture means $\{\mu_i\}$, covariances $\{\Sigma_i\}$, and weights $\{w_i\}$ via a Cholesky-like decomposition and logistic mapping, optionally constraining the second component’s mean to isolate the passive population ([Dempster et al. \(1977\)](#)). The KL divergence is estimated by constructing a 2D histogram of the data and summing $p_{\text{data}} \ln(p_{\text{data}}/p_{\text{GMM}})$, with a small regularization ϵ to avoid singularities ([Tojeiro et al. \(2009\)](#); [Bisigello et al. \(2018\)](#)). We optimize θ via L-BFGS-B (with fallback to Nelder–Mead) over multiple guided and random initializations, selecting the solution with lowest divergence ([Pedregosa et al. \(2011\)](#); [Bishop \(2006\)](#)). Initial means are seeded by a median split in $\log \text{sSFR}$ to improve convergence and robustness against local minima ([Rousseeuw \(1987\)](#)). In the final fit, one component naturally aligns with the high-sSFR “star-forming sequence,” while the other captures the low-sSFR “passive” cloud, providing a data-driven division that agrees with previous multi-Gaussian decompositions of the $\text{SFR} - M_*$ plane ([Wuyts et al. \(2011\)](#); [Hahn et al. \(2019\)](#)). The decision boundaries, defined as the loci where the posterior probabilities of adjacent components are equal, provide an objective criterion for delineating star-forming from passive galaxies.



The sSFR status for each sample is shown in Table 1. We find no significant difference in the Star forming fraction between CG_4 and the control samples; Fisher’s exact test gives p-values of 6.5×10^{-1} versus the $\text{Control}_{4\text{B}}$ sample, 2.5×10^{-1} versus $\text{Control}_{4\text{C}}$, and 1.8×10^{-2} versus RG_4 .

The same comparison restricted to BGGs and satellites is also shown in Table 1. For BGGs, the p-values obtained with two-sided Fisher exact test are 0.58 against $\text{Control}_{4\text{B}}$, 0.59 against $\text{Control}_{4\text{C}}$ and 0.37 against RG_4 . For satellites, the corresponding p-values are 0.30 against $\text{Control}_{4\text{B}}$, 0.03 against $\text{Control}_{4\text{C}}$ and 0.00 against RG_4 .


2.3. Morphology

Morphologies are determined via the Galaxy Zoo citizen-science decision tree, in which each galaxy image receives multiple independent volunteer classifications that are aggregated into debiased vote fractions ([Lintott et al. 2011](#)). Those fractions are provided in the `zooSpec` table of the SDSS database through the `p_el_debiased` and `p_cs_debiased` columns. We classify a galaxy as “elliptical” if `p_el_debiased` is greater than 0.5, and as “spiral” if `p_cs_debiased` is greater than 0.5. All other cases are classified as “uncertain”. The morphologies obtained for each

Sample		Quenched	Passive	Starforming
CG ₄		18 (7.3 %)	146 (58.9 %)	84 (33.9 %)
	<i>BGG</i>	6 (9.7 %)	45 (72.6 %)	11 (17.7 %)
	<i>Satellites</i>	12 (6.5 %)	101 (54.3 %)	73 (39.2 %)
Control _{4B}		181 (6.5 %)	1596 (57.1 %)	1019 (36.4 %)
	<i>BGG</i>	58 (8.3 %)	533 (76.4 %)	107 (15.3 %)
	<i>Satellites</i>	122 (5.8 %)	1061 (50.7 %)	911 (43.5 %)
Control _{4C}		191 (6.3 %)	1619 (53.8 %)	1198 (39.8 %)
	<i>BGG</i>	65 (8.7 %)	568 (75.6 %)	118 (15.7 %)
	<i>Satellites</i>	125 (5.5 %)	1050 (46.6 %)	1078 (47.8 %)
RG ₄		5 (2.2 %)	101 (45.1 %)	118 (52.7 %)
	<i>BGG</i>	3 (5.4 %)	38 (67.9 %)	15 (26.8 %)
	<i>Satellites</i>	2 (1.2 %)	63 (37.5 %)	103 (61.3 %)
<i>SDSS selection</i>		634 (1.2 %)	7705 (14.7 %)	44192 (84.1 %)

Table 1. Number of galaxies in each sSFR status for each sample.

Fig. 3. sSFR vs mass, star-forming/passive limit and Zoo morphologies for the SDSS control sample, excluding quenched galaxies.

Fig. 4. Distribution of residual sSFR offsets from the adopted star-forming main sequence.

sample are shown in Table 2. Barnard's two-sided exact tests were used to compare the fraction of each morphology between CG₄ and each control sample. The resulting p-values are 1.4×10^{-3} for ellipticals and 1.2×10^{-3} for spirals versus the Control_{4B} sample, 3.3×10^{-3} and 3.4×10^{-2} versus Control_{4C} and 2.2×10^{-7} and 5.9×10^{-7} versus RG₄.


Fig. 5. sSFR and morphologies of galaxies in CG₄ and our SDSS selection. Quenched galaxies are represented at an arbitrary low value of $10^{-14.0} \text{ Gyr}^{-1}$.

The galaxies in CG₄ systems, which we identified as cores of richer groups in Tricottet et al. (2025), have morphology distributions that differ significantly from those of ordinary four-member groups. Figure 5 displays sSFR versus stellar mass for galaxies in the SDSS selection and in CG₄.

2.4. Morphology versus sSFR

Among star-forming CG₄ galaxies, 58 are classified as spiral (84.1% after removing galaxies with uncertain morphology) and 11 are classified as elliptical (15.9% after removing galaxies with uncertain morphology). The corresponding numbers are 809 spiral galaxies (88.4%) and 106 elliptical galaxies (11.6%) for Control_{4B}, 851 spiral galaxies (84.3%) and 159 elliptical galaxies (15.7%) for Control_{4C}, and 91 spiral galaxies (91%) and 9 elliptical galaxies (9%) for RG₄. The probabilities that these fractions are drawn from the same distribution, estimated with Barnard two-sided exact test, are 0.29 against Control_{4B}, 0.97 against Control_{4C}, and 0.19 against RG₄. We therefore find neither

Sample	Elliptical	Spiral	Uncertain
CG ₄	124 (50.0 %)	86 (34.7 %)	38 (15.3 %)
Control _{4B}	1106 (39.6 %)	1266 (45.3 %)	420 (15.0 %)
Control _{4C}	1215 (40.4 %)	1249 (41.6 %)	540 (18.0 %)
RG ₄	60 (26.8 %)	129 (57.6 %)	35 (15.6 %)
<i>SDSS selection</i>	<i>10091 (19.2 %)</i>	<i>34715 (66.1 %)</i>	<i>7725 (14.7 %)</i>

Table 2. Number of galaxies of each morphology for each sample.

a significant excess nor a deficit of star-forming elliptical galaxies in CG₄ relative to any control sample.

When restricting the comparison to BGGs, we find 36 elliptical BGGs (58.1% of the BGGs) and 17 spiral BGGs (27.4%) in CG₄. The corresponding numbers are 405 elliptical BGGs (58.0%) and 195 spiral BGGs (27.9%) in Control_{4B}, 424 elliptical BGGs (56.5%) and 216 spiral BGGs (28.8%) in Control_{4C}, and 21 elliptical BGGs (37.5%) and 26 spiral BGGs (46.4%) in RG₄. The p -values obtained with two-sided Fisher exact test are 1.00 against Control_{4B}, 0.88 against Control_{4C} and 0.03 against RG₄. Thus, BGG morphologies differ between CG₄ and RG₄: CG₄ BGGs have a higher elliptical fraction than RG₄ BGGs.

3. sSFR

3.1. Main sequence

We model the star-forming main sequence as a function of stellar mass using orders 1–4 (Fig. 9). We then use order 2 to compute the residual of each star-forming galaxy with respect to the main sequence, and show their distributions in Fig. 10.

The median main-sequence residual of CG₄ galaxies is:

- significantly different from that of Control_{4B}, with a median offset of $\Delta = 0.056$ ($0.029 \leq \Delta \leq 0.099$, bootstrap), corresponding to a p -value of $p = 0.051$.
- significantly different from that of Control_{4C}, with a median offset of $\Delta = 0.082$ ($0.062 \leq \Delta \leq 0.122$, bootstrap), corresponding to a p -value of $p = 0.010$.
- not significantly different from that of RG₄, with a median offset of $\Delta = 0.046$ ($-0.035 \leq \Delta \leq 0.099$, bootstrap), corresponding to a p -value of $p = 0.535$.

A direct rank-sum comparison of the star-forming main-sequence offsets only identifies significant differences between CG₄ and

Control_{4C}, whose median offset is -0.060 with respect to the fitted sequence ($p = 0.03$).

Within CG₄, the star-forming main-sequence offsets of the Embedded, Isolated and Predom classes remain statistically consistent with one another.

3.2. Distance to the BGG

We also compare the sSFR of satellite galaxies with their distance to the BGG, expressed both in projected kpc and in units of $\langle R_{ij} \rangle$.

Significant monotonic trends are detected in the following cases.

In CG₄, sSFR correlates with projected distance to the BGG with Spearman coefficient $\rho_s = 0.24$ for 177 satellites




Fig. 6. Distributions of various quantities for galaxies according to their fertility.

($p = 1.2 \times 10^{-3}$) and Pearson coefficient $r = 0.21$ ($p = 4.1 \times 10^{-3}$).

In Control_{4B}, sSFR correlates with distance to the BGG normalized by $\langle R_{ij} \rangle$ with Spearman coefficient $\rho_s = 0.08$ for 1991 satellites ($p = 3.6 \times 10^{-4}$) and Pearson coefficient $r = 0.08$ ($p = 2.3 \times 10^{-4}$).

In Control_{4C}, sSFR correlates with projected distance to the BGG with Spearman coefficient $\rho_s = 0.25$ for 2143 satellites ($p = < 10^{-6}$) and Pearson coefficient $r = 0.20$ ($p = < 10^{-6}$).

In Control_{4C}, sSFR correlates with distance to the BGG normalized by $\langle R_{ij} \rangle$ with Spearman coefficient $\rho_s = 0.10$ for 2143 satellites ($p = 4.6 \times 10^{-6}$) and Pearson coefficient $r = 0.09$ ($p = 1.8 \times 10^{-5}$).

Sample	$\log_{10}(\text{sSFR} [\text{yr}^{-1}])$		M_r		$\log(M_\star/M_\odot)$	
	Passive	Star-forming	Passive	Star-forming	Passive	Star-forming
CG4	-12.08 ± 0.03	-10.19 ± 0.07	-21.16 ± 0.14	-20.4 ± 0.14	10.64 ± 0.07	10.05 ± 0.05
Control4B	-12.1 ± 0.01	-10.31 ± 0.02	-21.52 ± 0.04	-20.75 ± 0.04	10.79 ± 0.01	10.24 ± 0.02
Control4C	-11.92 ± 0.02	-10.16 ± 0.02	-20.81 ± 0.06	-19.93 ± 0.04	10.47 ± 0.03	9.82 ± 0.03
RG4	-11.91 ± 0.07	-10.16 ± 0.07	-21.11 ± 0.25	-20.32 ± 0.12	10.62 ± 0.09	9.97 ± 0.09

Table 3. Median values of galaxy properties for passive and star-forming populations in each sample. Uncertainties are estimated by bootstrapping (10 000 iterations).

Sample BGG	$\log_{10}(\text{sSFR} [\text{yr}^{-1}])$		M_r		$\log(M_\star/M_\odot)$	
	Passive	Star-forming	Passive	Star-forming	Passive	Star-forming
CG4	-11.53 ± 0.09	-10.85 ± 0.30	-20.93 ± 0.11	-20.77 ± 0.32	10.46 ± 0.06	10.32 ± 0.12
Control4B	-11.53 ± 0.03	-10.86 ± 0.06	-21.25 ± 0.03	-21.12 ± 0.07	10.61 ± 0.02	10.46 ± 0.04
Control4C	-11.3 ± 0.03	-10.62 ± 0.06	-20.42 ± 0.05	-20.55 ± 0.08	10.22 ± 0.02	10.15 ± 0.06
RG4	-11.14 ± 0.19	-10.58 ± 0.17	-20.69 ± 0.14	-20.74 ± 0.27	10.32 ± 0.07	10.19 ± 0.13

Table 4. Median values of various quantities for galaxies according to the fertility of their BGG (BGG included). Uncertainties are estimated by bootstrapping (10 000 iterations).

Sample	$\log_{10}(\text{sSFR} [\text{yr}^{-1}])$		M_r		$\log(M_\star/M_\odot)$	
	Passive	Star-forming	Passive	Star-forming	Passive	Star-forming
CG4	-11.23 ± 0.16	-10.92 ± 0.48	-20.45 ± 0.12	-20.34 ± 0.18	10.22 ± 0.05	10.09 ± 0.07
Control4B	-11.26 ± 0.06	-10.91 ± 0.13	-20.9 ± 0.03	-20.75 ± 0.05	10.43 ± 0.02	10.3 ± 0.03
Control4C	-10.96 ± 0.05	-10.6 ± 0.07	-19.8 ± 0.04	-20.02 ± 0.10	9.91 ± 0.02	9.89 ± 0.07
RG4	-10.82 ± 0.19	-10.61 ± 0.22	-20.23 ± 0.17	-20.3 ± 0.13	10.09 ± 0.08	10 ± 0.13

Table 5. Median values of various quantities for galaxies according to the fertility of their BGG (satellites only). Uncertainties are estimated by bootstrapping (10 000 iterations).

3.3. Dominance

We split groups into dominated systems, where the BGG contributes more than 60% of the group luminosity, and non-dominated systems. This split does not produce a strong compact-group signal by itself: CG₄ shows no significant compact/non-dominated difference after the Benjamini–Hochberg false-discovery-rate (FDR) correction. Here p_{adj} denotes the FDR-adjusted p-value, and Cliff’s δ is a rank-based effect size ranging from -1 to 1.

Across the control and reference samples, the clearest binary separation is instead in group luminosity. Dominated systems have lower L_{group} than their non-dominated counterparts: Control_{4B} gives 9.1×10^{10} versus $1.28 \times 10^{11} L_\odot$ ($p_{\text{adj}} < 10^{-6}$, $\delta = -0.55$); Control_{4C} gives 8.49×10^{10} versus $9.54 \times 10^{10} L_\odot$ ($p_{\text{adj}} < 10^{-6}$, $\delta = -0.25$); and RG₄ gives 7.09×10^{10} versus $9.82 \times 10^{10} L_\odot$ ($p_{\text{adj}} = 7.8 \times 10^{-5}$, $\delta = -0.59$). The negative δ values mean that dominated groups tend to occupy the lower-luminosity side of each comparison.

The spiral-fraction correlations show a broader environmental pattern rather than a clean dominated/non-dominated dichotomy. The most coherent set appears in Control_{4C}, where S_{frac} increases with $\langle R_{ij} \rangle$ in both dominated and non-dominated groups ($\rho_s = 0.38$, $p < 10^{-6}$ and $\rho_s = 0.35$, $p < 10^{-6}$), while weaker negative trends are found with σ_v and L_{group} . Control_{4B} contributes only weaker non-dominated-group correlations. In CG₄, the significant trends occur in small subsets, with S_{frac} decreasing with L_{group} among dominated groups ($\rho_s = -0.50$, 36 groups) and with $\Delta V_{\text{BGG}}/\sigma_v$ among non-dominated groups

($\rho_s = -0.53$, 26 groups); we therefore treat these as supporting diagnostics rather than the main dominance result. The full distributional and correlation diagnostics are shown in Figs. 13 and 14.

3.4. Morphology and domination

When we further split the samples by domination and by the morphology of the BGG, significant effects only appear in

Control_{4B} dominated groups, where the median satellite spiral fraction is 1.00 for spiral BGGs and 0.67 for elliptical BGGs. The corresponding group-level odds ratio is 1.84 ($p = 0.01$), while the satellite-level clustered fit gives an odds ratio of 1.84 ($p = 0.02$);

Control_{4C} dominated groups, where the median satellite spiral fraction is 0.67 for spiral BGGs and 0.50 for elliptical BGGs. The corresponding group-level odds ratio is 1.84 ($p = 2.1 \times 10^{-5}$), while the satellite-level clustered fit gives an odds ratio of 1.84 ($p = 2.7 \times 10^{-5}$).

4. Group dynamics

The crossing-time diagnostic mainly separates systems by dynamical scale rather than by luminosity alone. Shorter t_{cross} systems have larger virial mass-to-light ratios and virial masses: the correlations with $\log(M_{\text{VT}}/L_r)$ and $\log(M_{\text{VT}})$ are $\rho_s = -0.39$ ($p < 10^{-6}$) and $\rho_s = -0.33$ ($p < 10^{-6}$), respectively, over 816 groups. By contrast, the relation with total group luminosity is formally significant but very weak




Fig. 7. Distributions of various quantities for galaxies according to the fertility of their BGG (BGG included).




Fig. 8. Distributions of various quantities for galaxies according to the fertility of their BGG (satellites only).

($\rho_s = 0.08$, $p = 0.03$). We therefore read t_{cross} as a dynamical compactness indicator tied primarily to virial mass and mass-to-light ratio, with little independent leverage from L_{group} . Figures 15–17 show the corresponding projected and three-dimensional trends.

5. Colour

We derive the $(u - r)$, $(u - g)$, $(g - r)$ and $(r - i)$ colours from the matched SDSS photometry. The retained matches include 110 of 248 galaxies in CG₄, 931 of 2792 galaxies in Control_{4B}, 1478 of 3004 galaxies in Control_{4C}, 102 of 224 galaxies in RG₄. Stellar mass is a confounding variable in colour comparisons (Alpaslan et al. 2015); we therefore model each colour jointly with $\log(M_*/M_\odot)$ rather than comparing unadjusted colour distributions.

5.1. Catalogue colour–mass relations

The hypothesis of a common colour–mass slope across catalogues is rejected for $u-r$ ($p = 4.96 \times 10^{-4}$), $u-g$ ($p = 0.00203$), $g-r$ ($p = 1.46 \times 10^{-4}$), $r-i$ ($p = 0.00495$). After Holm correction for the directional comparisons, significantly steeper CG₄ slopes are found for $u-r$ relative to Control_{4C}, with $\Delta b = 0.289$ ($p_{\text{Holm}} = 0.00163$); for $u-r$ relative to SDSS, with




Fig. 9. Polynomial fits of orders 1–4 to the star-forming main sequence.

$\Delta b = 0.232$ ($p_{\text{Holm}} = 0.00163$); for $u-g$ relative to Control_{4C}, with $\Delta b = 0.237$ ($p_{\text{Holm}} = 0.00287$); for $u-g$ relative to SDSS, with $\Delta b = 0.186$ ($p_{\text{Holm}} = 0.00165$). These relations are shown in Fig. 18.




Fig. 10. Main-sequence residual distributions for the star-forming galaxies in each sample.




Fig. 11. sSFR as a function of projected distance to the BGG, in kpc, for satellite galaxies in each sample.

5.2. Satellite colours

To isolate an environmental difference, we pool satellites from the control catalogues as the ordinary-group population and compare them with CG₄ satellites at the common reference mass $\log(M_\star/M_\odot) = 9.885$. At this mass, compact-group satellites are 0.124 mag bluer in u-r ($p_{\text{Holm}} = 0.00909$) and 0.106 mag bluer in u-g ($p_{\text{Holm}} = 0.00737$). The compact and ordinary colour-mass slopes also differ in u-r ($b_{\text{compact}} = 0.819$, $b_{\text{ordinary}} = 0.344$, $p_{\text{Holm}} = 2.21 \times 10^{-5}$), u-g ($b_{\text{compact}} = 0.532$, $b_{\text{ordinary}} = 0.150$, $p_{\text{Holm}} = 1.42 \times 10^{-4}$), g-r ($b_{\text{compact}} = 0.287$, $b_{\text{ordinary}} = 0.195$, $p_{\text{Holm}} = 2.9 \times 10^{-4}$). Figure 19 displays the corresponding fitted relations.

5.3. Robustness of the satellite colour result

The SDSS-photometry subset is not representative of the full galaxy catalogues: matched galaxies are systemati-




Fig. 12. sSFR as a function of projected distance to the BGG normalized by $\langle R_{ij} \rangle$ for satellite galaxies in each sample.




Fig. 13. Distributions of the main group-scale observables in dominated and non-dominated systems.

cally less massive, more actively star-forming and more frequently satellites than unmatched galaxies. The fractions with all four colours are 44.4% in CG₄, 33.3% in Control4B, 49.2% in Control4C, 45.5% in RG₄.

We therefore refit the satellite colours with cluster-robust standard errors, using catalogue-qualified group identifiers. In the pooled comparison adjusted only for stellar mass and redshift, none of the four CG₄ coefficients is significantly negative after Holm correction. After additionally controlling for morphology and sSFR class, negative coefficients appear in ($u-r$) ($\Delta = -0.127$ mag, $p_{\text{Holm}} = 0.0177$), ($u-g$) ($\Delta = -0.101$ mag, $p_{\text{Holm}} = 0.0271$). These adjusted coefficients are control-sample dependent: the significant in-




Fig. 14. Bootstrap distributions of the Spearman coefficients for the group-property pairs selected by the XOR domination criterion.




Fig. 15. Crossing time versus virial mass-to-light ratio for the compact groups and their controls.

dividual comparisons are Control4C in $(u-r)$, Control4C in $(u-g)$, Control4C in $(g-r)$.

Within sSFR classes, no negative CG₄ residual difference survives correction for the four colours. Likewise, no within-morphology residual difference survives the same correction. Thus, the data do not localize the colour effect to the star-forming spiral population expected for a simple interaction-triggered star-formation scenario.

Figure 20 summarizes the model and control-sample dependence. Taken together, these tests do not establish a robust global blue offset for CG₄ satellites; the negative full-model coefficients should instead be treated as a control-sensitive exploratory result.




Fig. 16. Crossing time versus virial mass for the compact groups and their controls.




Fig. 17. Three-dimensional view of the crossing time, virial mass and group luminosity space.

5.4. Radial and group-level colour trends

The residual colours become bluer at larger projected distance from the BGG in CG₄. After Holm correction within each distance metric, significant CG₄ correlations are found for $(u-r)$ ($\rho_S = -0.379$, $p_{\text{Holm}} = 2.6 \times 10^{-4}$), for $(u-g)$ ($\rho_S = -0.336$, $p_{\text{Holm}} = 0.0011$), for $(g-r)$ ($\rho_S = -0.257$, $p_{\text{Holm}} = 0.00922$). This radial trend is consistent with stronger processing near the group centre and a less processed outer population, rather than centrally triggered blue colours.

At the group level, crossing time correlates with mean satellite $\Delta(u-r)$ ($\rho_S = -0.418$, $p_{\text{Holm}} = 0.013$), median satellite $\Delta(u-r)$ ($\rho_S = -0.409$, $p_{\text{Holm}} = 0.0167$), satellite blue fraction ($\rho_S = 0.369$, $p_{\text{Holm}} = 0.0393$). These group-level re-




Fig. 18. Optical colour as a function of stellar mass for CG₄, the three control catalogues and the matched SDSS reference sample. Dashed lines show the fitted catalogue relations.

lations remain exploratory because many groups contribute only a small number of colour-matched satellites.

any of the four colours. Figure 22 shows these group-level relations.

5.5. BGG and satellite colours

The mean satellite colour varies with the BGG colour across the combined group sample: for u-r, $r_p = 0.360$ ($p_{\text{Holm}} = 7.91 \times 10^{-7}$) and $\rho_S = 0.321$ ($p_{\text{Holm}} = 1.3 \times 10^{-5}$); for u-g, $r_p = 0.326$ ($p_{\text{Holm}} = 6.01 \times 10^{-6}$) and $\rho_S = 0.325$ ($p_{\text{Holm}} = 1.3 \times 10^{-5}$); for g-r, $r_p = 0.340$ ($p_{\text{Holm}} = 2.94 \times 10^{-6}$) and $\rho_S = 0.283$ ($p_{\text{Holm}} = 1.13 \times 10^{-4}$); for r-i, $r_p = 0.142$ ($p_{\text{Holm}} = 0.0464$) and $\rho_S = 0.232$ ($p_{\text{Holm}} = 0.00103$). After Holm correction, the BGG-satellite colour slope in CG₄ does not differ significantly from any control catalogue in

Taken together, these results suggest a limited degree of BGG-satellite concordance. The direct evidence is clearest for colour: redder BGGs are associated with redder mean satellite populations. A similar tendency appears in the satellite-only split by BGG sSFR class in Table 5, where satellites around passive BGGs have lower median sSFR than those around star-forming BGGs. The morphology signal is more restricted, appearing only in dominated control systems. We therefore treat concordance as a secondary pattern in the present analysis rather than as a separate population-level classification.




Fig. 19. Satellite colour–mass relations in compact and ordinary groups. The fitted lines show both colour offsets at the reference mass and differences in slope.

5.6. BGG colours and luminosity domination

The BGG stellar-mass distributions are significantly different between the domination classes ($p = 7.32 \times 10^{-4}$); the median $\log(M_{\star}/M_{\odot})$ values are 11.092 for dominated and 10.976 for non-dominated BGGs. Before mass adjustment, a significant colour difference is found in g-r ($\Delta_{\text{median}} = 0.021$, $p_{\text{Holm}} = 0.00157$). At the common reference mass $\log(M_{\star}/M_{\odot}) = 11.042$, the adjusted results are u-r: $\Delta = -0.002$ mag ($p_{\text{Holm}} = 1$); u-g: $\Delta = -0.010$ mag ($p_{\text{Holm}} = 1$); g-r: $\Delta = 0.007$ mag ($p_{\text{Holm}} = 1$); r-i: $\Delta = 0.001$ mag ($p_{\text{Holm}} = 1$). Thus, no dominated/non-dominated BGG colour offset remains significant after controlling for stellar mass. The mass dependence and class fits are shown in Fig. 23.

6. What drives the compact-group differences?

The previous sections establish that CG₄ galaxies differ from the control catalogues in their population mix, especially in morphology and in the star-forming fraction of satellites. We now ask whether these differences survive after controlling for stellar mass, rank and available group-scale quantities, and whether they are better interpreted as compact-group-specific evolution, preprocessing in dense halo cores, or catalogue-selection effects.

6.1. Mass-, rank- and environment-adjusted galaxy properties

We fit binomial models to the combined galaxy catalogues, standardizing the continuous predictors and using group-clustered standard errors. The common adjustment set in-




Fig. 20. Robustness of the satellite colour coefficients. Left: pooled ordinary-group comparison before and after controlling for morphology and sSFR class. Right: mass- and redshift-adjusted comparisons against each control catalogue separately. Points show the CG₄ minus ordinary coefficient and bars show 95% confidence intervals; negative values indicate bluer CG₄ satellites.




Fig. 21. Mass- and redshift-adjusted satellite $(u-r)$ and $(u-g)$ residuals as a function of projected distance to the BGG. Lines connect binned medians for CG₄ and the pooled ordinary-group satellites; extreme plotting tails are omitted for readability, while the reported rank correlations use the full matched sample. More negative residuals are bluer than the ordinary-satellite reference relation.

cludes stellar mass, redshift, satellite/BGG rank, group luminosity and velocity dispersion when sufficiently complete.

For passive status, the CG₄ odds ratio is 1.20, with 95% confidence interval [0.88, 1.63] and $p_{\text{Holm}} = 0.485$.

For passive satellite status, the CG₄ odds ratio is 1.31, with 95% confidence interval [0.95, 1.80] and $p_{\text{Holm}} = 0.390$.

For star-forming satellite status, the CG₄ odds ratio is 0.76, with 95% confidence interval [0.56, 1.05] and $p_{\text{Holm}} = 0.390$.

For elliptical morphology, the CG₄ odds ratio is 1.62, with 95% confidence interval [1.21, 2.16] and $p_{\text{Holm}} = 0.007$.




Fig. 22. Mean satellite colour versus BGG colour for groups with matched photometry. The dotted line is the one-to-one relation and dashed lines are catalogue-specific fits.

For elliptical BGG morphology, the CG₄ odds ratio is 0.90, with 95% confidence interval [0.46, 1.74] and $p_{\text{Holm}} = 0.751$.

For spiral morphology, the CG₄ odds ratio is 0.62, with 95% confidence interval [0.46, 0.83] and $p_{\text{Holm}} = 0.007$.

Compact-group membership therefore remains independently associated with morphology after the available adjustments: CG₄ galaxies have a higher probability of being elliptical and a lower probability of being spiral. In contrast, the adjusted passive and star-forming satellite coefficients are weaker and do not provide evidence for an instantaneous star-formation difference that remains significant after correction.

6.2. Matched-control comparison

We matched 234 CG₄ galaxies one-to-one to 234 ordinary-group galaxies without replacement, requiring common propensity-score support, a 0.2-standard-deviation caliper and exact rank strata. The propensity model uses logMstar, z numeric, rank, log group luminosity, velocity dispersion. Projected distance is retained as a phase-space and interaction diagnostic rather than matched away because it partly defines compactness and has limited overlap. The largest absolute standardized mean difference changed from 0.28 before matching to 0.14 afterwards.

The matched CG₄-minus-control difference in passive fraction is 0.098, with 95% bootstrap interval [0.017, 0.175] and $p_{\text{Holm}} = 0.054$.




Fig. 23. BGG colour–mass relations split by luminosity domination. Comparing the fitted relations at a common stellar mass separates colour effects from the different mass distributions.

This is suggestive but not formally significant after correction, so we treat the passive-fraction excess as borderline.

The corresponding CG₄–minus–control difference in star-forming fraction is -0.098 , with 95% bootstrap interval $[-0.175, -0.017]$ and $p_{\text{Holm}} = 0.054$.

The matched CG₄–minus–control difference in elliptical fraction is 0.197 , with 95% bootstrap interval $[0.090, 0.303]$ and $p_{\text{Holm}} < 10^{-3}$.

The matched CG₄–minus–control difference in spiral fraction is -0.197 , with 95% bootstrap interval $[-0.303, -0.090]$ and $p_{\text{Holm}} < 10^{-3}$.

Among matched star-forming galaxies, the residual-sSFR difference is -0.096 , with 95% bootstrap interval $[-0.221, 0.015]$ and $p = 0.084$.

The matched comparison is therefore consistent with the adjusted models: the most robust compact-group signal is

morphological, while the passive and star-forming-fraction signals are more model-dependent.

6.3. Satellite segregation around the brightest group galaxy

The previous sections show global differences between CG₄ and RG₄ satellites. We now ask whether these differences are concentrated near the BGG, as expected for accelerated evolution in the dense inner group, or whether they persist across the satellite population. The analysis uses 177 CG₄ and 166 RG₄ satellites, excludes rank-1 galaxies, ranks satellites by projected BGG-centric distance, and measures passive and early-type fractions with group-cluster bootstrap uncertainties. When available, we also use $|\Delta v_{\text{BGG}}|/\sigma_v$, with $\Delta v_{\text{BGG}} = c(z_{\text{gal}} - z_{\text{BGG}})/(1 + z_{\text{group}})$, as a projected




Fig. 24. Adjusted CG₄ odds ratios. Bars show 95% confidence intervals; the vertical line marks equal odds.




Fig. 25. CG₄-minus-control effects after one-to-one matching. Bars show 95% bootstrap confidence intervals.




Fig. 26. Absolute standardized mean differences before and after matching.

phase-space diagnostic. Similar projected phase-space coordinates are often used as observational proxies for satellite infall state or quenching time, but only statistically (Mahajan et al. 2011; Oman & Hudson 2016; Wetzel et al. 2013).

In the innermost satellite-distance bin, the passive fractions are 68.9% for CG₄ and 48.2% for RG₄, a CG₄-RG₄ difference of 20.6 percentage points. The corresponding early-type fractions are 65.4% and 35.6%, with a difference of 29.8 percentage points.




Fig. 27. Passive satellite fraction as a function of projected-distance rank from the BGG. Points show CG₄ and RG₄ fractions; bars show group-cluster bootstrap 16th-84th percentile intervals.




Fig. 28. Early-type satellite fraction as a function of projected-distance rank from the BGG. Because the catalogue separates ellipticals, spirals, and uncertain morphologies but not S0 galaxies, early type is equivalent to elliptical in this diagnostic.

The passive-fraction contrast is similar between the inner and outer distance ranks, so the data do not isolate the CG₄ offset to the immediate BGG environment.

After adjusting for stellar mass, redshift, distance bin, and available group-scale controls, the passive-status model gives a CG₄ odds ratio of 1.27 (95% CI 0.47-3.46; $p = 0.640$).

The analogous early-type model gives a CG₄ odds ratio of 2.57 ($p = 0.058$).

A nearest-neighbour mass-redshift matched check uses 128 CG₄-RG₄ satellite pairs. The matched passive-fraction difference is 9.4 percentage points, and the matched early-type difference is 19.6 percentage points.

Projected phase-space coordinates should not be read as orbital reconstructions for individual galaxies. In compact groups the member counts are small, velocity dispersions are noisy, and projection mixes recently accreted and long-resident satellites. We therefore treat these bins as an observational diagnostic of where the CG₄-RG₄ population contrast appears, rather than as a direct dynamical clock.




Fig. 29. Passive satellite fraction in coarse projected phase-space bins. Inner and outer refer to projected-distance rank from the BGG; low and high velocity are split at $|\Delta v_{\text{BGG}}|/\sigma_v = 1$.




Fig. 30. Distributions of the first-second and first-fourth magnitude gaps in CG₄ and the pooled controls.

6.4. Magnitude gaps and fossil-like assembly

We define $\Delta m_{12} = M_{r,2} - M_{r,1}$ and $\Delta m_{14} = M_{r,4} - M_{r,1}$, so that a more dominant BGG has a positive, larger gap.

For Δm_{12} , the CG₄ and control medians are 1.17 and 1.09 mag, respectively ($p_{\text{Holm}} = 0.705$).

For Δm_{14} , the CG₄ and control medians are 2.56 and 2.59 mag, respectively ($p_{\text{Holm}} = 0.705$).

The compact-group morphology signal is therefore not accompanied by a significant excess in these simple fossil-like magnitude-gap indicators.

The correlation between Δm_{12} and passive satellite fraction is $\rho_S = -0.15$ ($p_{\text{Holm}} < 10^{-3}$).

6.5. Recently quenched candidates

The catalogues do not contain both D_n4000 and $H\delta_A$, so a post-starburst classification is not possible. We therefore perform only a limited $H\alpha$ -equivalent-width comparison, using the SDSS convention in which negative equivalent width denotes emission. We classify strong $H\alpha$ emission using $\text{h_alpha_eqw} \leq -3$ Angstrom.

The strong-emission fraction is 32.3% in CG₄, 41.0% in Control4B, 45.7% in Control4C, 56.9% in RG4.

Relative to Control4B, the CG₄ strong-emission fraction differs by -0.087 ($p_{\text{Holm}} = 0.012$).

Relative to Control4C, the CG₄ strong-emission fraction differs by -0.135 ($p_{\text{Holm}} < 10^{-3}$).

Relative to RG4, the CG₄ strong-emission fraction differs by -0.246 ($p_{\text{Holm}} < 10^{-3}$).

These limited $H\alpha$ results indicate that CG₄ galaxies have a lower strong-emission fraction than in each control




Fig. 31. Passive satellite fraction as a function of the first-second magnitude gap.




Fig. 32. $H\alpha$ equivalent-width distributions used for the limited recent-quenching diagnostic.

sample in the current catalogue, but they do not provide a direct post-starburst or recently-quenched classification.

6.6. Emission-line and AGN-like populations

The complementary BPT analysis classifies star-forming, composite and broad AGN-like populations. The CG₄ AGN-like fraction is 38.0%.

After mass and environment adjustment, the CG₄ AGN-like odds ratio is 1.14 ($p = 0.451$).

Thus, AGN-like activity does not provide a robust independent compact-group discriminator in this analysis.

6.7. Pairwise tidal index

The pairwise analysis reconstructs projected separations and velocity differences for 6268 galaxies and tests whether adding $\sum_j M_{\star,j}/R_{ij}^3$ changes the adjusted CG₄ coefficient.

For passive status, the baseline CG₄ odds ratio is 1.20 ($p = 0.227$), while the model including the tidal index gives a CG₄ odds ratio of 0.80 ($p = 0.170$). The tidal-index term itself has odds ratio 1.65 ($p < 10^{-3}$).

For elliptical morphology, the baseline CG₄ odds ratio is 1.65 ($p < 10^{-3}$), whereas the model including the tidal index




Fig. 33. AGN-like fractions from the BPT/AGN-flag classification.




Fig. 34. Passive status as a function of the reconstructed pairwise tidal index. Colours distinguish CG₄ from control galaxies.

gives a CG₄ odds ratio of 1.15 ($p = 0.346$). The tidal-index term has odds ratio 1.56 ($p < 10^{-3}$). This suggests that the reconstructed local tidal-density proxy absorbs much of the unadjusted CG₄ morphology coefficient, although the result remains an exploratory projected-pair diagnostic.

6.8. Selection diagnostics

The fractions with complete measurements for the major derived quantities are shown in Fig. 35. The colour-matched versus unmatched comparisons quantify shifts in stellar mass, redshift, rank, star-formation class and morphology.

The median nearest-neighbour angular separation is 100.9 arcsec in CG₄ and 209.0 arcsec in the controls; the fractions below the 55-arcsec SDSS fibre-collision scale are 19.4% and 9.3%, respectively.

No major availability difference meets both criteria of corrected significance and a difference exceeding 10 percentage points.

The automated availability row for group-scale quantities should be interpreted as an alias audit for a specific set of column names rather than as evidence that no group-scale information is used elsewhere in the analysis.

7. Relation to recent structural studies of compact-group galaxies

A complementary interpretation of the present results is provided by the recent series of studies by Montaguth et al. (2023, 2025, 2026b,a). Using multiwavelength S-PLUS imaging and single-Sérsic structural fits, these authors identified a population of “transition” galaxies, broadly corresponding to red systems with relatively low or intermediate Sérsic index, whose location in the R_e – n plane suggests ongoing structural transformation in compact groups. In particular, Montaguth et al. (2023) found that compact-group transition galaxies include a population that is smaller and more centrally concentrated than analogous field galaxies, while Montaguth et al. (2025) showed that this signature is stronger in non-isolated compact groups embedded in larger structures. Montaguth et al. (2026b) further compared non-isolated compact groups with their surrounding host-group galaxies and found that the compact-group members display higher quenched fractions, a larger early-type fraction, and distinct transition-galaxy (R_e – n) distributions. Finally, Montaguth et al. (2026a) showed that the environmental dependence of the stellar mass–size relation is weak for early- and late-type galaxies, but strongest for transition galaxies, whose mass–size slopes and Sérsic-index distributions become more early-type-like in groups and compact groups.

Our analysis addresses a related but stricter question. Rather than comparing compact groups primarily with the field, or non-isolated compact groups with their immediate host structures, we ask whether compact-group galaxies remain different from galaxies in regular four-member group analogues after controlling for stellar mass, redshift, rank, and available group-scale quantities. In this framework, the most robust residual signal is morphological: compact-group membership remains associated with an enhanced probability of being classified as elliptical and a reduced probability of being classified as spiral. By contrast, the passive and star-forming satellite fractions are more sensitive to the adopted control sample and to the adjustment or matching procedure.

The two sets of results are therefore not contradictory. The structural work of Montaguth et al. (2023, 2025, 2026b,a) identifies the likely nature of the transformation: disk fading, increasing central concentration, and/or bulge growth in galaxies occupying an intermediate region between blue late-type disks and red early-type systems. Our results indicate that, when compact groups are compared with regular-group analogues rather than with the field alone, the clearest surviving signature is the accumulated morphological state of the galaxy population, whereas an instantaneous star-formation offset is less robust. This suggests that compact groups should not be interpreted simply as environments with ongoing enhanced quenching at the present epoch. They are more plausibly very dense regions in which past interactions, local tidal density, and preprocessing in larger haloes have already shifted part of the galaxy population toward earlier morphologies.

A useful way to reconcile these views is to regard the compact-group signal as primarily structural and cumulative. In that picture, the compact configuration may accelerate morphological transformation, but some of the observed differences may also reflect the fact that many compact groups are the dense cores or substructures of larger




Fig. 35. Availability fractions for the major derived quantities in each sample.




Fig. 36. Stellar-mass distributions of galaxies with and without complete matched colour measurements.

both fractions. This morphological difference is also present among BGGs: CG₄ BGGs are more often elliptical than RG₄ BGGs.

The star-formation difference is more specific. CG₄ satellites are significantly less frequently star-forming than RG₄ satellites, while the star-forming fraction of BGGs does not differ significantly between the two samples. Moreover, among galaxies that remain star-forming, CG₄ galaxies do not show a significant offset from RG₄ galaxies relative to the star-forming main sequence.

The colour analysis does not indicate a simple uniform reddening of compact-group galaxies. The pooled mass- and redshift-adjusted comparison does not support a statistically robust blue CG₄ satellite offset. Negative coefficients appear after adding morphology and sSFR class, but the result is sensitive to the choice of control catalogue. Neither the sSFR-class nor morphology splits isolate a significant effect among star-forming spirals. The radial residual trends instead indicate that outer CG₄ satellites are bluer than inner satellites, consistent with stronger processing toward the BGG. The BGG and mean satellite colours are correlated, but the corresponding CG₄ relations do not differ significantly from those of the control catalogues. Together with the satellite-only split by BGG sSFR class and the domination-conditioned morphology result, this suggests limited BGG–satellite concordance rather than a separately established general concordance signal. Likewise, the apparent colour difference between dominated and non-dominated BGGs disappears after controlling for stellar mass.

8. Conclusion

After removing systems classified as split, we find that compact groups differ from regular four-member groups mainly through their galaxy population mix. Galaxies in CG₄ are more frequently elliptical and less frequently spiral than galaxies in RG₄, with highly significant differences in

The most robust compact-group signal is morphological rather than an instantaneous star-formation offset. After controlling for stellar mass, redshift, rank and available group-scale properties, CG₄ membership remains associated with enhanced elliptical and reduced spiral probabilities, whereas passive status and satellite star-forming

status do not remain significant in the corresponding adjusted models. The matched-control analysis supports the same morphological excess and gives only borderline evidence for an enhanced passive fraction.

The projected phase-space test shows a passive-fraction contrast of similar size at inner and outer satellite ranks. This comparison is necessarily projected: with only a few satellites per group, $|\Delta v|/\sigma_v$ and projected radius cannot recover individual orbits, so the result should be read as a localization test for the population difference rather than as a dynamical reconstruction.

The additional diagnostics do not favour a simple fossil-group or AGN-driven explanation. The magnitude-gap distributions of CG₄ and the pooled controls are statistically similar, and AGN-like activity is not independently enhanced after mass and environment adjustment. The limited H α diagnostic shows a lower strong-emission fraction in CG₄ than in the control catalogues, but the absence of both D_n4000 and H δ_A prevents a direct post-starburst or recently-quenched classification.

Taken together, these results suggest that compact groups host a more morphologically evolved galaxy population, with an enhanced fraction of early-type galaxies and a reduced fraction of star-forming satellites. Because the passive CG₄ signal does not remain significant in both the adjusted and matched analyses, the apparent star-formation specialness is at least partly explained by environment, mass/rank mix, or pre-processing in dense halo cores. The morphology signal is more robust, but the projected tidal-index experiment suggests that local pairwise density may absorb part of that signal, so this result should be interpreted as evidence for accumulated transformation in very dense environments rather than as a uniquely isolated compact-group effect.

The selection audit does not remove all catalogue-systematic uncertainty, and the angular-neighbour diagnostic confirms the expected higher close-pair incidence in CG₄, but it does not identify a major availability imbalance under the adopted criterion.

Interpolation points for sSFR classification

References

- Alpaslan, M., Driver, S., Robotham, A. S. G., et al. 2015, *MNRAS*, 451, 3249
- Bishop, C. M. 2006, *Pattern Recognition and Machine Learning* (Springer)
- Bisigello, L., Caputi, K. I., Grogin, N., & Koekemoer, A. 2018, *Astron. Astrophys.*, 609, A82
- Dempster, A. P., Laird, N. M., & Rubin, D. B. 1977, *J. R. Statist. Soc. B*, 39, 1
- Díaz-Giménez, E., Zandivarez, A., & Taverna, A. 2018, *A&A*, 618, A157
- Hahn, C., Starkenburg, T. K., Choi, E., et al. 2019, *Astrophys. J.*, 872
- Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003, *Monthly Notices of the Royal Astronomical Society*, 346, 1055
- Kullback, S. & Leibler, R. A. 1951, *Ann. Math. Statist.*, 22, 79
- Lim, S. H., Mo, H. J., Lu, Y., Wang, H., & Yang, X. 2017, *MNRAS*, 470, 2982
- Lintott, C., Schawinski, K., Bamford, S., et al. 2011, *MNRAS*, 410, 166
- Mahajan, S., Mamon, G. A., & Raychaudhury, S. 2011, *MNRAS*, 416, 2882
- McLachlan, G. & Peel, D. 2000, *Finite Mixture Models* (Wiley)
- Montaguth, G. P., Mendes de Oliveira, C., Lima-Dias, C., et al. 2026a, *The Astrophysical Journal*, 999, 160
- Montaguth, G. P., Monachesi, A., Torres-Flores, S., et al. 2025, *Astronomy & Astrophysics*, 696, A240
- Montaguth, G. P., O’Mill, A. L., Mendes de Oliveira, C., et al. 2026b, *The Astrophysical Journal*, 998, 91
- Montaguth, G. P., Torres-Flores, S., Monachesi, A., et al. 2023, *Monthly Notices of the Royal Astronomical Society*, 524, 5340–5356
- Oman, K. A. & Hudson, M. J. 2016, *MNRAS*, 463, 3083
- Pedregosa, F., Varoquaux, G., Gramfort, A., et al. 2011, *J. Mach. Learn. Res.*, 12, 2825
- Rousseeuw, P. J. 1987, *Comput. Appl. Math.*, 20, 53
- Tempel, E., Tuvikene, T., Kipper, R., & Libeskind, N. I. 2017, *A&A*, 602, A100
- Tojeiro, R., Heavens, A. F., Jimenez, R., & Panter, B. 2009, *Mon. Not. R. Astron. Soc.*, 397, 1263
- Tricottet, M., Mamon, G. A., & Díaz-Giménez, E. 2025, *A&A*, 699, A329
- Veilleux, S. & Osterbrock, D. E. 1987, *Astrophysical Journal Supplement Series*, 63, 295
- Wetzel, A. R., Tinker, J. L., Conroy, C., & van den Bosch, F. C. 2013, *MNRAS*, 432, 336
- Wuyts, S., Förster Schreiber, N. M., Lutz, D., et al. 2011, *Astrophys. J.*, 742, 96
- Zheng, Y.-L. & Shen, S.-Y. 2021, *ApJ*, 911, 105