Octinoxate — co-fold MD binding energy (retention)

Peptide	Boltz / BoltzGen co-fol	OpenFold3 co-fold	RoseTTAFold3 co-fold
orig_f12	−13.71 (77.3%)	−5.90 (7.4%)	−19.51 (99.9%)
s2_esm2_control	−16.25 (59.6%)	−10.21 (33.9%)	−14.98 (57.2%)
s3_esm2_f4	−21.08 (99.6%)	−13.76 (84.8%)	−16.18 (88.2%)
s3_orig_f12	−24.33 (100.0%)	−10.10 (44.8%)	−12.10 (68.9%)
shuffle_control	−15.13 (75.9%)	−9.39 (25.7%)	−11.13 (38.7%)
shuffle_control_esm0	−14.32 (80.2%)	−15.62 (92.5%)	−21.53 (99.3%)
bg33_1	−11.31 (14.5%)	−14.90 (56.4%)	−12.43 (68.3%)
bg33_2	−8.02 (50.9%)	−11.58 (70.2%)	−11.52 (58.7%)
bg33_3	−11.86 (41.0%)	−10.10 (14.9%)	−12.44 (1.9%)
bg33_4	−19.66 (100.0%)	−17.11 (93.0%)	−16.13 (81.2%)

ΔG = MM/GBSA binding free energy (kcal/mol) over the whole 20 ns run; retention (in parentheses) = % of the run with the ligand within 10 Å of the peptide. Boltz/BoltzGen is the original co-fold; BoltzGen for bg* peptides, Boltz for the rest.

Oxybenzone — co-fold MD binding energy (retention)

Peptide	Boltz / BoltzGen co-fol	OpenFold3 co-fold	RoseTTAFold3 co-fold
ox1_esm1_f8	−10.70 (75.0%)	−17.95 (100.0%)	−12.12 (35.2%)
ox1_orig_f12	−10.31 (49.1%)	−3.96 (20.5%)	−12.27 (89.1%)
ox2_esm1_f8	−20.81 (100.0%)	−16.68 (93.2%)	−10.90 (90.4%)
ox2_orig_f8	−24.21 (100.0%)	−7.23 (49.5%)	−4.95 (19.6%)
ox2_shuffle	−5.30 (7.1%)	−2.98 (14.8%)	−13.35 (51.0%)
ox2_shuffle_esm0	−12.04 (94.7%)	−0.92 (1.1%)	−11.44 (32.6%)
ox3_orig_f8	−18.84 (100.0%)	−2.06 (2.4%)	−7.41 (45.6%)
bgox31_2	−10.89 (33.7%)	−7.45 (65.9%)	−9.06 (84.6%)
bgox31_3	−15.44 (99.7%)	−5.82 (15.8%)	−14.36 (99.2%)
bgox31_4	−14.05 (100.0%)	−5.74 (17.9%)	−11.02 (72.0%)
bgox31_5	−16.78 (100.0%)	−11.55 (70.7%)	−14.69 (96.4%)

ΔG = MM/GBSA binding free energy (kcal/mol) over the whole 20 ns run; retention (in parentheses) = % of the run with the ligand within 10 Å of the peptide. Boltz/BoltzGen is the original co-fold; BoltzGen for bg* peptides, Boltz for the rest.