

Protein’s geometry

Centre of mass

$$\vec{R}_{cm} = \frac{\sum_{i=1}^N m_i \vec{r}_i}{\sum_{i=1}^N m_i}$$

Radius of gyration

$$r_g = \sqrt{\frac{\sum_{i=1}^N m_i (\vec{r}_i - \vec{R}_{cm})^2}{\sum_{i=1}^N m_i}}$$

$$RMSD(t) = \sqrt{\frac{1}{N} \sum_{i=1}^N (\vec{r}_i(t) - \vec{r}_i(0))^2}$$

$$RMSF_i = \sqrt{\langle \Delta r_i^2 \rangle} = \sqrt{\frac{1}{M} \sum_{f=1}^M (\vec{r}_{i,f} - \langle \vec{r}_i \rangle)^2}$$

$$B_i = \frac{8\pi^2}{3} RMSF_i^2$$

Semi-empirical force fields

Bond stretching

Harmonic

$$U(r_{AB}) = \frac{1}{2} k_{AB} (r_{AB} - r_{AB,eq})^2$$

Anarmonic

$$U(r_{AB}) = \frac{1}{2} \left[k_{AB} + k_{AB}^{(3)} (r_{AB} - r_{AB,eq}) \right] (r_{AB} - r_{AB,eq})^2$$

Quartic correction

$$U(r_{AB}) = \frac{1}{2} \left[k_{AB} + k_{AB}^{(3)} (r_{AB} - r_{AB,eq}) + k_{AB}^{(4)} (r_{AB} - r_{AB,eq})^2 \right] \cdot (r_{AB} - r_{AB,eq})^2$$

Morse

$$U(r_{AB}) = D_{AB} \left[1 - e^{-\alpha_{AB} (r_{AB} - r_{AB,eq}^*)} \right]^2$$

Valence angle bending

Potential

$$U(\theta_{ABC}) = \frac{1}{2} [k_{ABC} + k_{ABC}^{(3)} (\theta_{ABC} - \theta_{ABC,eq}) + k_{ABC}^{(4)} (\theta_{ABC} - \theta_{ABC,eq})^2 + \dots] (\theta_{ABC} - \theta_{ABC,eq})^2$$

Fourier

$$U(\theta_{ABC}) = \sum_{\{j\}_{ABC}} k_{j,ABC}^{fourier} [1 + \cos(j\theta_{ABC} + \psi_j)]$$

Fourier

$$k_{j,ABC}^{fourier} = \frac{2k_{ABC}^{harmonic}}{j^2}$$

Torsions

Potential

$$U(\omega_{ABCD}) = \frac{1}{2} \sum_{\{j\}_{ABCD}} V_{j,ABCD} [1 + (-1)^{j+1} \cos(j\omega_{ABCD} + \psi_{j,ABCD})]$$

Improper

$$U(\omega_{ABCD}) = \frac{1}{2} \sum_{\{j\}_{ABCD}} V_{j,ABCD} [1 + (-1)^{j+1} \cos(j\omega_{ABCD} + \psi_{j,ABCD})]$$

Van der Waals

Lennard-Jones

$$U(r_{AB}) = 4\epsilon_{AB} \left[\left(\frac{\sigma_{AB}}{r_{AB}} \right)^{12} - \left(\frac{\sigma_{AB}}{r_{AB}} \right)^6 \right]$$

Morse

$$U(r_{AB}) = D_{AB} \left[1 - e^{-\alpha_{AB} (r_{AB} - r_{AB,eq}^*)} \right]^2$$

Hill

$$U(r_{AB}) = \epsilon \left[\frac{6}{\beta_{AB}-6} e^{\beta_{AB} \frac{1-r_{AB}}{r_{AB}}} - \frac{\beta_{AB}}{\beta_{AB}-6} \left(\frac{r_{AB}}{r_{AB}} \right)^6 \right]$$

Electrostatic interactions

Distribution of charges

$$U_{AB} = \sum_A \sum_{B>A} \vec{M}^{(A)} \vec{V}^{(B)}$$

Point like

$$U_{AB} = \frac{q_A q_B}{\epsilon_{AB} r_{AB}}$$

Dipolar interactions

$$U_{AB/CD} = \frac{\mu_{AB} \mu_{CD}}{\epsilon_{AB/CD} r_{AB/CD}^3} (\cos \chi_{AB/CD} - 3 \cos \alpha_{AB} \cos \alpha_{CD})$$

Parameterization

Parameters

$$Z = \sqrt{\sum_i \frac{observable\ occurrences}{j} \frac{(calc_{i,j} - expt_{i,j})^2}{w_i^2}}$$

$$\sigma_{AB} = \sigma_A + \sigma_B \qquad \epsilon_{AB} = \sqrt{\epsilon_A \epsilon_B}$$

Classical mechanics (contd)

Legendre transforms

$$s = f'(x) \equiv g(x) \qquad f'(x) = g(x) = s \Rightarrow x = g^{-1}(s)$$

$$b(g^{-1}(s)) = f(g^{-1}(s)) - sg^{-1}(s) \equiv \tilde{f}(s) = f(x(s)) - sx(s)$$

$$\tilde{f}(s_1, \dots, s_n) = f(x_1(s_1, \dots, s_n), \dots, x_n(s_1, \dots, s_n)) - \sum_i s_i x_i(s_1, \dots, s_n)$$

Hamiltonian formulation

$$\mathcal{H}(\vec{r}_1, \dots, \vec{r}_N, \vec{p}_1, \dots, \vec{p}_N) = -\tilde{\mathcal{L}}(\vec{r}_1, \dots, \vec{r}_N, \vec{p}_1, \dots, \vec{p}_N)$$

$$\mathcal{H}(\vec{r}_1, \dots, \vec{r}_N, \vec{p}_1, \dots, \vec{p}_N) = \sum_{i=1}^N \frac{\vec{p}_i^2}{2m_i} + U(\vec{r}_1, \dots, \vec{r}_N)$$

$$\mathcal{H}(q_1, \dots, q_{3N}, p_1, \dots, p_{3N}) = \frac{1}{2} \sum_{\alpha} \sum_{\beta} p_{\alpha} G_{\alpha\beta}^{-1} p_{\beta} + U(q_1, \dots, q_{3N})$$

Hamilton equations

$$\dot{q}_{\alpha} = \frac{\partial \mathcal{H}}{\partial p_{\alpha}} \qquad \dot{p}_{\alpha} = -\frac{\partial \mathcal{H}}{\partial q_{\alpha}} \qquad \frac{\mathcal{H}}{dt} = 0 \qquad \mathcal{H} = const$$

Some properties

Conservation laws

$$\frac{da}{dt} = \frac{\partial a}{\partial x_i} \dot{x}(t) = \{a, \mathcal{H}\} = 0$$

Incompressibility

$$\nabla_x(x) = 0$$

Symplectic structure

$$M = J^T M J \qquad J_{kl} = \frac{\partial x_k(t)}{\partial x_l(0)}$$

Theoretical foundations of statistical mechanics

Thermodynamics

Equilibrium

$$g(N, P, V, T) = 0$$

State function

$$f(n, P, V, T)$$

Reversible work

$$dW_{rev} = -PdV + \mu dN$$

Heat

$$dQ_{rev} = CdT$$

First law

$$\Delta E = \Delta Q + \Delta W$$

Entropy

$$\Delta S = \int_1^2 \frac{dQ_{rev}}{T}$$

The ensemble

Average

$$A = \frac{1}{Z} \sum_{\lambda=1}^N a(x_{\lambda}) \equiv \langle a \rangle$$

Microstate

$$x_0 = (q_1(0), \dots, q_{3N}(0), p_1(0), \dots, p_{3N}(0))$$

Phase space volume

$$dx_t = J(x_t; x_0) dx_0 \qquad \frac{dJ}{dt} = 0 \Rightarrow J(x_t; x_0) = 1 \Rightarrow dx_t = dx_0$$

$$f(x_t) : \int f(x) dx = 1 \wedge \frac{df(x_t, t)}{dt} = 0 \Rightarrow$$

Distribution function

$$f(x_t, t) dx_t = f(x_0, 0) dx_0 \Rightarrow$$

$$\frac{\partial f(x, t)}{\partial t} + \{f(x, t), \mathcal{H}(x, t)\} = 0$$

Equilibrium

$$A = \int a(x) f(x, t) dx \Rightarrow \frac{\partial f(x, t)}{\partial t} = 0 \wedge \{f(x, t), \mathcal{H}(x, t) = 0\} \Rightarrow$$

$$f(x) \propto \mathcal{F}(\mathcal{H}(x))$$

$$Z = \int dx \mathcal{F}(\mathcal{H}(x)) \Rightarrow f(x) = \frac{1}{Z} \mathcal{F}(\mathcal{H}(x))$$

Introduction to molecular dynamics

Verlet algorithm

$$\vec{r}_i(t + \Delta t) = 2\vec{r}_i(t) - \vec{r}_i(t - \Delta t) + \frac{\Delta t^2}{m_i} \vec{F}_i(t)$$

$$\vec{v}_i(t + \Delta t) = \vec{v}_i(t) + \frac{\Delta t}{2m_i} \left[\vec{F}_i(t) + \vec{F}_i(t + \Delta t) \right]$$

Initial conditions

$$f(v) = \sqrt{\frac{m}{2\pi kT}} e^{-\frac{mv^2}{2kT}} \qquad f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{x^2}{2\sigma^2}}$$

Action integral

$$Q \equiv \{q_1, \dots, q_{3N}\} \qquad \dot{Q} \equiv \{\dot{q}_1, \dots, \dot{q}_{3N}\}$$

$$A[Q] = \int_{t_1}^{t_2} \mathcal{L}(Q(t), \dot{Q}(t)) dt$$

$$\delta Q(t_1) = \delta Q(t_2) = 0 \qquad \delta \dot{Q}(t_1) = \delta \dot{Q}(t_2) = 0$$

$$\delta A = \int_{\alpha=1}^{3N} \frac{\partial \mathcal{L}}{\partial \dot{q}_{\alpha}} \delta q_{\alpha}(t) \Big|_{t_1}^{t_2} dt + \int_{t_1}^{t_2} \sum_{\alpha=1}^{3N} \left[\frac{\partial \mathcal{L}}{\partial q_{\alpha}} \delta q_{\alpha}(t) - \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}_{\alpha}} \right) \delta q_{\alpha}(t) \right] dt = 0$$

Constraints

$$\sum_{\alpha=1}^{3N} a_{k\alpha} dq_{\alpha} + a_{kt} dt = 0, k = 1, \dots, N_C$$

Holonomic

$$a_{k\alpha} = \frac{\partial \sigma_k}{\partial q_{\alpha}} \qquad a_{kt} = \frac{\partial \sigma_k}{\partial t}$$

$$\frac{1}{2} \sum_i m_i \dot{r}_i^2 - C = 0 \Rightarrow \frac{1}{2} \sum_i m_i \dot{r}_i d\vec{r}_i - C dt = 0$$

Non-holonomic

$$\Rightarrow a_{1i} = \frac{1}{2} m_i \dot{r}_i \wedge a_{1t} = -C$$

Lagrange multiplier

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}_{\alpha}} \right) - \frac{\partial \mathcal{L}}{\partial q_{\alpha}} = \sum_{k=1}^{N_C} \lambda_k a_{k\alpha}$$

$$\dot{q}_{\alpha} = \frac{\partial \mathcal{H}}{\partial p_{\alpha}} \qquad \dot{p}_{\alpha} = -\frac{\partial \mathcal{H}}{\partial q_{\alpha}} - \sum_{k=1}^{N_C} \lambda_k a_{k\alpha} \qquad \sum_{\alpha=1}^{3N} a_{k\alpha} \frac{\partial \mathcal{H}}{\partial p_{\alpha}} = 0$$

Classical mechanics

Newton’s laws

$$\vec{F} = m\vec{a} \qquad \vec{F}_{BA} = -\vec{F}_{AB}$$

$$\vec{v}(t) = \frac{d\vec{r}}{dt} \qquad \vec{a}(t) = \frac{d\vec{v}}{dt} = \frac{d^2\vec{r}}{dt^2} \qquad m \frac{d^2\vec{r}}{dt^2} = \vec{F}$$

Force acting on atom

$$\vec{F}_i(\vec{r}_1, \dots, \vec{r}_N, \vec{r}_i) = \sum_{j \neq i} \vec{F}_{ij}(\vec{r}_i - \vec{r}_j) + \vec{F}^{(ext)}(\vec{r}_i, \dot{\vec{r}}_i)$$

Bond stretching:

$$U = \frac{k_l}{2} (l - l^0)^2$$

Bond bending:

$$U = \frac{k_{\theta}}{2} (\theta - \theta^0)^2$$

Bond torsion:

$$U = k_{\phi} [1 + \cos(n\phi - \phi^0)]$$

Van der Waals interactions:

$$U = \left[\frac{a_{ij}}{r_{ij}^{12}} - \frac{b_{ij}}{r_{ij}^6} \right]$$

Electrostatic interactions:

$$U = \frac{332 q_i q_j}{\epsilon r_{ij}}$$

$$\vec{p}_i = m_i \vec{v}_i = m_i \dot{\vec{r}}_i \qquad \vec{F}_i = m_i \ddot{\vec{r}}_i = \dot{\vec{p}}_i$$

$$\vec{x}(t) = \{\vec{r}_1(t), \dots, \vec{r}_N(t), \vec{p}_1(t), \dots, \vec{p}_N(t)\}$$

Lagrangian formulation

$$\vec{F}_i(\vec{r}_1, \dots, \vec{r}_N) = -\Delta_i U(\vec{r}_1, \dots, \vec{r}_N)$$

$$W_{AB} = \int_A^B \vec{F}_i d\vec{l} = U_A - U_B = -\Delta U_{AB} \qquad \oint \vec{F}_i d\vec{l} = 0$$

Kinetic energy

$$K(\dot{\vec{r}}_1, \dots, \dot{\vec{r}}_N) = \frac{1}{2} \sum_i m_i \dot{\vec{r}}_i^2$$

$$\mathcal{L}(\vec{r}_1, \dots, \vec{r}_N, \dot{\vec{r}}_1, \dots, \dot{\vec{r}}_N) = K(\dot{\vec{r}}, \dot{\vec{r}}_N) - U(\vec{r}_1, \dots, \vec{r}_N)$$

Euler-Lagrange

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\vec{r}}_i} \right) - \frac{\partial \mathcal{L}}{\partial \vec{r}_i} = 0$$

$$E = \frac{1}{2} \sum_i m_i \dot{\vec{r}}_i^2 + U(\vec{r}_1, \dots, \vec{r}_N)$$

$$quad \frac{dE}{dt} = 0$$

Generalized coordinates

$$q_{\alpha} = f_{\alpha}(\vec{r}_1, \dots, \vec{r}_N) \qquad \alpha = 1, \dots, 3N \qquad \vec{r}_i = \vec{q}_i(q_1, q_2, \dots, q_{3N}) \qquad i = 1, \dots, N$$

$$\dot{\vec{r}}_i = \sum_{\alpha=1}^{3N} \frac{\partial \vec{r}_i}{\partial q_{\alpha}} \dot{q}_{\alpha} \qquad \mathcal{L}(q, \dot{q}) = \frac{1}{2} \sum_{\alpha=1}^{3N} \sum_{\beta=1}^{3N} G_{\alpha\beta} \dot{q}_{\alpha} \dot{q}_{\beta} - U(q_1, \dots, q_{3N})$$

Microcanonical ensemble

State and distribution function

State function

$$dS = \frac{1}{T} dE + \frac{P}{T} dV - \frac{\mu}{T} dN$$

$$\left(\frac{\partial S}{\partial E} \right)_{V, N} = \frac{1}{T} \qquad \left(\frac{\partial S}{\partial V} \right)_{N, E} = \frac{P}{T} \qquad \left(\frac{\partial S}{\partial N} \right)_{V, N} = \frac{\mu}{T}$$

Boltzmann relation

$$S(N, V, E) = k \ln \omega(N, V, E)$$

$$\Omega(N, V, E) = M_N \int d\vec{p} \int_{D(V)} d\vec{r} \delta(\mathcal{H}(\vec{r}, \vec{p}) - E)$$

Distribution function

$$= M_N \int dx \delta(\mathcal{H}(x) - E)$$

$$M_N = \frac{E_0}{N! h^{3N}}$$

$$A = \langle a \rangle = \frac{M_N}{\Omega(N, V, E)} \int dxa(x) \delta(\mathcal{H}(x) - E) = \frac{\int dxa(x) \delta(\mathcal{H}(x) - E)}{\int dx \delta(\mathcal{H} - E)}$$

$$\left\langle x_i \frac{\partial \mathcal{H}}{\partial x_j} \right\rangle = \frac{M_N}{\Omega(N, V, E)} \frac{\partial}{\partial E} \int_{\mathcal{H}(x) < E} dx x_i \frac{\partial (\mathcal{H} - E)}{\partial x_j}$$

Virial theorem

$$\left\langle x_i \frac{\partial \mathcal{H}}{\partial x_j} \right\rangle = \delta_{ij} \frac{\Sigma(E)}{\frac{1}{\partial E}}$$

$$\Sigma(N, V, E) = \frac{1}{N! h^{3N}} \int dx \theta(E - \mathcal{H})$$

$$\Omega(N, V, E) = E_0 \frac{\partial \Sigma(N, V, E)}{\partial E} \quad \left\langle x_i \frac{\partial \mathcal{H}}{\partial x_j} \right\rangle = \delta_{ij} \left(\ln \frac{\Sigma(E)}{\partial E} \right)^{-1}$$

$$S(N, V, E) = k \ln \Omega(N, V, E) \simeq k \ln \Sigma(N, V, E) = \tilde{S}(N, V, E)$$

$$\left\langle x_i \frac{\partial \mathcal{H}}{\partial x_j} \right\rangle \simeq \delta_{ij} \left(\frac{S(E)}{\partial E} \right)^{-1} = kT \delta_{ij}$$

Thermal contact

$$\Omega(N, V, E) = M_N \int dx \delta(\mathcal{H}_1(x_1) + \mathcal{H}_2(x_2) - E)$$

$$\Omega(N, V, E) = \int dE_1 \Omega_1(N_1, V_1, E_1) \Omega_2(N_2, V_2, E - E_1)$$

$$S(N, V, E) = k \ln \Omega_1(N_1, V_1, \bar{E}_1) + k \ln \Omega_2(N_2, V_2, E - \bar{E}_1)$$

$$= S_1(N_1, V_1, \bar{E}_1) + S_2(N_2, V_2, E - \bar{E}_1)$$

$$T_1 = T_2$$