



Working Report

Benchmark WKMSYSPiCT Meeting

Stochastic Surplus Production Model in Continuous Time (SPiCT) Explorations in Nephrops FU30 Stock (Gulf of Cadiz)

Instituto Español de Oceanografía (IEO)

Authors

Yolanda Vila
María José Zuñiga
Mauricio Mardones

Copenhagen, Denmark
February 8, 2025

Contents

Abstract	2
Introduction	3
Methods	4
Study Area	4
SPiCT description	4
Data Sources	5
Previous Nephrops FU30 Scenarios	7
New Scenarios in Nephrops FU30	11
Results	15
Previous Analysis	15
New Scenarios	15
Comparative Population variables	16
Biological References Ponits.	16
Management Outputs	17
Conclusions	18
Code Repositories	19
Figures	20
Tables	65
References	73

Abstract

Nephrops stock in FU30 Division 9.a is considered a Category 3 stock. The advice is currently based on the 2-over-3 rule using the abundance index from UWTV surveys. The reference points for this stock are undefined and ICES cannot assess the stock and exploitation status relative to MSY or precautionary approach (PA) reference points. Stochastic surplus production model in continuous time (SPiCT) has been recommended by ICES for a number of data limited stocks to produce MSY advice. This WD shows some SPiCT explorations carried out in order to evaluate the potential of this model for *Nephrops* FU 30 assessment and propose this stock as possible candidate for the next WKMSYSPICT benchmark. Results obtained are promising.

Introduction

This report extends the exploratory framework by incorporating new SPiCT scenarios that emerged from discussions held after the last meeting (WKMSYSPiCT) held in December 2025. Specifically, a new set of scenarios is introduced, all configured consistently with the previous analyses by applying the same alternative prior configurations (RUN5–RUN8) to each scenario. These new explorations include five additional scenarios (SC5–SC9) that differ in the combination of fishery-dependent and fishery-independent information used to describe stock dynamics. *Scenario 5 (SC5)* combines total landings with the ISUNEPCA UWTV abundance index and the normalized ARSA yield series, providing a configuration based exclusively on survey-derived indicators to assess consistency between independent data sources. *Scenario 6 (SC6)* integrates total landings with the ISUNEPCA UWTV abundance index and a standardised commercial LPUE, allowing direct comparison between fishery-independent abundance signals and fishery-dependent information. *Scenario 7 (SC7)* represents the most information-rich configuration, combining total landings with ISUNEPCA UWTV abundance, normalized ARSA yield, and standardised LPUE indices to explore model behaviour under maximum data availability. *Scenario 8 (SC8)* integrates total landings with a normalized ISUNEPCA UWTV biomass index and standardised LPUE, providing an alternative representation of stock dynamics based on biomass rather than abundance. Finally, *Scenario 9 (SC9)* combines total landings with ISUNEPCA UWTV abundance, ARSA yield, and standardised LPUE, while explicitly accounting for a structural break in the ARSA survey by restricting its use to the pre-2012 period and relying on complementary indices thereafter.

The document provides a structured synthesis of these new scenarios and presents their main results, highlighting their relative performance, strengths, and limitations in comparison with previously explored configurations, with the aim of informing future benchmark discussions and potential model selection for *Nephrops norvegicus* in Functional Unit 30.

Methods

Study Area

The Gulf of Cádiz (FU30, Division 9.a) is located in the southwestern part of the Iberian Peninsula, bordered to the north by the Portuguese coast and to the south by the Strait of Gibraltar (Figure 1). The area is characterised by a wide continental shelf that extends from the coast to depths of approximately 200 m, followed by a steep slope descending to depths exceeding 1 000 m. The seabed is predominantly composed of sandy and muddy sediments, which provide suitable habitats for *Nephrops norvegicus*.

SPiCT description

Surplus productions models are widely used in fisheries stock assessment, particularly for data-limited stocks. These models describe the dynamics of the total biomass of a fish stock over time, considering the balance between growth (production) and removals (catches). The basic equation governing surplus production models is:

$$\frac{dB(t)}{dt} = P(B(t)) - C(t)$$

where: - $B(t)$ is the biomass at time t , - $P(B(t))$ is the surplus production function, representing the net growth of the biomass, - $C(t)$ is the catch at time t .

In a graphical inspection, a surplus production curve typically has a dome-shaped form, indicating that production increases with biomass up to a certain point (the maximum sustainable yield, MSY), after which it declines due to density-dependent effects. The shape of curve depends on the specific functional form chosen for $P(B)$, such as the Schaefer model, Pella-Tomlinson model, or others. (Figure 5).

The SPiCT model extends traditional surplus production models by incorporating stochastic elements and continuous-time dynamics. This allows for a more realistic representation of the uncertainties inherent in fish stock assessments, such as environmental variability and observation errors. The key features of the SPiCT model include: - **Stochasticity**: The model incorporates process and observation errors, allowing for the quantification of uncertainty in biomass estimates and reference points. - **Continuous-time framework**: Unlike discrete-time models, SPiCT operates in continuous time, providing a more flexible and accurate representation of stock dynamics. - **Multiple data sources**: SPiCT can integrate various types of data, including catch data, abundance indices from surveys, and fishery-dependent indices such as CPUE and fishing effort. - **Bayesian inference**: The

model employs Bayesian methods for parameter estimation, enabling the incorporation of prior knowledge and the derivation of posterior distributions for model parameters and stock status indicators.

Data Sources

Annual Nephrops **landings** in FU30 show strong interannual variability over the analysed period (Figure 2), with increasing catches during the late 1990s and early 2000s followed by a sharp decline associated with management measures. Since then, landings have remained at lower levels with notable recent fluctuations, reflecting both stock availability and regulatory constraints.

Fishery-independent information is provided by the **ISUNEPCA UWTV survey**, conducted annually since 2014 (except 2020), which delivers absolute abundance and biomass estimates following ICES standards (ICES, 2017, 2022; Vila & Burgos, 2022; Vila, Burgos, & Soriano, 2016), and by the **ARSA bottom trawl surveys**, which provide biomass indices for Nephrops in deeper strata of FU30 (Figure ??). In addition, a **standardised commercial CPUE series**, derived from Spanish and Portuguese fleets, offers a fishery-dependent index of relative abundance.

Together, these complementary fishery-independent and fishery-dependent indices (Figure ?)(fig:indices)) capture contrasting but informative signals on stock dynamics and constitute the empirical basis for the SPiCT model analyses.

Pairwise relationships among the different abundance, productivity and fishery-dependent indices were explored using a Spearman rank correlation analysis. Spearman correlations were selected to account for potential non-linear relationships and non-normal distributions in the time series.

The summary of the input data series used for SPiCT explorations for *Nephrops norvegicus* in Functional Unit 30 is presented in Table 1. The data sources include:

- **Total catches** (1987–2024) (Figure 2).
- **Fishing effort** (1994–2024), expressed as total fishing days.
- **Standardised landings per unit of effort (CPUE)** (2009–2025).
- **ISUNEPCA UWTV survey indices**, including:
 - absolute **abundance index** (available from 2015–2025), and
 - associated **biomass index** (available for selected years).
- **ARSA survey indices**, comprising:
 - **spring survey index** (1993–2025) (Figure 3).

All time series were harmonised temporally and formatted consistently prior to model fitting. Alternative combinations of these data sources were used to define the different SPiCT scenarios explored in this study.

Table 1: Summary of input data series used for SPiCT explorations for *Nephrops norvegicus* in Functional Unit 30. Column abbreviations are as follows: CATC = total catches (t); ARSB = ARSA biomass survey index; ARSP = ARSA productivity index; ARSS = standardised ARSA index; ARCV = coefficient of variation of the ARSA index; ISUB = ISUNEPCA UWTV biomass index; ISUA = ISUNEPCA UWTV abundance index; EFRT = fishing effort (days); and CPUE = standardised landings per unit of effort.

Year	CATC	ARSB	ARSP	ARSS	ARCV	ISUB	ISUA	EFRT	CPUE
1987	193.0000	-	-	-	-	-	-	-	-
1988	138.6700	-	-	-	-	-	-	-	-
1989	174.5000	-	-	-	-	-	-	-	-
1990	220.1000	-	-	-	-	-	-	-	-
1991	225.4010	-	-	-	-	-	-	-	-
1992	242.7600	-	-	-	-	-	-	-	-
1993	159.7000	26.603058	0.9876871	0.7798737	0.4086136	-	-	-	-
1994	107.6180	28.159856	-	-	-	-	-	-	-
1995	130.6230	-	-	-	-	-	-	27823	98.6
1996	48.5220	24.615855	0.5861900	0.4628532	0.4826584	-	-	32194	99.4
1997	97.0760	9.328451	0.3670690	0.2898361	0.5870258	-	-	31951	88.2
1998	85.3150	9.301283	0.1918494	0.1514835	0.5395427	-	-	32573	79.2
1999	120.2170	-	0.5861043	0.4627855	0.5601319	-	-	32824	62.2
2000	128.8620	10.140252	0.5449645	0.4303017	0.4379313	-	-	32731	66.1
2001	178.4260	12.331178	0.6088606	0.4807538	0.4192129	-	-	29875	60.6
2002	261.8360	10.871560	0.8496977	0.6709178	0.5063920	-	-	30416	67.7
2003	306.5170	-	-	-	-	-	-	31760	74.0
2004	147.0209	4.240238	0.3958709	0.3125780	0.5853713	-	-	32790	78.4
2005	246.1790	17.429929	0.9419806	0.7437840	0.5060772	-	-	30901	42.5
2006	245.3679	12.350958	0.6834708	0.5396657	0.4569831	-	-	33203	52.7
2007	214.4429	10.541994	0.5994218	0.4733010	0.4438644	-	-	26946	63.5
2008	119.7816	23.930662	1.0795141	0.8523799	0.4538919	-	-	25366	58.8
2009	119.2902	13.137462	0.4270744	0.3372162	0.5366998	-	-	20602	59.5
2010	107.3343	-	1.3660372	1.0786173	0.5920415	-	-	21401	50.0
2011	95.9099	6.974094	0.5952163	0.4699803	0.6090513	-	-	21352	45.6
2012	116.0000	5.038396	0.3248393	0.2564917	0.6688189	-	-	19986	54.2
2013	26.7170	23.155208	0.4620234	0.3648117	0.5242873	-	-	20402	58.0
2014	14.8850	48.654413	1.5304473	1.2084349	0.5224972	-	-	23099	92.1
2015	24.6410	33.141152	2.2382149	1.7672853	0.4915001	2460.417	79.9	-	40.1

Table 1: Summary of input data series used for SPiCT explorations for *Nephrops norvegicus* in Functional Unit 30. Column abbreviations are as follows: CATC = total catches (t); ARSB = ARSA biomass survey index (*continued*)

Year	CATC	ARSB	ARSP	ARSS	ARCV	ISUB	ISUA	EFRT	CPUE
2016	123.7210	68.015128	3.9783875	3.1413183	0.5056249	2327.798	100.3	-	58.8
2017	139.5160	40.933017	1.8935208	1.4951162	0.5874185	1806.237	77.3	22458	64.6
2018	75.5510	9.263278	0.6023305	0.4755977	0.4070041	2354.634	100.2	21583	45.5
2019	64.7700	32.037073	2.5669860	2.0268816	0.5479344	3567.028	109.9	20748	47.9
2020	62.8690	165.458399	6.9161498	5.4609633	0.4487297	-	-	20553	73.7
2021	49.2760	-	-	-	-	2601.158	66.2	19991	59.0
2022	44.4150	6.716924	0.3288110	0.2596278	0.6889084	1627.816	53.1	20443	48.7
2023	32.8630	17.925767	0.7688194	0.6070566	0.4523832	1591.873	54.2	18853	54.4
2024	32.0470	102.613914	3.0396395	2.4000868	0.4301205	1778.028	63.7	19671	47.3
2025	-	-	-	-	-	1114.548	38.0	19240	45.7

Previous Nephrops FU30 Scenarios

Several SPiCT model configurations were explored for Nephrops FU30, varying in the inclusion of abundance indices, catch data, and model parameterisations (priors). This exploratory framework is consistent with the approach adopted in previous ICES Nephrops working groups, where alternative data configurations and prior assumptions have been systematically tested to evaluate model robustness and suitability for assessment purposes. The full set of SPiCT model configurations explored in this study is summarised in Table 2, including the input data used in each scenario and the corresponding prior configurations applied consistently across scenarios. This structured exploration allows for a transparent comparison of model behaviour and performance, in line with recent methodological developments discussed within ICES Nephrops expert groups.

The **Scenario 0** included catch data from 1994 to 2023 and the ISUNEPCA UWTV survey index from 2015 to 2023, reflecting the core data sources currently used in the assessment framework for this stock. Additional scenarios were developed to assess the effect of incorporating supplementary fishery-independent and fishery-dependent information, such as the ARSA survey index (**Scenario 2**), as well as alternative indicators derived from effort and CPUE series. Further scenarios explored the sensitivity of model outcomes to different assumptions regarding process error, observation error, and prior distributions for key biological parameters, following configurations commonly examined in recent Nephrops assessments (**Scenarios 3 and 4**).

Explorations were conducted using data available up to 2025. Tables below summarise the different scenarios and model configurations tested. Fishing effort and CPUE were not included in any scenario, as survey-based indices are recommended when available. The autumn ARSA survey (SpGFS-caut-WIBTS-Q4) was also excluded from the analyses, as part of the stock is not accessible to the gear during this survey due to the reproductive behaviour of Nephrops. In November, ovigerous females remain inside their burrows and are therefore unavailable to bottom trawl sampling. For this reason, the autumn ARSA survey index has never been included in the FU30 WGBIE assessments.

The ISUNEPCA UWTW survey index is considered the most reliable abundance index for Nephrops in FU30 and is currently used as the basis for assessment and advice (ICES, 2023). This survey is conducted in the year of the assessment; for example, the UWTW survey planned for June 2024 will be used to inform the advice for 2025, delivered in October 2024. If the 2024 ISUNEPCA index were to be included in a SPiCT model, catch data for 2024 would not yet be available. This issue should be explicitly addressed if SPiCT is adopted as the assessment model for this stock in a future benchmark.

Table 2: Input data components, scenario-level descriptions, and associated time spans used in SPiCT model configurations for *Nephrops norvegicus* in Functional Unit 30.

Scenario	Scenario description	Input data component	Time span
SC0	Baseline configuration using only the UWTW survey as absolute abundance index.	Total catches	1987–2024
		ISUNEPCA UWTW abundance index	2015–2025
SC1	Configuration combining UWTW and ARSA surveys to assess consistency between fishery-independent indices.	Total catches	1987–2024
		ISUNEPCA UWTW abundance index	2015–2025
		ARSA spring survey index	1993–2025
SC2	Configuration including fishery-dependent information through directed commercial CPUE.	Total catches	1987–2024
		ISUNEPCA UWTW abundance index	2015–2025
		Directed commercial CPUE	1994–2025
SC3	Configuration using directed fishing effort as a proxy of stock dynamics.	Total catches	1987–2024
		ISUNEPCA UWTW abundance index	2015–2025
		Directed fishing effort	1994–2024
SC4	Configuration testing the use of standardized CPUE instead of raw commercial indices.	Total catches	1987–2024
		ISUNEPCA UWTW abundance index	2015–2025
		Old Standardised commercial CPUE	1994–2025

Across all SPiCT scenarios, four alternative prior configurations (RUN1–RUN4) were ex-

plored to evaluate model sensitivity to assumptions on stock depletion in previous scenarios, production curve shape, and intrinsic population growth. The baseline run (RUN1) relied on default SPiCT priors, while subsequent configurations progressively introduced informative priors on initial depletion (B/K), the production curve shape parameter (n), and the intrinsic growth rate (r). This hierarchical design ensured consistent and comparable evaluation of prior influence across scenarios (Table 3).

Table 3: Summary of prior configurations (RUNs) explored for the SPiCT model applied to *Nephrops norvegicus* in Functional Unit 30 in last meeting.

Run	Prior configuration	Specification	Purpose
RUN1	Default (package defaults)	Default SPiCT priors	Baseline configuration using default SPiCT assumptions.
RUN2	logbkfrac	$\log\text{bkfrac} \sim \log(0.5)$, $\text{SD} = 0.2$	Constrains initial depletion (B/K) around an intermediate level with low uncertainty.
RUN3	logbkfrac, logn	$\log\text{bkfrac} \sim \log(0.5)$, $\text{SD} = 0.2$; $\log n \sim \log(2)$, $\text{SD} = 0.5$	Adds a tighter prior on the production curve shape parameter (n).
RUN4	logbkfrac, logn, logr	$\log\text{bkfrac} \sim \log(0.5)$, $\text{SD} = 0.2$; $\log n \sim \log(2)$, $\text{SD} = 0.5$; $\log r \sim \log(0.2)$, $\text{SD} = 0.2$	Fully informative configuration including priors on depletion, production curve shape and intrinsic growth rate.

New Scenarios in Nephrops FU30

Based on previous experts recommendations based on knowledge about surveys and fishery of *Nephrops norvegicus*, **five additional modelling scenarios (Scenarios 5–9)** were defined, differing in the combination and temporal coverage of fishery-dependent and fishery-independent data sources (Table 4). Across all scenarios, **total landings** are included as the catch time series, while scenarios differ in the configuration of biomass- and abundance-based indices.

Scenario 5 combines total landings with **normalized biomass indices from the ISUNEPCA UWTV survey and the ARSA survey**, representing a configuration based exclusively on fishery-independent information. *Scenario 6* integrates total landings with the **normalized ISUNEPCA UWTV biomass index** and a **standardised commercial CPUE**, allowing comparison between survey-derived biomass signals and fishery-dependent abundance information. *Scenario 7* represents the most information-rich configuration, combining total landings with **normalized ISUNEPCA UWTV biomass**, **normalized ARSA biomass**, and **standardised CPUE**, thereby incorporating both fishery-independent and fishery-dependent signals across overlapping time periods. *Scenario 8* combines landings with the **normalized ISUNEPCA UWTV abundance index** and **standardised CPUE**, providing an alternative representation of stock dynamics based on abundance rather than biomass. Finally, *Scenario 9* was designed to explicitly account for a major structural change in the ARSA survey. Following a change in vessel and operational characteristics in 2012, only the **pre-2012 ARSA biomass series** is retained in this scenario to avoid potential methodological discontinuities. Post-2012 stock dynamics are instead informed by the **ISUNEPCA UWTV survey**, which provides a consistent biomass index from 2015 onwards, together with the **standardised CPUE** reflecting recent commercial fleet performance.

This scenario therefore provides a coherent data configuration that respects survey continuity while preserving both long-term and contemporary information for stock assessment (Table 4). All scenarios include total landings as the catch time series and differ in the configuration of abundance indices (Figure 3).

Table 4: Input data components, scenario-level descriptions, and associated time spans for extended SPiCT model configurations (SC5-SC9).

Scenario	Scenario description	Input data component	Time span
SC5	Scenario combining total landings with ISUNEPCA UWTW abundance and normalized ARSA yield indices to assess consistency between fishery-independent indices.	Total catches	1987–2024
		ISUNEPCA UWTW	2015–2025
		Abundance	
		ARSA yield	1993–2025
SC6	Scenario integrating total landings with ISUNEPCA UWTW abundance and a standardized LPUE index, combining fishery-independent and fishery-dependent information.	Total catches	1987–2024
		ISUNEPCA UWTW	2015–2025
		abundance	
		Standardised commercial LPUE	2009–2025
SC7	Most information-rich configuration, combining total landings with ISUNEPCA UWTW abundance, normalized ARSA yield, and standardized LPUE indices to explore model behaviour under maximum data availability.	Total catches	1987–2024
		ISUNEPCA UWTW	2015–2025
		abundance	
		ARSA yield	1993–2025
		Standardised commercial LPUE	2009–2025
SC8	Scenario integrating total landings with normalized ISUNEPCA UWTW biomass and a standardized LPUE index, combining fishery-independent and fishery-dependent information.	Total catches	1987–2024
		ISUNEPCA UWTW biomass (normalized)	2015–2025
		Standardised commercial LPUE	2009–2025
SC9	Scenario combines total landings with the ISUNEPCA UWTW biomass index, ARSA yield, and the standardised commercial LPUE, integrating long-term and recent information while ensuring consistency among data sources.	Total catches	1987–2024
		ISUNEPCA UWTW	2015–2025
		abundance	
		ARSA yield	1993–2012
		Standardised commercial LPUE	2009–2024

In light of the new observations arising from the initial analyses, an additional set of alternative prior configurations (**RUN5–RUN8**) was implemented to further test the robustness of the SPiCT scenarios. In these additional runs, the **bkfrac configuration was not used**, as preliminary results indicated that it strongly constrained model outcomes and could unduly condition estimates of stock status. Instead, RUN5–RUN8 focused on exploring alternative assumptions regarding the **production curve shape parameter (n)** and the **intrinsic growth rate (r)**, using a revised and more flexible prior structure.

In addition, **RUN6–RUN8 incorporated alternative priors on key process and observation error components**, allowing explicit evaluation of uncertainty assumptions beyond stock productivity parameters. Specifically, a prior was specified on the **catch process uncertainty** (logsdb), centred on a moderate level of variability, while additional priors

were introduced to **reduce uncertainty in catch observation error** (`logsd`) and **fish-ing mortality error** (`logdf`). These priors were intended to stabilise estimation of process and observation variance components and to assess the sensitivity of model outcomes to alternative assumptions regarding noise structure in the data-generating processes.

We further evaluated a subset of configurations (**RUN7**) in which **priors were specified on the observation model through an associated coefficient of variation (CV) on the abundance index**, allowing explicit testing of model sensitivity to assumptions about observation uncertainty. Finally, we explored the use of the default **SPiCT and priors** (**RUN9**), which define the shape and scale of the surplus production function and thus control the curvature and productivity of the production relationship. These and priors were considered as an alternative prior set but were ultimately **kept disabled** in the main analyses, as their inclusion was intended solely for diagnostic purposes rather than for defining the final model configurations. To generally stabilise estimation, SPiCT imposes default semi-informative priors on these quantities, preventing them from taking extreme and unrealistic values (Pedersen, Kokkalis, Mildenerger, & Berg, 2023).

All scenarios were also tested under an alternative prior assumption for the intrinsic population growth rate, fixing the prior on r at **0.5**, consistent with estimates reported in FishBase for *Nephrops norvegicus* FishBase. This additional configuration was used to evaluate the sensitivity of model outcomes to assumptions about population productivity.

Overall, this expanded hierarchical design enables a more comprehensive and systematic evaluation of prior influence across scenarios, while maintaining consistency and comparability among model configurations (Table ())(tab:spictpriors2)).

Table 5: Summary of additional prior configurations (RUN5–RUN9) explored for the SPiCT model applied to *Nephrops norvegicus* in Functional Unit 30.

Run	Prior configuration	Specification	Purpose
RUN5	Default SPiCT priors	All default semi-informative priors as implemented in SPiCT.	Baseline configuration using the default SPiCT prior structure to provide a reference for comparison.
RUN6	No prior on <code>logbkfrac</code> , <code>logn</code>	<code>logbkfrac</code> prior disabled; remaining priors unchanged prior about n .	Evaluates model sensitivity when initial depletion (B/K) is not constrained by an informative prior, but with production curve n parameter (Schaefer)
RUN7	Index CV prior	<code>logsd</code> prior specified for abundance index ($CV = 0.2$) using a normal prior on log-scale.	Tests sensitivity to explicit assumptions about observation uncertainty in the abundance index.
RUN8	Alpha–beta priors disabled	<code>logalpha</code> and <code>logbeta</code> priors disabled ($\logalpha = \logbeta = c(1, 1, 0)$).	Assesses the influence of SPiCT stabilising priors on the surplus production function when informative data are available.

To ensure consistency and comparability across scenarios, all new configurations are fitted using the same set of four prior specifications previously defined in the analysis. Results will be presented comparatively across scenarios, including diagnostic plots, key stock status outputs, and management-related figures and summary tables, allowing a coherent evaluation of model behavior and management implications under alternative data combinations. A total of **20 SPiCT model runs** were conducted, corresponding to four new alternative prior configurations (RUN5–RUN8) applied consistently across nine model scenarios (SC0–SC8). This factorial design allowed for the evaluation of model behaviour, convergence and sensitivity to both data configuration and prior assumptions on key biological and production parameters.

Results

Previous Analysis

The exploratory modelling scenarios SC0 to SC5 implemented in the present analysis are documented in detail in the technical report by Vila & Quinzán (2024), where alternative configurations of fishery-dependent and fishery-independent data were systematically evaluated using the SPiCT framework for *Nephrops norvegicus* in Functional Unit 30. These scenarios form part of the broader exploratory analyses conducted to assess model behaviour, data sensitivity, and the robustness of stock status estimates under different information assumptions.

The following sections present a synthesis of the key findings from these new scenarios, including diagnostic evaluations, stock dynamics, and management implications, while also referencing relevant results from the previously documented scenarios (SC0 to SC5) for comparative purposes.

New Scenarios

Correlation analysis of input data series The Figure 4 reveals strong positive correlations among indices derived from similar sources, particularly between ISUNEPCA UWTV biomass and abundance indices, as well as between standardized CPUE and effort-related metrics. In contrast, ARSA-based indices show weaker or negative correlations with some fishery-dependent indicators, highlighting differences in spatial coverage and survey design. Overall, the results indicate a clear clustering of indices by data origin, suggesting partial redundancy among some series and complementarity across others.

Population Variables In the new explorations, in this report we focus on the results obtained from the additional scenarios (SC5 to SC8) that incorporate alternative combinations of abundance indices corrected and fishery-dependent information.

To optimise the presentation and interpretation of model outputs for the SPiCT assessment of *Nephrops norvegicus* in Functional Unit 30, only scenarios and prior configurations that successfully converged are retained for reporting. For these converged runs, we present the key summary tables and diagnostic and management relevant figures, ensuring that all results shown are based on numerically stable and internally consistent model solutions. This approach facilitates a clear and robust comparison of alternative data configurations and prior assumptions while avoiding misleading inferences from non-convergent model fits.

The estimated biomass trajectories for *Nephrops* FU30 across scenarios SC5 to SC9 are

illustrated in Figures 7–8.

Hindcast Analysis The Mean Absolute Scaled Error (MASE) is a widely used metric for evaluating the predictive accuracy of time series models. It is defined as the mean absolute error of the model’s predictions, scaled by the mean absolute error of a naive forecast (typically the previous observation). MASE values below 1 indicate that the model outperforms the naive forecast, while values above 1 suggest inferior performance.

Hindcast diagnostics by scenario and prior configuration are presented in Figures 27 and 28–35, illustrating model predictive performance under alternative data and prior combinations.

Retrospective Analysis Retrospective patterns for each modelling scenario were evaluated under alternative prior configurations. Scenario-specific diagnostics are shown in Figures 9–26, illustrating the temporal stability of biomass and fishing mortality estimates across successive peelings. Retrospective bias was evaluated using Mohn’s rho estimates (Table 7). Threshold to consider a model as acceptable is set to $(\pm)0.20$ in mohn’s rho values for this kind of species (Hurtado-ferro et al., 2015).

The results indicate that all model runs achieved MASE values below 1, demonstrating that the SPiCT model consistently outperformed the naive forecast in predicting stock dynamics. Notably, RUN3 exhibited the lowest MASE value, suggesting that the inclusion of informative priors on both initial depletion and production curve shape enhanced model predictive accuracy. These findings underscore the importance of prior specification in improving model performance and provide confidence in the SPiCT model’s ability to capture key stock dynamics for Nephrops FU30.

Comparative Population variables

Scenario-specific stock dynamics derived from the SPiCT model are illustrated through estimated biomass (Figures ??–??) and fishing mortality trajectories (Figures ??–??), allowing direct comparison of model behaviour across alternative data configurations.

Biological Reference Points.

The Figure 6 compares key biological reference points across **Scenarios SC5–SC9** and **prior configurations (RUN1–RUN4)** for Nephrops FU30. Several clear patterns emerge. Estimates of **Bmsy** and **MSY** vary widely across RUNs within the same scenario. RUN1–RUN3 typically yield **much higher Bmsy and MSY** than RUN4, indicating that the more informative priors in RUN4 (on depletion, productivity, and growth) strongly constrain stock

productivity and carrying capacity. This implies that data alone are not sufficient to tightly identify absolute scale, and priors play a major role in determining long-term productivity. Across scenarios, **Fmsy** is far less variable than **Bmsy** or **MSY**, particularly under RUN4, where Fmsy consistently clusters around $\sim 0.12\text{--}0.15$. This suggests that exploitation rate at MSY is relatively well informed by the time series, even when absolute biomass is uncertain.

Scenarios **SC7** and **SC8**, which combine **normalized UWTV biomass**, **ARSA biomass**, and/or **standardized CPUE**, show tighter clustering across RUNs compared with SC5–SC6. Adding complementary fishery-independent and fishery-dependent indices improves identifiability, especially for Fmsy and Cp. SC9 displays extreme or unstable MSY and Bmsy for some RUNs, reflecting the mixed time coverage (UWTV 2015–2025, ARSA biomass 1993–2012, CPUE 2009–2024) and the survey break in 2012. This configuration increases structural uncertainty and makes results more prior-dependent.

Catch suggested is broadly similar across RUNs within scenarios, indicating that recent catches are well constrained by data and largely independent of prior assumptions. Main values in Table 6.

Management Outputs

Management strategy evaluation results by scenario are summarised in Table 8. Values correspond to equilibrium projections for relative biomass (B/B_{msy}), fishing mortality (F/F_{msy}), and expected catches under each management rule.

CHECK!!! Key reference points, including MSY, BMSY, FMSY, and current stock status indicators (B/B_{MSY} and F/F_{MSY}), are presented alongside uncertainty estimates (95% credible intervals). The results indicate that all scenarios suggest the stock is currently below BMSY, with varying degrees of fishing pressure relative to FMSY. Notably, Scenario 2 under RUN4 configuration indicates the most optimistic stock status, with B/B_{MSY} approaching 1 and F/F_{MSY} below 1, suggesting potential for recovery under current management measures. In contrast, Scenario 0 under RUN1 configuration indicates a more depleted stock status, with B/B_{MSY} significantly below 1 and F/F_{MSY} exceeding 1, highlighting the need for continued management interventions. Overall, the results underscore the sensitivity of stock status estimates to both data configuration and prior assumptions, emphasizing the importance of robust model selection in fisheries management.

Conclusions

(todo esto es preliminar, falta integrar resultados de las nuevas corridas)

WGBIE considered the results obtained for Nephrops FU 30 stock promising and will propose it as possible candidate for the next WKMSYSPiCT benchmark. However, further work should be conducted to improve the model according to the most recent SPiCT guidelines (Mildenberger, Kokkalis, & Berg, 2023). The WKMSYSPiCT benchmark, where a close contact with SPiCT expert is possible, could help to find the best model configuration to provide the assessment for this stock. Further work is needed to improve the model, including:

- Extend the time series of landings to earlier years in order to improve the stability and robustness of the model
- Add uncertainty in some years of the biomass indices time series or catches
- Fine-tune model configuration.

The results indicate that all model runs achieved MASE values below 1, demonstrating that the SPiCT model consistently outperformed the naive forecast in predicting stock dynamics. Notably, RUN3 exhibited the lowest MASE value, suggesting that the inclusion of informative priors on both initial depletion and production curve shape enhanced model predictive accuracy. These findings underscore the importance of prior specification in improving model performance and provide confidence in the SPiCT model's ability to capture key stock dynamics for Nephrops FU30.

Implication for management and benchmarking RUN4 provides the most conservative and stable estimates of productivity and fishing mortality across scenarios. SC7–SC8 appear to offer the best balance between information content and robustness, whereas SC9 should be treated cautiously due to survey discontinuity–driven uncertainty.

Code Repositories

All this work has been developed using R language. The code and data used for the SPiCT explorations for Nephrops FU30 can be found in the following GitHub repository:

All updates and improvements can be followed and obtained at the [SPiCT GitHub repository](#). Bugs and issues can be reported via the [SPiCT Releases page](#).

Nephrops Stock Assessment and Exploratory Data Analysis (ICES Division 9A) codes to replicate part of this analysis, is available at the following GitHub repository: [Nephros SPiCT codes](#)

Figures

(Que figura sería interesante incorporar aparte de las q están?)




Figure 1: Map of the Gulf of Cádiz (FU30, Division 9.a) showing the main fishing grounds for *Nephrops norvegicus*.




Figure 2: Time series of annual Nephrops landings (tons) for FU30 (Division 9.a). Bars represent reported annual catches, while the dashed horizontal line indicates the long-term mean catch over the period analysed.




Figure 3: Time series of abundance and fishery-dependent indices used in the SPiCT explorations for Nephrops FU30 (Division 9.a). Panels A–H show the different survey-based and commercial indices. Points represent observed annual values, while solid lines correspond to smoothed trends with associated uncertainty envelopes.




Figure 4: Spearman rank correlation heatmap among abundance, productivity and fishery-dependent indices for Nephrops FU30 (Division 9.a). Colours indicate the strength and direction of correlations (red: positive; blue: negative), and hierarchical clustering highlights groups of highly correlated indices.




Figure 5: Comparison of surplus production functions: Schaefer ($n=1$), Pella - Tomlinson ($n=2$) and Fox (log).





Figure 6: Comparison of stochastic biological reference points (Bmsy, Fmsy, MSY) and current catch (Cp) across SPiCT Scenarios SC5–SC9 and prior configurations (RUN5–RUN8) for Nephrops FU30. Points represent scenario–prior combinations, illustrating the sensitivity of stock productivity and exploitation reference points to alternative data configurations and prior assumptions.



spict_v1.3.8@107a32

Figure 7: Plot Spict Scenario 5 RUN 5




Figure 8: Plot Spict Scenario 9 RUN 8




Figure 9: Retrospective analysis diagnostics for Scenario SC0 under prior configuration RUN4.




Figure 10: Retrospective analysis diagnostics for Scenario SC1 under prior configuration RUN1.




Figure 11: Retrospective analysis diagnostics for Scenario SC1 under prior configuration RUN2.




Figure 12: Retrospective analysis diagnostics for Scenario SC1 under prior configuration RUN3.




Figure 13: Retrospective analysis diagnostics for Scenario SC1 under prior configuration RUN4.




Figure 14: Retrospective analysis diagnostics for Scenario SC2 under prior configuration RUN1.




Figure 15: Retrospective analysis diagnostics for Scenario SC2 under prior configuration RUN2.




Figure 16: Retrospective analysis diagnostics for Scenario SC2 under prior configuration RUN3.





Figure 17: Retrospective analysis diagnostics for Scenario SC2 under prior configuration RUN4.



spict_v1.3.8@107a32

Figure 18: Retrospective analysis diagnostics for Scenario SC3 under prior configuration RUN4.






Figure 19: Retrospective analysis diagnostics for Scenario SC4 under prior configuration RUN1.




spict_v1.3.8@107a32

Figure 20: Retrospective analysis diagnostics for Scenario SC4 under prior configuration RUN2.



spict_v1.3.8@107a32

Figure 21: Retrospective analysis diagnostics for Scenario SC4 under prior configuration RUN3.



spict_v1.3.8@107a32

Figure 22: Retrospective analysis diagnostics for Scenario SC4 under prior configuration RUN4.




Figure 23: Retrospective analysis diagnostics for Scenario SC5 under prior configuration RUN1.




Figure 24: Retrospective analysis diagnostics for Scenario SC5 under prior configuration RUN2.




Figure 25: Retrospective analysis diagnostics for Scenario SC5 under prior configuration RUN3.




Figure 26: Retrospective analysis diagnostics for Scenario SC5 under prior configuration RUN4.




Figure 27: Hindcast diagnostics for Scenario SC0 under prior configuration RUN4.




Figure 28: Hindcast diagnostics for Scenario SC1 under prior configuration RUN1.




Figure 29: Hindcast diagnostics for Scenario SC1 under prior configuration RUN2.




Figure 30: Hindcast diagnostics for Scenario SC1 under prior configuration RUN3.




Figure 31: Hindcast diagnostics for Scenario SC1 under prior configuration RUN4.




Figure 32: Hindcast diagnostics for Scenario SC2 under prior configuration RUN1.




Figure 33: Hindcast diagnostics for Scenario SC2 under prior configuration RUN2.




Figure 34: Hindcast diagnostics for Scenario SC2 under prior configuration RUN3.




Figure 35: Hindcast diagnostics for Scenario SC2 under prior configuration RUN4.




Figure 36: Estimated relative biomass trajectory for Scenario SC0 derived from the SPiCT model.




Figure 37: Estimated relative biomass trajectory for Scenario SC1 derived from the SPiCT model.




Figure 38: Estimated relative biomass trajectory for Scenario SC2 derived from the SPiCT model.




Figure 39: Estimated relative biomass trajectory for Scenario SC3 derived from the SPiCT model.

Estimated biomass trajectories – SC4




Figure 40: Estimated relative biomass trajectory for Scenario SC4 derived from the SPiCT model.

Estimated biomass trajectories – SC5




Figure 41: Estimated relative biomass trajectory for Scenario SC5 derived from the SPiCT model.




Figure 42: Estimated fishing mortality trajectory for Scenario SC0 derived from the SPiCT model.

Estimated fishing mortality trajectories – SC1




Figure 43: Estimated fishing mortality trajectory for Scenario SC1 derived from the SPiCT model.




Figure 44: Estimated fishing mortality trajectory for Scenario SC2 derived from the SPiCT model.




Figure 45: Estimated fishing mortality trajectory for Scenario SC3 derived from the SPiCT model.




Figure 46: Estimated fishing mortality trajectory for Scenario SC4 derived from the SPiCT model.




Figure 47: Estimated fishing mortality trajectory for Scenario SC5 derived from the SPiCT model.

Tables

Table 7: Mohn’s rho estimates derived from retrospective analyses of the SPiCT models for *Nephrops norvegicus* in Functional Unit 30. Values are reported for relative spawning biomass (B/B_{MSY}) and fishing mortality (F/F_{MSY}) by scenario and prior configuration (RUN).

Scenario	Run	Mohn_BBmsy	Mohn_FFmsy
SC0	RUN4	-0.22	1.87
SC1	RUN1	-0.28	1.19
SC1	RUN2	-0.08	0.19
SC1	RUN3	-0.07	0.12
SC1	RUN4	-0.07	0.15
SC2	RUN1	0.02	0.87
SC2	RUN2	-	-
SC2	RUN3	-	-
SC2	RUN4	0.12	-0.15
SC3	RUN4	0.00	0.00
SC4	RUN1	-0.31	0.12

Table 7: Mohn's rho estimates derived from retrospective analyses of the SPiCT models for *Nephrops norvegicus* in Functional Unit 30. Values are reported for relative spawning biomass (B/B_{MSY}) and fishing mortality (F/F_{MSY}) by scenario and prior configuration (RUN). (*continued*)

Scenario	Run	Mohn_BBmsy	Mohn_FFmsy
SC4	RUN2	0.00	-0.09
SC4	RUN3	0.01	-0.10
SC4	RUN4	0.07	-0.27
SC5	RUN1	45.73	-
SC5	RUN2	0.83	-
SC5	RUN3	-	0.01
SC5	RUN4	0.01	-0.07

Table 8: Projected stock status and catch outcomes under alternative management rules by scenario and prior configuration (RUN) for the SPiCT-based *Nephrops* FU30 assessment.

Management_rule	Catch	B/Bmsy	F/Fmsy
Scenario SC0 – RUN1			
1. Keep current catch	33.0	1.45	0.16
2. Keep current F	28.1	1.46	0.14
3. Fish at Fmsy	186.2	1.19	1.00
4. No fishing	0.0	1.51	0.00
5. Reduce F by 25%	21.2	1.47	0.10
6. Increase F by 25%	35.0	1.45	0.17
7. MSY hockey-stick rule	186.2	1.19	1.00
8. ICES advice rule	146.5	1.25	0.77
Scenario SC0 – RUN2			
1. Keep current catch	32.9	1.45	0.14
2. Keep current F	28.9	1.46	0.12
3. Fish at Fmsy	207.9	1.17	1.00
4. No fishing	0.0	1.50	0.00
5. Reduce F by 25%	21.8	1.47	0.09
6. Increase F by 25%	36.0	1.45	0.16
7. MSY hockey-stick rule	207.9	1.17	1.00
8. ICES advice rule	163.1	1.25	0.76
Scenario SC0 – RUN3			
1. Keep current catch	32.9	1.45	0.14
2. Keep current F	28.8	1.45	0.13
3. Fish at Fmsy	206.1	1.18	1.00
4. No fishing	0.0	1.50	0.00
5. Reduce F by 25%	21.7	1.46	0.09
6. Increase F by 25%	35.8	1.44	0.16
7. MSY hockey-stick rule	206.1	1.18	1.00
8. ICES advice rule	161.6	1.25	0.76
Scenario SC0 – RUN4			
1. Keep current catch	33.1	1.54	0.05
2. Keep current F	24.0	1.54	0.04

Table 8: Projected stock status and catch outcomes under alternative management rules by scenario and prior configuration (RUN) for the SPiCT-based Nephrops FU30 assessment.
(continued)

Management_rule	Catch	B/Bmsy	F/Fmsy
3. Fish at Fmsy	586.0	1.40	1.00
4. No fishing	0.0	1.54	0.00
5. Reduce F by 25%	18.0	1.54	0.03
6. Increase F by 25%	30.0	1.54	0.05
7. MSY hockey-stick rule	586.0	1.40	1.00
8. ICES advice rule	460.0	1.43	0.78
Scenario SC1 – RUN1			
1. Keep current catch	32.5	3.23	0.05
2. Keep current F	26.7	3.26	0.04
3. Fish at Fmsy	473.1	1.44	1.00
4. No fishing	0.0	3.37	0.00
5. Reduce F by 25%	20.1	3.29	0.03
6. Increase F by 25%	33.2	3.23	0.05
7. MSY hockey-stick rule	473.1	1.44	1.00
8. ICES advice rule	384.9	1.79	0.73
Scenario SC1 – RUN2			
1. Keep current catch	32.5	3.05	0.05
2. Keep current F	26.2	3.07	0.04
3. Fish at Fmsy	462.0	1.56	1.00
4. No fishing	0.0	3.16	0.00
5. Reduce F by 25%	19.7	3.10	0.03
6. Increase F by 25%	32.6	3.05	0.05
7. MSY hockey-stick rule	462.0	1.56	1.00
8. ICES advice rule	371.7	1.87	0.74
Scenario SC1 – RUN3			
1. Keep current catch	32.6	1.71	0.10
2. Keep current F	26.5	1.72	0.08
3. Fish at Fmsy	280.8	1.38	1.00
4. No fishing	0.0	1.76	0.00
5. Reduce F by 25%	19.9	1.73	0.06
6. Increase F by 25%	33.1	1.71	0.11
7. MSY hockey-stick rule	280.8	1.38	1.00
8. ICES advice rule	216.7	1.46	0.75
Scenario SC1 – RUN4			
1. Keep current catch	32.5	1.62	0.10
2. Keep current F	24.7	1.62	0.08
3. Fish at Fmsy	309.4	1.45	1.00
4. No fishing	0.0	1.64	0.00
5. Reduce F by 25%	18.6	1.62	0.06
6. Increase F by 25%	30.9	1.62	0.09
7. MSY hockey-stick rule	309.4	1.45	1.00
8. ICES advice rule	235.2	1.50	0.75
Scenario SC2 – RUN1			
1. Keep current catch	32.2	2.00	0.01
2. Keep current F	32.3	2.00	0.01

Table 8: Projected stock status and catch outcomes under alternative management rules by scenario and prior configuration (RUN) for the SPiCT-based Nephrops FU30 assessment.

(continued)

Management_rule	Catch	B/Bmsy	F/Fmsy
3. Fish at Fmsy	3103.3	1.82	1.00
4. No fishing	0.0	2.01	0.00
5. Reduce F by 25%	24.2	2.01	0.01
6. Increase F by 25%	40.4	2.00	0.01
7. MSY hockey-stick rule	3103.3	1.82	1.00
8. ICES advice rule	2351.1	1.86	0.75
Scenario SC2 – RUN2			
1	-	-	-
Scenario SC2 – RUN3			
1	-	-	-
Scenario SC2 – RUN4			
1. Keep current catch	32.3	0.51	0.52
2. Keep current F	32.7	0.51	0.53
3. Fish at Fmsy	61.6	0.50	1.00
4. No fishing	0.0	0.52	0.00
5. Reduce F by 25%	24.6	0.51	0.39
6. Increase F by 25%	40.7	0.51	0.66
7. MSY hockey-stick rule	60.3	0.50	0.98
8. ICES advice rule	44.2	0.51	0.71
Scenario SC3 – RUN1			
1	-	-	-
Scenario SC3 – RUN2			
1	-	-	-
Scenario SC3 – RUN3			
1	-	-	-
Scenario SC3 – RUN4			
1. Keep current catch	90.7	0.51	1.00
2. Keep current F	91.1	0.51	1.01
3. Fish at Fmsy	90.9	0.51	1.00
4. No fishing	0.2	0.51	0.00
5. Reduce F by 25%	68.3	0.51	0.75
6. Increase F by 25%	113.8	0.51	1.26
7. MSY hockey-stick rule	90.9	0.51	1.00
8. ICES advice rule	87.1	0.51	0.96
Scenario SC4 – RUN1			
1. Keep current catch	40.4	0.00	12.60
2. Keep current F	55.2	0.00	4.18
3. Fish at Fmsy	31.9	0.00	1.00
4. No fishing	0.2	0.01	0.00
5. Reduce F by 25%	54.2	0.00	3.13
6. Increase F by 25%	53.9	0.00	5.22
7. MSY hockey-stick rule	0.1	0.01	0.00
8. ICES advice rule	0.0	0.01	0.00
Scenario SC4 – RUN2			

Table 8: Projected stock status and catch outcomes under alternative management rules by scenario and prior configuration (RUN) for the SPiCT-based Nephrops FU30 assessment.
(continued)

Management_rule	Catch	B/Bmsy	F/Fmsy
1. Keep current catch	35.5	0.68	0.62
2. Keep current F	41.6	0.66	0.73
3. Fish at Fmsy	55.3	0.62	1.00
4. No fishing	0.0	0.77	0.00
5. Reduce F by 25%	31.8	0.69	0.55
6. Increase F by 25%	51.0	0.64	0.91
7. MSY hockey-stick rule	55.3	0.62	1.00
8. ICES advice rule	44.2	0.65	0.78
Scenario SC4 – RUN3			
1. Keep current catch	35.4	0.66	0.64
2. Keep current F	41.3	0.65	0.76
3. Fish at Fmsy	53.3	0.61	1.00
4. No fishing	0.0	0.75	0.00
5. Reduce F by 25%	31.5	0.67	0.57
6. Increase F by 25%	50.6	0.62	0.95
7. MSY hockey-stick rule	53.3	0.61	1.00
8. ICES advice rule	42.6	0.64	0.78
Scenario SC4 – RUN4			
1. Keep current catch	35.3	0.92	0.36
2. Keep current F	36.4	0.91	0.37
3. Fish at Fmsy	95.7	0.86	1.00
4. No fishing	0.0	0.95	0.00
5. Reduce F by 25%	27.4	0.92	0.28
6. Increase F by 25%	45.3	0.91	0.46
7. MSY hockey-stick rule	95.7	0.86	1.00
8. ICES advice rule	77.0	0.88	0.80
Scenario SC5 – RUN1			
1	-	-	-
Scenario SC5 – RUN2			
1	-	-	-
Scenario SC5 – RUN3			
1. Keep current catch	32.5	2.61	0.06
2. Keep current F	34.3	2.60	0.06
3. Fish at Fmsy	371.0	1.35	1.00
4. No fishing	0.0	2.71	0.00
5. Reduce F by 25%	25.9	2.63	0.05
6. Increase F by 25%	42.6	2.58	0.08
7. MSY hockey-stick rule	371.0	1.35	1.00
8. ICES advice rule	303.5	1.64	0.73
Scenario SC5 – RUN4			
1. Keep current catch	32.5	0.62	0.41
2. Keep current F	26.0	0.63	0.33
3. Fish at Fmsy	78.0	0.60	1.00
4. No fishing	0.0	0.64	0.00
5. Reduce F by 25%	19.6	0.63	0.24

Table 8: Projected stock status and catch outcomes under alternative management rules by scenario and prior configuration (RUN) for the SPiCT-based Nephrops FU30 assessment.

(continued)

Management_rule	Catch	B/Bmsy	F/Fmsy
6. Increase F by 25%	32.4	0.62	0.41
7. MSY hockey-stick rule	78.0	0.60	1.00
8. ICES advice rule	58.8	0.61	0.75

Table 6: Stochastic biological reference points and estimated current catch (Cp) by SPiCT scenario and prior configuration (RUN) for Nephrops FU30.

Scenario	Run	$B_{\{MSY\}}$	$F_{\{MSY\}}$	MSY	Cp
SC0	RUN1	551.406	0.276	151.027	28.099
SC0	RUN2	589.548	0.293	171.299	28.898
SC0	RUN3	603.704	0.284	169.102	28.781
SC0	RUN4	3964.471	0.103	393.814	23.973
SC1	RUN1	194.002	1.213	235.274	26.692
SC1	RUN2	242.131	0.907	219.600	26.197
SC1	RUN3	663.834	0.289	190.983	26.511
SC1	RUN4	1609.166	0.128	204.279	24.745
SC2	RUN1	15102.859	0.108	1627.234	32.302
SC2	RUN2	-	-	-	31.078
SC2	RUN3	-	-	-	31.078
SC2	RUN4	2817.064	0.044	120.904	32.674
SC3	RUN1	59940.847	-	-	0.893
SC3	RUN2	69.328	6.884	1239.830	33.734
SC3	RUN3	-	-	1911.312	32.171
SC3	RUN4	1458135.555	0.000	179.351	90.834
SC4	RUN1	33996.785	0.522	17739.924	55.213
SC4	RUN2	417.520	0.223	78.345	41.586
SC4	RUN3	424.280	0.215	74.045	41.268
SC4	RUN4	1205.914	0.092	46.054	36.394
SC5	RUN1	578.317	1.980	1144.969	32.065
SC5	RUN2	151.642	1.917	290.676	33.190
SC5	RUN3	158.706	1.406	227.578	34.272
SC5	RUN4	1769.054	0.075	126.318	26.033

Table 9: Summary of SPiCT model diagnostics and outputs by scenario and prior configuration (RUN).

Scenario	Run	Convergence	PDHess	AIC	Biomass_2025	FishingMortality_2025	ProdCurve	Mohn_BBmsy	Mohn_FFmsy
SC0	RUN1	FALSE	FALSE	81.74	11581.04	0.04	0.46	-	-
SC0	RUN2	FALSE	FALSE	75.16	11944.54	0.04	0.51	-	-
SC0	RUN3	FALSE	FALSE	79.84	12193.22	0.04	0.52	-	-
SC0	RUN4	TRUE	FALSE	85.69	105352.14	0.00	0.47	-0.22	1.87
SC1	RUN1	TRUE	FALSE	161.43	9751.92	0.04	0.23	-0.28	1.19
SC1	RUN2	TRUE	FALSE	156.39	11686.70	0.04	0.23	-0.08	0.19
SC1	RUN3	TRUE	FALSE	155.01	17693.59	0.02	0.44	-0.07	0.12
SC1	RUN4	TRUE	FALSE	154.64	43935.02	0.01	0.45	-0.07	0.15
SC2	RUN1	TRUE	FALSE	93.81	449899.34	0.00	0.48	0.02	0.87
SC2	RUN2	TRUE	FALSE	93.36	94100.83	0.01	0.50	-	-
SC2	RUN3	TRUE	FALSE	90.59	94100.95	0.01	0.50	-	-
SC2	RUN4	TRUE	FALSE	95.20	20637.30	0.02	0.59	0.12	-0.15
SC3	RUN1	FALSE	FALSE	-1101.78	9559775.29	0.00	0.09	-	-
SC3	RUN2	FALSE	FALSE	57.04	5780.24	0.08	0.68	-	-
SC3	RUN3	FALSE	FALSE	49.65	6403.39	0.07	0.74	-	-
SC3	RUN4	TRUE	FALSE	-874.48	11046283.28	0.00	0.99	0.00	0.00
SC4	RUN1	TRUE	FALSE	63.88	250.38	2.13	0.40	-0.31	0.12
SC4	RUN2	TRUE	FALSE	112.47	3286.40	0.16	0.49	0.00	-0.09
SC4	RUN3	TRUE	FALSE	109.71	3288.82	0.16	0.50	0.01	-0.10
SC4	RUN4	TRUE	FALSE	109.59	15493.14	0.03	0.45	0.07	-0.27
SC5	RUN1	TRUE	FALSE	316.98	21792.41	0.02	0.31	45.73	-
SC5	RUN2	TRUE	FALSE	322.34	5403.90	0.08	0.30	0.83	-
SC5	RUN3	TRUE	FALSE	320.42	5194.90	0.09	0.45	-	0.01
SC5	RUN4	TRUE	FALSE	317.43	17709.07	0.02	0.51	0.01	-0.07

References

- Hurtado-ferro, F., Szuwalski, C. S., Valero, J. L., Anderson, S. C., Cunningham, C. J., Johnson, K. F., ... Muradian, M. L. (2015). *Marine Science*. 72, 99–110.
- ICES. (2017). *Report of the benchmark workshop on nephrops stocks (WKNEP)* (No. CM 2016/ACOM:38). ICES.
- ICES. (2022). *Working group for the bay of biscay and the iberian waters ecoregion (WGBIE)* (No. 52; Vol. 4). ICES Scientific Reports. <https://doi.org/10.17895/ices.pub.20068988>
- ICES. (2023). *Advice on fishing opportunities*. International Council for the Exploration of the Sea. <https://doi.org/10.17895/ices.advice.22240624>
- Mildenberger, T. K., Kokkalis, A., & Berg, C. W. (2023). *Guidelines for the stochastic production model in continuous time (SPiCT)*. Retrieved from https://github.com/DTUAqua/spict/raw/master/spict/inst/doc/spict_guidelines.pdf
- Pedersen, M. W., Kokkalis, A., Mildenberger, T. K., & Berg, C. W. (2023). *Handbook for the stochastic production model in continuous time (SPiCT)*. Retrieved from https://github.com/DTUAqua/spict/blob/master/spict/inst/doc/spict_handbook.pdf
- Vila, Y., & Burgos, C. (2022). *New area proposed for the ISUNEPCA UWTV survey in the gulf of cadiz (FU 30)* (No. 52; p. 847). International Council for the Exploration of the Sea. <https://doi.org/10.17895/ices.pub.20068988>
- Vila, Y., Burgos, C., & Soriano, M. M. (2016). *Nephrops (FU 30) UWTV survey on the gulf of cadiz grounds*. International Council for the Exploration of the Sea.
- Vila, Y., & Quinzán, M. (2024). *Stochastic surplus production model in continuous time (SPiCT) explorations in the nephrops FU30 stock (gulf of cadiz)* (Technical Report No. WGBIE 2024). Lisbon, Portugal: Instituto Español de Oceanografía (IEO-CSIC), Centro Oceanográfico de Cádiz.