

p_T resolution fit vs $|\eta|^{\text{truth}}|$: r_1, r_2 with $r_0 = 0$ fixed

$$\sigma_{p_T}/p_T = \sqrt{r_0^2/p_T^2 + r_1^2 + (r_2 \times p_T)^2} \quad -- \quad \sigma_{68} = (q_{84} - q_{16})/2 \quad -- \quad Z + Z' \text{ combined}$$

$ \eta $ bin	N points	χ^2/ndf (r_0 free)	χ^2/ndf ($r_0 = 0$)	r_1 ($r_0 = 0$)	r_2 ($r_0 = 0$) [GeV^{-1}]
$0.0 \leq \eta < 0.1$	16	0.16	0.14	0.0182 ± 0.0014	0.311 ± 0.015 ($\times 10^{-3}$)
$0.1 \leq \eta < 1.05$	16	0.67	0.59	0.0209 ± 0.0011	0.192 ± 0.011 ($\times 10^{-3}$)
$1.05 \leq \eta < 1.3$	15	0.29	0.25	0.0229 ± 0.0014	0.270 ± 0.014 ($\times 10^{-3}$)
$1.3 \leq \eta < 1.7$	14	0.20	0.17	0.0290 ± 0.0016	0.283 ± 0.016 ($\times 10^{-3}$)
$1.7 \leq \eta < 2.5$	13	0.05	0.04	0.0269 ± 0.0015	0.278 ± 0.015 ($\times 10^{-3}$)
$2.5 \leq \eta < 2.8$	11	0.06	0.10	0.0331 ± 0.0021	0.425 ± 0.022 ($\times 10^{-3}$)

In red: $\chi^2/\text{ndf} > 3$, unreliable fit

In blue: r_2 , the nominal ATLAS resolution at high p_T