

Fit risoluzione in p_T vs $|\eta^{\text{truth}}|$: r_0 libero vs $r_0 = 0$

$$\sigma_{p_T}/p_T = \sqrt{r_0^2/p_T^2 + r_1^2 + (r_2 \times p_T)^2} \quad -- \quad \sigma_{68} = (q_{84} - q_{16})/2 \quad -- \quad Z + Z. \text{ combinati}$$

$ \eta $ bin	n punti	r_0 libero [GeV]	χ^2/ndf (r_0 libero)	χ^2/ndf ($r_0 = 0$)	r_2 [GeV ⁻¹]
$0.0 \leq \eta < 0.1$	16	0.000 ± 36.345	0.58	0.51	$0.224 \pm 0.012 \ (\times 10^{-3})$
$0.1 \leq \eta < 1.05$	16	0.000 ± 8.983	1.54	1.35	$0.126 \pm 0.008 \ (\times 10^{-3})$
$1.05 \leq \eta < 1.3$	15	0.000 ± 37.178	0.51	0.45	$0.124 \pm 0.008 \ (\times 10^{-3})$
$1.3 \leq \eta < 1.7$	14	0.000 ± 32.319	1.68	1.47	$0.161 \pm 0.011 \ (\times 10^{-3})$
$1.7 \leq \eta < 2.5$	13	0.000 ± 32.491	0.64	0.56	$0.089 \pm 0.008 \ (\times 10^{-3})$
$2.5 \leq \eta < 2.8$	11	0.223 ± 0.334	0.08	0.09	$0.104 \pm 0.010 \ (\times 10^{-3})$

In rosso: $\chi^2/\text{ndf} > 3$, fit poco affidabile

In blu: r_2 , la risoluzione nominale ATLAS ad alto p_T