

Fit risoluzione in p_T vs $|\eta|^{\text{truth}}$: r_0 libero vs $r_0 = 0$

$$\sigma_{p_T}/p_T = \sqrt{r_0^2/p_T^2 + r_1^2 + (r_2 \times p_T)^2} \text{ -- tutti i sample combinati (Z + Z')}$$

$ \eta $ bin	n pt	r_0 free	χ^2/ndf free	χ^2/ndf fix $r_0=0$	r_2 [GeV^{-1}]
$0.0 \leq \eta < 0.1$	14	0.000 ± 4.164	13.64	12.51	$0.351 \pm 0.019 \ (\times 10^{-3})$
$0.1 \leq \eta < 1.05$	14	0.000 ± 0.215	13.21	12.11	$0.247 \pm 0.012 \ (\times 10^{-3})$
$1.05 \leq \eta < 1.3$	14	0.000 ± 3.294	13.99	12.82	$0.270 \pm 0.014 \ (\times 10^{-3})$
$1.3 \leq \eta < 1.7$	14	0.000 ± 3.462	13.16	12.06	$0.288 \pm 0.015 \ (\times 10^{-3})$
$1.7 \leq \eta < 2.5$	13	0.000 ± 5.266	8.09	7.35	$0.289 \pm 0.018 \ (\times 10^{-3})$
$2.5 \leq \eta < 2.8$	11	0.304 ± 0.092	5.23	5.99	$0.262 \pm 0.023 \ (\times 10^{-3})$

In rosso: $\chi^2/\text{ndf} > 3$, fit poco affidabile

In blu: r_2 , la risoluzione nominale ATLAS ad alto p_T