

Fit risoluzione in p_T vs $|\eta^{\text{truth}}|$: r_0 libero vs $r_0 = 0$

$$\sigma_{p_T}/p_T = \sqrt{r_0^2/p_T^2 + r_1^2 + (r_2 \times p_T)^2} \quad -- \quad \sigma_{68} = (q_{84} - q_{16})/2 \quad -- \quad Z + Z. \text{ combinati}$$

$ \eta $ bin	n punti	r_0 libero [GeV]	χ^2/ndf (r_0 libero)	χ^2/ndf ($r_0 = 0$)	r_2 [GeV $^{-1}$]
$0.0 \leq \eta < 0.1$	16	0.000 ± 45.213	0.16	0.14	0.311 ± 0.015 ($\times 10^{-3}$)
$0.1 \leq \eta < 1.05$	16	0.000 ± 30.900	0.67	0.59	0.192 ± 0.011 ($\times 10^{-3}$)
$1.05 \leq \eta < 1.3$	15	0.001 ± 38.310	0.29	0.25	0.270 ± 0.014 ($\times 10^{-3}$)
$1.3 \leq \eta < 1.7$	14	0.000 ± 45.304	0.20	0.17	0.283 ± 0.016 ($\times 10^{-3}$)
$1.7 \leq \eta < 2.5$	13	0.000 ± 49.423	0.05	0.04	0.278 ± 0.015 ($\times 10^{-3}$)
$2.5 \leq \eta < 2.8$	11	0.404 ± 0.347	0.06	0.10	0.430 ± 0.024 ($\times 10^{-3}$)

In rosso: $\chi^2/\text{ndf} > 3$, fit poco affidabile

In blu: r_2 , la risoluzione nominale ATLAS ad alto p_T