

Fit risoluzione in p_T vs $|\eta^{\text{truth}}|$: r_0 libero vs $r_0 = 0$

$$\sigma_{p_T}/p_T = \sqrt{r_0^2/p_T^2 + r_1^2 + (r_2 \times p_T)^2} \text{ -- tutti i sample combinati (Z + Z')}$$

$ \eta $ bin	n pt	r_0 free	χ^2/ndf free	χ^2/ndf fix $r_0=0$	r_2 [GeV ⁻¹]
$0.0 \leq \eta < 0.1$	14	0.000 ± 0.029	1331.76	1220.78	$0.184 \pm 0.001 (\times 10^{-3})$
$0.1 \leq \eta < 1.05$	14	0.000 ± 0.002	18370.18	16839.33	$0.090 \pm 0.000 (\times 10^{-3})$
$1.05 \leq \eta < 1.3$	14	0.000 ± 0.037	1276.25	1169.90	$0.075 \pm 0.000 (\times 10^{-3})$
$1.3 \leq \eta < 1.7$	14	0.000 ± 0.015	4905.88	4497.05	$0.103 \pm 0.000 (\times 10^{-3})$
$1.7 \leq \eta < 2.5$	13	0.000 ± 0.022	1910.93	1737.21	$0.080 \pm 0.000 (\times 10^{-3})$
$2.5 \leq \eta < 2.8$	11	0.248 ± 0.003	26.01	166.41	$0.100 \pm 0.001 (\times 10^{-3})$

In rosso: $\chi^2/\text{ndf} > 3$, fit poco affidabile

In blu: r_2 , la risoluzione nominale ATLAS ad alto p_T