

# Properties of Elementary Particles

Max Bock

Marvin Hoffmann

Email s6mabock@uni-bonn.de

Email marvin.hoffmann@uni-bonn.de

April 19, 2017

Nature has always looked like a horrible mess, but as we go along we see patterns and put theories together; a certain clarity comes and things get simpler. - Richard P. Feynman [1]

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Theoretical Background</b>	<b>2</b>
2.1	Basics of particle physics . . . . .	2
2.1.1	Basics of Standard Model of Particle Physics . . . . .	2
2.1.2	Relativistic Kinematics . . . . .	3
2.2	Bubble Chamber . . . . .	3
2.3	Proton Proton interaction . . . . .	4
2.3.1	Elastic and Inelastic Scattering . . . . .	4
2.3.2	Cross-Section . . . . .	4
2.3.3	Radiation Length . . . . .	4
2.4	$\omega$ Meson . . . . .	5
2.4.1	Breit-Wigner Distribution . . . . .	5
2.4.2	Gauss Distribution . . . . .	5
2.4.3	Dalitz Plots . . . . .	5
<b>3</b>	<b>Experimental Set-Up and Measurements</b>	<b>5</b>
<b>4</b>	<b>Analysis</b>	<b>5</b>
<b>5</b>	<b>Conclusion</b>	<b>5</b>
<b>6</b>	<b>Appendix</b>	<b>5</b>

# 1 Introduction

In this experiment typical interactions in elementary particle physics will be studied. Therefore, first proton-proton interactions observed in a bubble chamber are analysed. Further, the films from the bubble chamber are used to study  $\pi$ - $\mu$ - $e$  decays and observe strange neutral particles which have to be identified. In the end certain characteristics of  $\omega$  mesons are determined in a computer-based analysis of many  $pp \rightarrow pp\pi^+\pi^-\pi^0$  events.

## 2 Theoretical Background

### 2.1 Basics of particle physics

#### 2.1.1 Basics of Standard Model of Particle Physics

The Standard Model of particle physics implies the unified theory of the electromagnetic, weak and strong interaction and divides the elementary particles into two classes, namely bosons and fermions.

The bosons act as exchange particles of the fundamental forces and have an integer spin. These ones are often called gauge bosons. There are eight massless gluons, which carry the colour charge of the strong force. Due to the fact that the gluons themselves have a colour charge, they interact with each other. The interaction range of the gluon is the smallest of all gauge bosons and is in the order of 1 fm. The massless photon is the exchange particle of the electromagnetic force, which has no electromagnetic charge. The range of this gauge boson is infinity. For the weak interaction there are two exchange particles, the  $Z$  boson which carries no electromagnetic charge, whereby the  $W^\pm$  bosons have a charge of  $\pm 1$ . The last boson is the so called Higgs particle- This is a scalar boson which delivers the mass of all particles which are coupled to the Higgs field.

The fermions are divided into two subclasses, the quarks and the leptons which are also subdivided into three generations. Each fermion has a corresponding anti particle with the same mass but opposite charge, including electric, weak and strong charge.

The quarks are the only particles which interact strongly and carry colour charge but also interact electromagnetically and weakly. Due to the fact that the quarks can not appear alone apart they must form colour neutral formations. These formations are called hadrons which are divided into two classes, the mesons and the baryons. The mesons consist of a quark-antiquark pair and the baryons consist of three quarks. Each quark has either a charge of  $2/3$  or  $-1/3$  and further a spin of  $1/2$ .

The leptons consist of electron, muon and tau with the corresponding neutrino. Whereas the neutrinos have no charge and only interact weakly, the electron, muon and tau have a charge of  $-1$  and interact weakly and electromagnetically. Additionally, all leptons have a spin of  $1/2$ . Due to the fact that the muon and the tau have a higher mass than the electron, free muons and taus will decay most likely into an electron and further neutrinos in the last stable decay step, whereas the electron is stable. [2]

# Standard Model of Elementary Particles




Figure 1: Standard Model of Elementary Particles.

## 2.1.2 Relativistic Kinematics

The ratio between moving mass  $m$  and energy  $E$  in relativistic kinematics is given by Einsteins famous equation [3]

$$E = mc^2 \quad (1)$$

where  $c$  is the speed of light. This equation can also be written as [3]

$$E^2 c^4 = m_0^2 c^4 + \vec{p}^2 c^2 \quad (2)$$

with the momentum vector  $\vec{p}$  and the rest mass  $m_0$ , which is directly related to the mass  $m$  via [3]

$$m = \gamma m_0 \quad (3)$$

with the Lorentz factor  $\gamma$  [3]

$$\gamma = \frac{1}{\sqrt{1 - \beta^2}} \quad \text{with} \quad \beta = \frac{v}{c} \quad (4)$$

If one defines the four-momentum  $p$  as vector [3]

$$p = \begin{pmatrix} E \\ \vec{p} \end{pmatrix} \quad (5)$$

the invariant mass  $M_n$  of a  $n$ -particle system will be given by [3]

$$M_n^2 = \left( \sum p_i \right)^2 = \left( \sum E_i \right)^2 - \left( \sum \vec{p}_i \right)^2 \quad (6)$$

## 2.2 Bubble Chamber

Till the early 80's, the bubble chamber was one of the most popular devices for detecting high energy elementary particles at accelerators. Therefore, in a chamber that can be seen in figure 2 a liquid which is nearly below the critical boiling temperature gets sequentially decompressed and compressed by a piston. While the liquid is decompressed it reaches an overheated state in which ionizing particles cause bubbles along their tracks. A particle will be deflected by a magnetic field if the particle has a electric charge. These bubbles can be detected by the use of cameras at different angles.




Figure 2: Illustration of a bubble chamber. [4] - edited

## 2.3 Proton Proton interaction

### 2.3.1 Elastic and Inelastic Scattering

In the elastic scattering the particles before and after the scattering process are identical. Further, the sum of kinetic energy and the momentum stay unchanged during the event. In the inelastic case the particles after the action must not be the same as before. In addition, the momentum is conserved but the kinetic energy is transferred into for instance binding or ionizing energy. [2]

### 2.3.2 Cross-Section

The cross-section is one of the most important quantities for description and interpretation in scattering experiments. This is related to the quantum mechanically probability of a reaction between the incoming particle and the target. In a proton-proton interaction the cross-section  $\sigma$  is given by [5]

$$\sigma = \frac{1}{nl} \ln \left( \frac{N_p}{N_p - N_i} \right) \quad (7)$$

with the concentration of scattering centres  $n$ , the observed length of the bubble chamber  $l$ , the number of incoming protons  $N_p$  and the number of prime interactions between two protons  $N_i$ . [6]

### 2.3.3 Radiation Length

The radiation length is defined as the average range where the energy of an electron is reduced to  $1/e$  by bremsstrahlung. Further, the mean free path of the  $e^+e^-$  pair production process is approximately  $7/9$  of the radiation length. The radiation length  $X_0$  is given by [7]

$$X_0 \approx \frac{1}{4\alpha n_n Z^2 r_e^2 \ln(287/Z^{1/2})} \quad (8)$$

where  $n_n$  is the number density of nuclei,  $r_e$  is the classical radius of an electron,  $Z$  is the atomic number and the fine structure constant  $\alpha$ . [7]

## 2.4 $\omega$ Meson

The  $\omega$  meson is a vector meson ( $J^P = 1^-$ ) to which an isospin of  $I = 0$  can be assigned. The corresponding wave function can be written as [2]

$$|\omega\rangle = \frac{1}{\sqrt{2}} \{ |u^\uparrow \bar{u}^\uparrow\rangle + |d^\uparrow \bar{d}^\uparrow\rangle \} \quad (9)$$

### 2.4.1 Breit-Wigner Distribution

A resonance is a system of shortly lived bounded states which are formed by two partners in a collision. In this case, the mass dependence of the cross-section in the vicinity of the resonance mass of  $m_\omega$  is described by the Breit-Wigner formula [5]

$$\sigma_{BW}(m) = \frac{\Gamma_\omega}{2\pi} \frac{1}{(m - m_\omega)^2 + \Gamma_\omega^2/4} \quad (10)$$

### 2.4.2 Gauss Distribution

The Gauss distribution is a good approximation to describe many phenomena in physics. The continuous distribution is given by [8]

$$\sigma_G(m) = \frac{1}{s\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{m-m_\omega}{s}\right)^2} \quad (11)$$

whereupon  $s$  is the standard deviation.

### 2.4.3 Dalitz Plots

To classify decays of three particles with the same mass Dalitz plots are often used to determine the spin and parity of resonances. Therefore the kinetic energies of the three particles  $T_1$ ,  $T_2$  and  $T_3$  are presented in a two dimensional diagram as shown in figure.

## 3 Experimental Set-Up and Measurements

## 4 Analysis

## 5 Conclusion

## 6 Appendix

## References

- [1] R. P. Feynman: *QED- The Strange Theory of Light and Matter* (Princeton University Press, 1985)
- [2] B. Povh, K. Rith, C. Scholz, F. Zetsche, W. Rodejohann: *Teilchen und Kerne - Eine Einführung in die physikalischen Konzepte* (Springer Spektrum, 2014) 9th. edition
- [3] E. Reibhan: *Theoretische Physik: Relativitätstheorie und Kosmologie* (Spektrum, 2012) 1st. edition
- [4] [https://en.wikipedia.org/wiki/Bubble\\_chamber#/media/File:Bubble-chamber.svg](https://en.wikipedia.org/wiki/Bubble_chamber#/media/File:Bubble-chamber.svg)  
(last downloaded at 7th. April 2017)
- [5] G. Seul: *Praktikumsversuche zur Einführung in die Hochenergiephysik* (Bonn, 1977)
- [6] B. R. Martin: *Nuclear and Particle Physics - An Introduction* (Wiley, 2009) 2nd. edition
- [7] M. Thomson: *Modern Particle Physics* (Cambridge University Press, 2015) 3rd. edition
- [8] L. Fahrmeir, A. Hamerle, G. Tutz: *Multivariate statistische Verfahren* (Walter de Gruyter, 1996) 2nd. edition