

PROBABILITY III

Lakshmi Priya M E, notes by Ramdas Singh

Fifth Semester

Contents

1	REFINING THE NOTION OF PROBABILITY	1
1.1	σ -algebras	2
1.1.1	Other Systems of Sets	4
1.2	The Lebesgue Measure	5
1.2.1	Carathéodory's Extension Theorem	6
1.3	Measurable Maps and Random Variables	7
	Index	9

Chapter 1

REFINING THE NOTION OF PROBABILITY

July 21st.

Consider the random walk on the line; a drunk ant sits at the origin and takes steps of length 1 to the left or right with equal probability. We wish to obtain the probability that after an even n steps, the ant returns to the origin. Of course, the natural approach is to frame this problem in a discrete probability space.

Definition 1.1. A *discrete probability space* is a pair (Ω, q) where Ω is a non-empty finite or countably infinite set called the sample space, and $q : \Omega \rightarrow [0, 1]$ is a function satisfying $\sum_{\omega \in \Omega} q(\omega) = 1$. Any subset of Ω is termed an event, and the probability of an event $A \subseteq \Omega$ is defined as $\mathbb{P}(A) = \sum_{\omega \in A} q(\omega)$.

For the problem above, we can take the sample space to be

$$\Omega = \{\mathbf{x} = (x_0 = 0, x_1, x_2, \dots, x_n) \in \mathbb{Z}^{n+1} \mid x_{i+1} - x_i \in \{-1, 1\} \text{ for all } 0 \leq i < n\}, \quad (1.1)$$

and the mapping q to be $q(\mathbf{x}) = 2^{-n}$ for all $\mathbf{x} \in \Omega$. Then, the event that the ant returns to the origin after n steps is given by $A = \{\mathbf{x} \in \Omega \mid x_n = 0\}$, and we have

$$\mathbb{P}(A) = \sum_{\mathbf{x} \in A} q(\mathbf{x}) = 2^{-n} |A| = 2^{-n} \binom{n}{n/2}. \quad (1.2)$$

Reframing the problem, let us consider a random walk on the line starting at the origin, and continuing indefinitely. In this case, the sample space is given by

$$\Omega = \{\mathbf{x} = (x_0 = 0, x_1, x_2, \dots) \in \mathbb{Z}^{\mathbb{N}} \mid x_{i+1} - x_i \in \{-1, 1\} \text{ for all } i \in \mathbb{N}\}. \quad (1.3)$$

This Ω is uncountably infinite. In such a case, we cannot define a probability measure by assigning a positive probability to each element of Ω . As another example of difficulty in dealing with uncountable sample spaces, consider the problem of choosing a real number uniformly at random from the interval $[0, 1]$. If we attempt to assign a positive probability to each element of $[0, 1]$, we would have to assign a probability of 0 to each element, since there are uncountably many elements in $[0, 1]$. However, this would lead to a total probability of 0, which is not valid. Of course, we know that the probability of choosing a specific element or a countable set of elements from $[0, 1]$ is 0, but there then arises the question of how to assign a probability to an uncountable set of elements, such as the Cantor set. We can work in a slightly different viewpoint.

Definition 1.2. A *probability space* is a tuple $(\Omega, \mathcal{F}, \mathbb{P})$ where Ω is a non-empty set called the sample space, $\mathcal{F}$ is the collection of all subsets of Ω ($\sim 2^{\Omega}$), and $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$ is a function satisfying $\mathbb{P}(\Omega) = 1$ and countable additivity, that is, for any countable collection of pairwise disjoint sets $A_1, A_2, \dots \in \mathcal{F}$,

we have

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i). \quad (1.4)$$

Using this definition, we look to define a $\mathbb{P}$ on subsets of $[0, 1]$ that satisfies the above properties, along with the property that $\mathbb{P}([a, b]) = b - a$. Thus, we have $\mathbb{P}([0, 1]) = 1$. If we have two disjoint intervals $I_1, I_2 \subseteq [0, 1]$, then $\mathbb{P}(I_1 \cup I_2) = \mathbb{P}(I_1) + \mathbb{P}(I_2)$. This is a natural property to expect from a probability measure. What about a general set $S \subseteq [0, 1]$? One can observe that if we have a sequence of intervals $I_1, I_2, \dots$ such that $S \subseteq \bigcup_{k=1}^{\infty} I_k$, then we would have $\mathbb{P}(S) \leq \sum_{k=1}^{\infty} \mathbb{P}(I_k)$. This is because the probability of S should be less than or equal to the probability of any set that contains S . Thus, we can define

$$\mathbb{P}_*(S) = \inf \left\{ \sum_{k=1}^{\infty} |I_k| \mid S \subseteq \bigcup_{k=1}^{\infty} I_k, I_k \subseteq [0, 1] \right\}. \quad (1.5)$$

We have not verified that this is a valid definition of a probability measure, hence the subscript $*$. Via this, let us try computing $\mathbb{P}_*(\mathbb{Q} \cap [0, 1])$. Enumerate the rationals in $[0, 1]$ as $\{q_1, q_2, \dots\}$. For any $\epsilon > 0$, we can choose intervals $I_k = [q_k - \epsilon/2^{k+1}, q_k + \epsilon/2^{k+1}] \cap [0, 1]$ for each $k \in \mathbb{N}$. Note that $\bigcup_{k=1}^{\infty} I_k$ contains all the rationals in $[0, 1]$, and $\sum_{k=1}^{\infty} |I_k| \leq \sum_{k=1}^{\infty} \epsilon/2^k = \epsilon$. Since ϵ was arbitrary, we have $\mathbb{P}_*(\mathbb{Q} \cap [0, 1]) = 0$. This is consistent with our intuition that the probability of choosing a rational number from $[0, 1]$ is 0. So, $\mathbb{P}_*$ seems to be a reasonable definition of a probability measure on $([0, 1], 2^{[0, 1]})$. However, we will show later that there exists a set $\mathcal{A} \subseteq [0, 1]$ such that $\mathbb{P}_*(\mathcal{A}) = \mathbb{P}_*([0, 1] \setminus \mathcal{A}) = 1$, contradicting the property that $\mathbb{P}_*([0, 1]) = 1$. We state the following theorem without proof.

Theorem 1.3. *There is no probability measure $\mathbb{P}$ on $([0, 1], 2^{[0, 1]})$ such that $\mathbb{P}([a, b]) = b - a$ for all $[a, b] \subseteq [0, 1]$.*

If we remove the requirement that the probability measure of a subinterval $[a, b]$ is equal to its length, then we can define a probability measure on $([0, 1], 2^{[0, 1]})$. For example, we can define $\mathbb{P}(S) = 1$ if $1/2 \in S$ and $\mathbb{P}(S) = 0$ otherwise. This is a valid probability measure, but it is not very useful. Keeping the requirement that the probability measure of a subinterval $[a, b]$ is equal to its length, we can instead restrict the collection of subsets of $[0, 1]$ on which we define the probability measure.

Definition 1.4. A *probability space* is a tuple $(\Omega, \mathcal{F}, \mathbb{P})$ where

- Ω is a non-empty set called the sample space,
- $\mathcal{F}$ is a collection of subsets of Ω satisfying the following properties:
 - $\Omega \in \mathcal{F}$,
 - if $A \in \mathcal{F}$, then $\Omega \setminus A \in \mathcal{F}$,
 - if $A_1, A_2, \dots \in \mathcal{F}$, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$,
- $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$ is a function satisfying:
 - $\mathbb{P}(\Omega) = 1$,
 - if $A_1, A_2, \dots \in \mathcal{F}$ are pairwise disjoint, then $\mathbb{P}(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$.

Such an $\mathcal{F}$ is termed a σ -algebra on Ω , and such a $\mathbb{P}$ is termed a *probability measure* on $(\Omega, \mathcal{F})$.

1.1 σ -algebras

Example 1.5. Given Ω , trivial possible choices for $\mathcal{F}$ are $\{\emptyset, \Omega\}$ and 2^Ω . A more interesting example is the following: suppose Ω is uncountable. Then $\mathcal{F} = \{A \subseteq \Omega \mid A \text{ is countable or } \Omega \setminus A \text{ is countable}\}$ is a σ -algebra on Ω .

Let $S \subseteq 2^\Omega$ be a collection of subsets of Ω . We claim that there exists a notion of the smallest σ -algebra containing S . This is called the σ -algebra generated by S , and is denoted by $\sigma(S)$.

Lemma 1.6. *For any collection $S \subseteq 2^\Omega$, there exists a unique smallest σ -algebra containing S satisfying*

$$\sigma(S) = \bigcap \{ \mathcal{F} \subseteq 2^\Omega \mid \mathcal{F} \text{ is a } \sigma\text{-algebra and } S \subseteq \mathcal{F} \}. \quad (1.6)$$

Proof. It is easy to check that $\sigma(S)$ is indeed a σ -algebra containing S . If $\mathcal{F}$ is any σ -algebra containing S , then $\sigma(S) \subseteq \mathcal{F}$. Thus, $\sigma(S)$ is the unique smallest σ -algebra containing S . ■

Definition 1.7. Let (X, d) be a metric space (or a topological space), and let $\mathcal{S} \subseteq 2^X$ be the collection of all open sets in X . Then, the σ -algebra $\sigma(\mathcal{S})$ is called the *Borel σ -algebra* on X , and its elements are called *Borel sets*.

We will show that there exists a probability measure on $([0, 1], \mathcal{B})$, where $\mathcal{B}$ is the Borel σ -algebra on $[0, 1]$, which extends the notion of length of intervals to Borel sets.

Let us return to the problem of indefinite random walks on the line. Let $\Omega = \{ \omega = (\omega_1, \omega_2, \dots) \mid \omega_i \in \{0, 1\} \}$, where $\omega_i = 0$ corresponds to a step to the left and $\omega_i = 1$ corresponds to a step to the right. If we let $A = \{0, 1\}$, then $\Omega = A^\mathbb{N}$; consider the discrete topology on A , and the product topology on Ω . If we let

$$\mathcal{S} = \{ U_1 \times U_2 \times \dots \times U_k \times A \times A \times \dots \mid U_i = \{0\} \text{ or } \{1\} \} \quad (1.7)$$

then one can show that the σ -algebra generated by $\mathcal{S}$ is the Borel σ -algebra on Ω . We can set $\mathcal{F} = \sigma(\mathcal{S})$, and define a probability measure $\mathbb{P}$ on $(\Omega, \mathcal{F})$ by setting $\mathbb{P}(U_1 \times U_2 \times \dots \times U_k \times A \times A \times \dots) = 2^{-k}$ for all $U_i = \{0\}$ or $\{1\}$.

July 24th.

Example 1.8. Let (X, d) be a separable metric space (that is, there exists a countable dense subset). The Borel σ -algebra on X is then

$$\mathcal{B}(X) = \sigma(\{ B(p, r) \}_{p \in X, r > 0}). \quad (1.8)$$

Note that the usual containment is $\sigma(\{ B(p, r) \}_{p \in X, r > 0}) \subseteq \mathcal{B}(X)$ for a metric space X . So, it suffices to show that if $V \in \tau$, the topology on X , then $V \in \sigma(\{ B(p, r) \}_{p \in X, r > 0})$. Let Q be a countable dense subset of X . Then, for any $V \in \tau$, we have

$$V = \bigcup_{n=1}^{\infty} B(q_n, r_n), \quad (1.9)$$

where $q_n \in Q \cap V$ and $r_n > 0$ (where r_n is the largest possible radius such that $B(q_n, r_n) \subseteq V$). Since $\sigma(\{ B(p, r) \}_{p \in X, r > 0})$ is a σ -algebra, we have $V \in \sigma(\{ B(p, r) \}_{p \in X, r > 0})$. Thus, we have $\mathcal{B}(X) = \sigma(\{ B(p, r) \}_{p \in X, r > 0})$.

Example 1.9. Let $(\Sigma, \mathcal{F}, \mathbb{P})$ be a probability space.

- $\mathcal{F}$ is closed under countable unions, countable intersections, differences, and symmetric differences. Countable unions and intersections follow from the definition of a σ -algebra. If $A, B \in \mathcal{F}$, then $A \setminus B = A \cap (\Omega \setminus B) \in \mathcal{F}$. Finally, $A \triangle B = (A \setminus B) \cup (B \setminus A) \in \mathcal{F}$.
- Let $(\Sigma, \mathcal{F}, \mathbb{P})$ be a probability space, and let $A_1, A_2, \dots \in \mathcal{F}$. Define

$$\limsup_{n \rightarrow \infty} A_n := \{ x \in \Omega \mid x \text{ belongs to infinitely many } A_n \} \quad (1.10)$$

and

$$\liminf_{n \rightarrow \infty} A_n := \{ x \in \Omega \mid x \text{ belongs to all but finitely many } A_n \}. \quad (1.11)$$

Then, both the $\limsup$ and $\liminf$ belong in $\mathcal{F}$. To see this, simply note that

$$\limsup_{n \rightarrow \infty} A_n = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k, \quad \liminf_{n \rightarrow \infty} A_n = \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k. \quad (1.12)$$

- One may show that $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(A) \leq \mathbb{P}(B)$ if $A \subseteq B$, $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$, and $\mathbb{P}(\bigcup A_n) \leq \sum \mathbb{P}(A_n)$ for any countable collection of sets $A_1, A_2, \dots \in \mathcal{F}$.
- For a nested sequence of sets $A_1 \subseteq A_2 \subseteq \dots$ in $\mathcal{F}$, let $A = \bigcup_{n=1}^{\infty} A_n$ (this is usually denoted by $A_n \uparrow A$). Then, $\mathbb{P}(A) = \lim_{n \rightarrow \infty} \mathbb{P}(A_n)$. To this end, let $B_n = A_n \setminus A_{n-1}$ for $n > 1$, and $B_1 = A_1$. Then $A = \bigcup_{n=1}^{\infty} B_n$. Thus,

$$\mathbb{P}(A) = \sum_{n=1}^{\infty} \mathbb{P}(B_n) = \lim_{N \rightarrow \infty} \sum_{n=1}^N \mathbb{P}(B_n) = \lim_{N \rightarrow \infty} \sum_{n=2}^N \mathbb{P}(A_n) - \mathbb{P}(A_{n-1}) + \mathbb{P}(A_1) = \lim_{N \rightarrow \infty} \mathbb{P}(A_N). \quad (1.13)$$

1.1.1 Other Systems of Sets

The following are other interesting collection of subsets of Ω besides the Borel σ -algebra.

1. π -system: a collection of subsets $S \subseteq \Omega$ such that $A, B \in S$ implies $A \cap B \in S$.
2. λ -system: a collection Λ of subsets of Ω such that $\Omega \in \Lambda$, if $A, B \in \Lambda$ with $A \subseteq B$, then $B \setminus A \in \Lambda$, and if $A_1, A_2, \dots \in \Lambda$, then $\bigcup_{n=1}^{\infty} A_n \in \Lambda$.
3. algebra: a collection $\mathcal{A}$ of subsets of Ω such that $\Omega \in \mathcal{A}$, if $A \in \mathcal{A}$, then $\Omega \setminus A \in \mathcal{A}$, and if $A, B \in \mathcal{A}$, then $A \cup B \in \mathcal{A}$.

We had defined $\omega(S)$, the σ -algebra generated by a collection of subsets $S \subseteq 2^\Omega$. We can similarly define $\pi(S)$, the π -system generated by S , $\lambda(S)$, the λ -system generated by S , and $\mathcal{A}(S)$, the algebra generated by S .

Theorem 1.10 (*Sierpiński-Dynkin π - λ Theorem*). *Let $S, \mathcal{F} \subseteq 2^\Omega$ be collections of subsets of Ω . Then,*

1. $\mathcal{F}$ is a σ -algebra if and only if $\mathcal{F}$ is both a π -system and a λ -system.
2. If S is a π -system, then $\lambda(S) = \sigma(S)$.

Proof. 1. If $\mathcal{F}$ is a σ -algebra, then it is trivially both a π -system and a λ -system. Conversely, if $\mathcal{F}$ is both a π -system and a λ -system, we need to verify the axioms of a σ -algebra. We have $\Omega \in \mathcal{F}$ since $\mathcal{F}$ is a λ -system. If $A \in \mathcal{F}$, then $\Omega \setminus A \in \mathcal{F}$ since $\mathcal{F}$ is a λ -system. Finally, if $A_1, A_2, \dots \in \mathcal{F}$, then we can write $B_n = \bigcup_{i=1}^n A_i \in \mathcal{F}$ since $A_i^c \in \mathcal{F}$ and $B_n^c = \bigcap_{i=1}^n A_i^c \in \mathcal{F}$ since $\mathcal{F}$ is a π -system. Then, $B_n \uparrow B = \bigcup_{i=1}^{\infty} A_i$, and since $\mathcal{F}$ is a λ -system, we have $B \in \mathcal{F}$. Thus, $\mathcal{F}$ is a σ -algebra.

2. Let S be a π -system. Since $\lambda(S) \subseteq \sigma(S)$, it suffices to show that $\lambda(S)$ is a π -system. Fix $A \in S$, and define $\mathcal{C}_A = \{B \in \lambda(S) \mid A \cap B \in \lambda(S)\}$. Note that $S \subseteq \mathcal{C}_A$ because S is a π -system. We show that $\mathcal{C}_A$ is a λ -system. We have $\Omega \in \mathcal{C}_A$ since $A \cap \Omega = A \in \lambda(S)$. If $B_1, B_2 \in \mathcal{C}_A$ with $B_1 \subseteq B_2$, then $A \cap B_1 \subseteq A \cap B_2$, and since $A \cap B_2 \in \lambda(S)$, we have $(A \cap B_2) \setminus (A \cap B_1) = A \cap (B_2 \setminus B_1) \in \lambda(S)$, and thus $B_2 \setminus B_1 \in \mathcal{C}_A$. Finally, if $B_1, B_2, \dots$ are in $\mathcal{C}_A$, then $A \cap B_n \in \lambda(S)$ for all n , and thus $\bigcup (A \cap B_n) = A \cap (\bigcup B_n) \in \lambda(S)$, and hence $\bigcup B_n \in \mathcal{C}_A$. Thus, $\mathcal{C}_A$ is a λ -system. Since $\lambda(S)$ is the smallest λ -system containing S , we have $\lambda(S) = \mathcal{C}_A$, and hence for any $B, C \in S$, we have $B, C \in S$ implies $B \cap C = A \cap C \in S$. Thus, $\lambda(S)$ is a π -system. ■

Lemma 1.11. *Let S be a π -system, and let $\mathbb{P}$ and $\mathbb{Q}$ be two probability measures on $(\Sigma, \sigma(S))$ such that $\mathbb{P}(A) = \mathbb{Q}(A)$ for all $A \in S$. Then, $\mathbb{P}(A) = \mathbb{Q}(A)$ for all $A \in \sigma(S)$.*

Proof. Let $\mathcal{C} = \{B \in \sigma(S) \mid \mathbb{P}(B) = \mathbb{Q}(B)\}$. Then, $S \subseteq \mathcal{C}$, and we show that $\mathcal{C}$ is a λ -system. We have $\Omega \in \mathcal{C}$ since $\mathbb{P}(\Omega) = 1 = \mathbb{Q}(\Omega)$. If $B_1, B_2 \in \mathcal{C}$ with $B_1 \subseteq B_2$, then $\mathbb{P}(B_2 \setminus B_1) = \mathbb{P}(B_2) - \mathbb{P}(B_1) = \mathbb{Q}(B_2) - \mathbb{Q}(B_1) = \mathbb{Q}(B_2 \setminus B_1)$, and hence $B_2 \setminus B_1 \in \mathcal{C}$. Finally, if $B_1, B_2, \dots$ are in $\mathcal{C}$, then $\mathbb{P}(A) = \lim_{n \rightarrow \infty} \mathbb{P}(A_n)$, and since $\mathbb{Q}(A) = \lim_{n \rightarrow \infty} \mathbb{Q}(A_n)$, we have $\mathbb{P}(A) = \mathbb{Q}(A)$. ■

July 28th.

Example 1.12. We wish to obtain an example of S which is *not* a π -system, and two probability measures $\mathbb{P}$ and $\mathbb{Q}$ on $(\Sigma, \sigma(S))$ such that $\mathbb{P}(A) = \mathbb{Q}(A)$ for all $A \in S$, but $\mathbb{P}(A) \neq \mathbb{Q}(A)$ for some $A \in \sigma(S)$. Let $\Omega = \{1, 2, 3, 4\}$, and let $S = \{\{1, 2\}, \{2, 3\}\}$. Then, $\sigma(S) = 2^\Omega$. Define $\mathbb{P}$ and $\mathbb{Q}$ on $(\Omega, \sigma(S))$ by

$$\mathbb{P}(\{1\}) = \mathbb{P}(\{2\}) = \mathbb{P}(\{3\}) = \mathbb{P}(\{4\}) = 1/4, \quad (1.14)$$

and

$$\mathbb{Q}(\{1\}) = \mathbb{Q}(\{3\}) = 1/6, \quad \mathbb{Q}(\{2\}) = \mathbb{Q}(\{4\}) = 1/3. \quad (1.15)$$

Then, $\mathbb{P}(\{1, 2\}) = \mathbb{Q}(\{1, 2\}) = 1/2$ and $\mathbb{P}(\{2, 3\}) = \mathbb{Q}(\{2, 3\}) = 1/2$, but $\mathbb{P}(\{3\}) = 1/4 \neq 1/6 = \mathbb{Q}(\{3\})$.

Example 1.13. Let $\Omega \neq \emptyset$, and A_λ be a collection of subsets of Ω indexed by $\lambda \in \Lambda$. Consider the σ -algebra $\sigma(\{A_\lambda\}_{\lambda \in \Lambda})$, and pick $B \in \sigma(\{A_\lambda\}_{\lambda \in \Lambda})$. We will show that there exists a countable subcollection $\{A_{\lambda_n}\}_{n=1}^\infty$ such that $B \in \sigma(\{A_{\lambda_n}\}_{n=1}^\infty)$. Let $\mathcal{C} = \{B \in \sigma(\{A_\lambda\}_{\lambda \in \Lambda}) \mid B \in \sigma(\{A_{\lambda_n}\}_{n=1}^\infty) \text{ for some countable subcollection } \{A_{\lambda_n}\}_{n=1}^\infty\}$. Then, $\mathcal{C}$ is a σ -algebra containing $\{A_\lambda\}_{\lambda \in \Lambda}$, and hence $\sigma(\{A_\lambda\}_{\lambda \in \Lambda}) \subseteq \mathcal{C}$. Thus, $B \in \mathcal{C}$, and hence there exists a countable subcollection $\{A_{\lambda_n}\}_{n=1}^\infty$ such that $B \in \sigma(\{A_{\lambda_n}\}_{n=1}^\infty)$.

1.2 The Lebesgue Measure

Theorem 1.14. *There exists a unique Borel measure λ on $[0, 1]$ such that $\lambda(I) = |I|$ for all intervals $I \subseteq [0, 1]$.*

Proof. For the π -system $S = \{(a, b) \mid 0 \leq a < b \leq 1\}$, we know that $\sigma(S)$ is a Borel σ -algebra on $[0, 1]$. If λ and μ are two Borel measures on $[0, 1]$ such that $\lambda(I) = \mu(I) = |I|$ for all intervals $I \subseteq [0, 1]$, then by the previous lemma, we have $\lambda(A) = \mu(A)$ for all $A \in \sigma(S)$. Thus, the Borel measure is unique.

To show the existence of such a measure, we will define a pseudomeasure λ_* that satisfies the interval property, and then show that it is a measure on the Borel σ -algebra. For any $E \subseteq [0, 1]$, define

$$\lambda_*(E) = \inf \left\{ \sum_{i=1}^\infty |I_i| \mid I_i \text{ are intervals and } E \subseteq \bigcup_{i=1}^\infty I_i \right\} \quad (1.16)$$

Infer that $0 \leq \lambda_*(E) \leq 1$ for all $E \subseteq [0, 1]$. Moreover, let A and B be two sets in $[0, 1]$. Then, for any $\varepsilon > 0$, we have

$$\lambda_*(A \cup B) \leq \sum_{k=1}^\infty |I_k| + \sum_{k=1}^\infty |J_k| \leq \lambda_*(A) + \varepsilon + \lambda_*(B) + \varepsilon \implies \lambda_*(A \cup B) \leq \lambda_*(A) + \lambda_*(B). \quad (1.17)$$

Finally, let $(a, b) \subseteq [0, 1]$ be an interval. Then, $\lambda_*((a, b)) = b - a$. To see this, suppose $(a, b) \subseteq \bigcup_{k=1}^\infty I_k$ for some intervals I_k . Now consider $[a + \varepsilon, b]$ for some $\varepsilon > 0$ such that $[a + \varepsilon, b] \subseteq \bigcup_{k=1}^\infty I_k$. This is a compact subinterval, so we can extract a finite subcover $[a + \varepsilon, b] \subseteq \bigcup_{k=1}^N I_k$. Then,

$$b - a - \varepsilon = |[a + \varepsilon, b]| \leq \sum_{k=1}^N |I_k| \leq \sum_{k=1}^\infty |I_k| \implies b - a \leq \lambda_*((a, b)). \quad (1.18)$$

Now, since $(a, b] \subseteq (a, b + \varepsilon)$, a similar line of reasoning shows that $\lambda_*((a, b]) \leq b - a$. Thus, we have $\lambda_*((a, b]) = b - a$. The property of finite subadditivity also implies the property of countable subadditivity (show this). We now construct the σ -algebra $\mathcal{F}$ on which λ_* is a measure. We pick

$$\mathcal{F} = \{A \subseteq [0, 1] \mid \lambda_*(E) = \lambda_*(E \cap A) + \lambda_*(E \cap A^c) \text{ for all } E \subseteq [0, 1]\}. \quad (1.19)$$

We show that $\mathcal{F}$ is a σ -algebra containing the Borel σ -algebra, and that λ_* is a measure on $\mathcal{F}$. It is left as a minor exercise to show that $\mathcal{F}$ is closed under intersections and complements (making $\mathcal{F}$ a π -system). We show $\mathcal{F}$ is a λ -system. Let $A, B \in \mathcal{F}$. Then $B \setminus A = B \cap A^c$ is in $\mathcal{F}$. Then, for any $E \subseteq [0, 1]$, we have

$$\lambda_*(E) = \lambda_*(E \cap A_n) + \lambda_*(E \cap A_n^c) \quad (1.20)$$

$$\geq \lambda_*(E \cap A_n) + \lambda_*(E \cap A^c), \quad (1.21)$$

because $A^c \subseteq A_n^c$. Since $A_n \uparrow A$, we have $E \cap A_n \uparrow E \cap A$, and hence

$$\lambda_*(E \cap A_n) \uparrow \lambda_*(E \cap A).$$

Passing to the limit gives

$$\lambda_*(E) \geq \lambda_*(E \cap A) + \lambda_*(E \cap A^c).$$

On the other hand, subadditivity of the outer measure gives

$$\lambda_*(E) \leq \lambda_*(E \cap A) + \lambda_*(E \cap A^c).$$

Therefore,

$$\lambda_*(E) = \lambda_*(E \cap A) + \lambda_*(E \cap A^c),$$

so $A \in \mathcal{F}$. ■

July 30th.

Note that $\mathcal{F} \subsetneq 2^{[0,1]}$; that is, there exist subsets of $[0, 1]$ that are not in $\mathcal{F}$. We shall assume that $\lambda_*(E) = \lambda_*(E + x)$ for all $E \subseteq [0, 1]$ and $x \in \mathbb{R}$ such that $E + x \subseteq [0, 1]$. This is a reasonable assumption, since we expect the Lebesgue measure to be translation invariant (one can prove this). Moreover, instead of looking at $[0, 1]$ we shall look at $([0, 1], \oplus)$, where $\oplus$ is addition modulo 1. Now suppose we manage to break $[0, 1]$ into countably many disjoint pieces $A_1, A_2, \dots$ such that $\bigcup_{n=1}^{\infty} A_n = [0, 1]$, and $A_i = A_1 \oplus r_i$ for some $r_i \in [0, 1]$. Then, we would have $\lambda_*(A_i) = \lambda_*(A_1)$ for all i , and hence $\lambda_*([0, 1]) = \sum_{n=1}^{\infty} \lambda_*(A_n) = \sum_{n=1}^{\infty} \lambda_*(A_1)$. This is a contradiction, since $\lambda_*([0, 1]) = 1$ and $\sum_{n=1}^{\infty} \lambda_*(A_1)$ is either 0 or ∞ . Thus, we have shown that there exists a set $A \subseteq [0, 1]$ such that $A \notin \mathcal{F}$.

To construct these A_i 's, simply look at the equivalence relation $\sim$ on $[0, 1]$ defined by $x \sim y$ if and only if $x - y \in \mathbb{Q}$. Then, we can pick a representative from each equivalence class to form a set A_1 . Then, we can define $A_i = A_1 \oplus r_i$ for some $r_i \in [0, 1]$ such that $\{r_i\}_{i=1}^{\infty}$ is a complete set of representatives of $\mathbb{Q} \cap [0, 1]$. This gives us the desired decomposition of $[0, 1]$ into disjoint pieces.

1.2.1 Carathéodory's Extension Theorem

August 4th.

Suppose Ω is non-empty, and $\mathcal{A}$ is an *algebra*; that is, $\mathcal{A}$ is a collection of subsets of Ω such that $\Omega \in \mathcal{A}$, and $\mathcal{A}$ is closed under complements and finite unions. Further say that μ is a *pre-measure* on $\mathcal{A}$; that is, $\mu : \mathcal{A} \rightarrow [0, 1]$ satisfies $\mu(\emptyset) = 0$, and if $A_1, A_2, \dots \in \mathcal{A}$ are pairwise disjoint, then $\mu(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mu(A_i)$. Then, we can extend μ to a measure on the σ -algebra generated by $\mathcal{A}$. This is the content of *Carathéodory's extension theorem*.

Remark 1.15. 1. Note that since $\mathcal{A}$ is a π -system, such an extension must be unique.

2. For a π -system S , a pre-measure on S gives a pre-measure on $\mathcal{A}(S)$, the algebra generated by S , since every set $B \in \mathcal{A}(S)$ can be written as a finite disjoint union of sets in S .

We now wish to assign a 'correct' setting to talk about infinite coin tosses (or simple random walks on $\mathbb{Z}$). Let $\Omega = \{0, 1\}^{\mathbb{N}}$, and we define

$$S = \{\{\mathbf{x} \mid x_1 = a_1, x_2 = a_2, \dots, x_n = a_n\} \mid n \in \mathbb{N}, a_i \in \{0, 1\}\} \cup \{\emptyset\} \quad (1.22)$$

and let $\mathcal{F} = \sigma(S)$. Verify that S is a π -system; we can set

$$\mu(\{\mathbf{x} \mid x_1 = a_1, x_2 = a_2, \dots, x_n = a_n\}) = \prod_{i=1}^n p^{a_i} (1-p)^{1-a_i}. \quad (1.23)$$

For now, assume that μ is a pre-measure on S . Then there'll exist a unique probability measure on $(\Omega, \sigma(S))$ that extends μ .

1.3 Measurable Maps and Random Variables

Note that a space $(\Omega, \mathcal{F})$ is called a *measurable space*.

Definition 1.16. A map $T : (\Omega, \mathcal{F}) \rightarrow (\tilde{\Omega}, \tilde{\mathcal{F}})$ between two measurable spaces is termed a *measurable function* if $T^{-1}(A) \in \mathcal{F}$ for all $A \in \tilde{\mathcal{F}}$. When $(\tilde{\Omega}, \tilde{\mathcal{F}}) = (\mathbb{R}, \mathcal{B})$, we call T a *random variable*.

In most situations, X is usually a topological space and $\mathcal{B}(X)$ is the Borel σ -algebra on X . For the case of $X = \mathbb{R}$, a random variable can be interpreted as a meaningful quantity which can be studied.

Proposition 1.17. $T : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}, \mathcal{B})$ is measurable if and only if $T^{-1}(U) \in \mathcal{F}$ whenever U is open in $\mathbb{R}$.

Proof. One direction is clear; for the converse direction, let $\tilde{\mathcal{F}} = \{W \subseteq \mathbb{R} \mid T^{-1}(W) \in \mathcal{F}\}$. It is relatively easy to show that $\tilde{\mathcal{F}}$ is a σ -algebra on $\mathbb{R}$. Thus, since $\mathcal{B} \subseteq \tilde{\mathcal{F}}$, we have $T^{-1}(A) \in \mathcal{F}$ for all $A \in \mathcal{B}$. ■

Note that the image space being in $\mathbb{R}$ did not play a role in the above proof; for any image space $\tilde{\Omega}$, we can define $\tilde{\mathcal{F}} = \{W \subseteq \tilde{\Omega} \mid T^{-1}(W) \in \mathcal{F}\}$, and show that $\tilde{\mathcal{F}}$ is a σ -algebra on $\tilde{\Omega}$. Such a $\tilde{\mathcal{F}}$ is called the *pushforward σ -algebra* of $\mathcal{F}$ under T , and is the largest σ -algebra on $\tilde{\Omega}$ such that T is measurable. One can think in the converse way, where we are given a map $T : \Omega \rightarrow (\tilde{\Omega}, \tilde{\mathcal{F}})$; to make such a T measurable, one can simply equip Ω with the powerset σ -algebra 2^Ω , and then T is automatically measurable. Alternatively, we can define $\mathcal{F} = \{T^{-1}(A) \mid A \in \tilde{\mathcal{F}}\}$, and then T is measurable. This is called the *pullback σ -algebra* of $\tilde{\mathcal{F}}$ under T , and is the smallest σ -algebra on Ω such that T is measurable.

The most non-trivial yet basic example of a random variable is the *indicator function* of a set $A \in \mathcal{F}$, which is defined as $\mathbf{1}_A(\omega) = 1$ if $\omega \in A$, and 0 otherwise. It is easy to check that $\mathbf{1}_A$ is a random variable.

Lemma 1.18. If $X, Y : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}, \mathcal{B})$ are random variables, then $X + Y$ is also a random variable. More generally, if $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is continuous, then $f(X, Y)$ is a random variable.

August 7th.

Example 1.19. Consider the infinite simple random walk from before; for $a \in \mathbb{Z}$, define $\tau_a = \inf\{n > 0 \mid S_n = a\}$, the first hitting time of a . Then we claim that τ_a is measurable and, hence, a random variable. To see this, note that $\tau_a^{-1}((n, \infty)) = \{\omega \in \Omega \mid \tau_a(\omega) > n\} = \{\omega \in \Omega \mid S_1(\omega) \neq a, S_2(\omega) \neq a, \dots, S_n(\omega) \neq a\} = \bigcap_{k=1}^n \{\omega \in \Omega \mid S_k(\omega) \neq a\}$, which is in $\mathcal{F}$ since S_k is measurable for all k . Thus, τ_a is measurable.

Definition 1.20. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let $T : (\Omega, \mathcal{F}) \rightarrow (\tilde{\Omega}, \tilde{\mathcal{F}})$ be a measurable map. Then, the *pushforward measure* of $\mathbb{P}$ under T is the measure $\tilde{\mathbb{P}}$ on $(\tilde{\Omega}, \tilde{\mathcal{F}})$ defined by $\tilde{\mathbb{P}}(A) = \mathbb{P}(T^{-1}(A))$ for all $A \in \tilde{\mathcal{F}}$.

Indeed, one can show that the pushforward measure $\tilde{\mathbb{P}}$ is a probability measure on $(\tilde{\Omega}, \tilde{\mathcal{F}})$.

Definition 1.21. Given a Borel probability measure μ on $(\mathbb{R}, \mathcal{B})$, we define the *cumulative distribution function* of μ to be the function $F_\mu : \mathbb{R} \rightarrow [0, 1]$ defined by $F_\mu(x) = \mu((-\infty, x])$ for all $x \in \mathbb{R}$.

Proposition 1.22. *With μ as above, the following hold true.*

1. F_μ is non-decreasing.
2. $\lim_{x \rightarrow \infty} F_\mu(x) = 1$ and $\lim_{x \rightarrow -\infty} F_\mu(x) = 0$.
3. F_μ is right-continuous.

Proof. 1. If $x_1 \leq x_2$, then $(-\infty, x_1] \subseteq (-\infty, x_2]$, and hence $F_\mu(x_1) = \mu((-\infty, x_1]) \leq \mu((-\infty, x_2]) = F_\mu(x_2)$.

2. Note that $\mathbb{R} = \bigcup_{n=1}^{\infty} (-\infty, n]$; defining $A_k = (-\infty, k]$, we have $A_1 \subseteq A_2 \subseteq \dots$ and $A_k \uparrow \mathbb{R}$. Since μ is a measure, $\mu(A_k) \uparrow \mu(\mathbb{R}) = 1$, and hence $\lim_{x \rightarrow \infty} F_\mu(x) = 1$. Similarly, $\lim_{x \rightarrow -\infty} F_\mu(x) = 0$.

3. Let x_n be a decreasing sequence of numbers such that $x_n \downarrow x$. Then, $(-\infty, x_1] \supseteq (-\infty, x_2] \supseteq \dots$ and $(-\infty, x_n] \downarrow (-\infty, x]$. Since μ is a measure, $\mu((-\infty, x_n]) \downarrow \mu((-\infty, x])$, and hence $F_\mu(x_n) \downarrow F_\mu(x)$. Thus, F_μ is right-continuous. ■

Theorem 1.23. *Suppose $F : \mathbb{R} \rightarrow [0, 1]$ is a non-decreasing, right-continuous function such that $\lim_{x \rightarrow -\infty} F(x) = 0$ and $\lim_{x \rightarrow \infty} F(x) = 1$. Then, there exists a unique Borel probability measure μ on $(\mathbb{R}, \mathcal{B})$ such that $F = F_\mu$.*

Thus, there is a one-to-one correspondence between Borel probability measures on $(\mathbb{R}, \mathcal{B})$ and cumulative distribution functions. In general, it is easier to study the cumulative distribution function of a probability measure than the measure itself.

Theorem 1.24. *Given any probability measure μ on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$, there exists a measurable map $T : ([0, 1], \mathcal{B}([0, 1]), \lambda) \rightarrow (\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$ such that $\mu = \lambda \circ T^{-1}$. That is, the pushforward measure of λ under T is μ .*

Index

- λ -system, 4
- π -system, 4
- σ -algebra, 2
- algebra, 4, 6
- Borel σ -algebra, 3
- Borel set, 3
- Carathéodory's extension theorem, 6
- cumulative distribution function, 7
- discrete probability space, 1
- indicator function, 7
- measurable function, 7
- measurable space, 7
- pre-measure, 6
- probability measure, 2
- probability space, 2
- pullback σ -algebra, 7
- pushforward σ -algebra, 7
- pushforward measure, 7
- random variable, 7
- Sierpiński-Dynkin π - λ Theorem, 4