

# Analysis of Variance

Cathy Maugis-Rabusseau

INSA Toulouse / IMT  
GMM 116  
cathy.maugis@insa-toulouse.fr

2023-2024

# Outline

- 1 **Introduction**
- 2 One-way ANOVA
- 3 Two-way ANOVA
- 4 Conclusion

- ANOVA = analysis of variance
- Aim: Explain a **quantitative variable**  $Y$  using **qualitative** explanatory variables called **factors**
- The modalities of a factor = **levels** (sub-groups in the sample)

- Here we will not address the issue of **experimental design**, just this vocabulary:

## Definition

- ① A **block** of an experimental design = group of observations associated to a combination of controlled factors
- ② An experimental design is called **full** if there is at least one observation in each block
- ③ An experimental design is called **repeated** if there are several observations per block
- ④ An experimental design is called **balanced** if there is the same number of observations per block

# Outline

- 1 Introduction
- 2 One-way ANOVA**
- 3 Two-way ANOVA
- 4 Conclusion

## 2 One-way ANOVA

- Context and Example
- Regular model
- Singular model
- Predictions, residuals and variance
- Confidence interval
- Test: effect of the factor?

- Data: One quantitative response variable  $Y$  and **one** factor having  $I$  levels
- Notation:
  - $Y_{ij}$  = value for individual  $j$  in group  $i$  (level of the factor)
  - Group  $i$  has  $n_i$  individuals
  - $Y_{i.}$  is the mean value for group  $i$ :  $Y_{i.} = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$
  - $n = \sum_{i=1}^I n_i$  is the total number of individuals
- Question: potential effect of the factor on the response  $Y$  ?  $\Leftrightarrow$   
Difference of the average response per group


# Example

- We are interested in the marks obtained by students in an oral examination.
- Is there a potential effect of the examiner on the mark obtained?

Examiner (i)	A	B	C
Mark $Y_{ij}$	10, 11, 11 12, 13, 15	8, 10, 11, 12 14, 15, 16, 16	10, 13, 14, 14 15, 16, 16
Number $n_i$	6	8	7
Average $\bar{Y}_i$	12	12.75	14



# Example



## 2 One-way ANOVA

- Context and Example
- Regular model
- Singular model
- Predictions, residuals and variance
- Confidence interval
- Test: effect of the factor?

# Regular model

- Regular model:

$$\begin{cases} Y_{ij} = m_i + \varepsilon_{ij}, \quad \forall i = 1, \dots, I, \quad \forall j = 1, \dots, n_i \\ \varepsilon_{ij} \text{ i.i.d } \mathcal{N}(0, \sigma^2) \end{cases}$$

$$Y = \begin{pmatrix} Y_{1,1} \\ \vdots \\ Y_{1n_1} \\ Y_{21} \\ \vdots \\ Y_{In_I} \end{pmatrix} = \begin{pmatrix} 1_{n_1} & 0_{n_1} & 0_{n_1} & \cdots & 0_{n_1} \\ 0_{n_2} & 1_{n_2} & 0_{n_2} & \cdots & 0_{n_2} \\ 0_{n_3} & 0_{n_3} & 1_{n_3} & \cdots & 0_{n_3} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0_{n_I} & 0_{n_I} & 0_{n_I} & \cdots & 1_{n_I} \end{pmatrix} \begin{pmatrix} m_1 \\ m_2 \\ \vdots \\ m_I \end{pmatrix} + \varepsilon$$

with  $\varepsilon \sim \mathcal{N}_n(0_n, \sigma^2 I_n)$ .

- Unknown parameters:  $\theta = (m_1, \dots, m_I)'$  [ $k = I$ ] and  $\sigma^2$ .



- $X'X = \text{diag}(n_1, \dots, n_I)$  is invertible  $\Rightarrow$  regular model
- $\hat{\theta} = (X'X)^{-1}X'Y$  thus  $\hat{m}_i = Y_{i.} = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$

```
anReg<-lm(Marks~Exam -1)
summary(anReg)
```

Call:

```
lm(formula = Marks ~ Exam - 1)
```

Residuals:

Min	1Q	Median	3Q	Max
-4.75	-1.00	0.00	2.00	3.25

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
ExamA	12.0000	0.9789	12.26	3.58e-10 ***
ExamB	12.7500	0.8478	15.04	1.23e-11 ***
ExamC	14.0000	0.9063	15.45	7.88e-12 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.398 on 18 degrees of freedom

Multiple R-squared: 0.9716, Adjusted R-squared: 0.9668

F-statistic: 205 on 3 and 18 DF, p-value: 4.226e-14

# Estimation of $\theta$



```
import statsmodels.api as sm
from statsmodels.formula.api import ols
anRegpy = ols('Marks ~ Exam-1', data=Datapy).fit();
anRegpy.summary()
```

```
<class 'statsmodels.iolib.summary.Summary'>
```

```
"""
                                OLS Regression Results
=====
Dep. Variable:                  Marks    R-squared:                  0.115
Model:                            OLS    Adj. R-squared:              0.017
Method:                           Least Squares    F-statistic:                1.170
Date:                            Mar, 22 aoû 2023    Prob (F-statistic):         0.333
Time:                            09:29:59    Log-Likelihood:             -46.546
No. Observations:                  21    AIC:                        99.09
Df Residuals:                      18    BIC:                        102.2
Df Model:                          2
Covariance Type:                  nonrobust
=====
                                coef    std err          t      P>|t|      [0.025    0.975]
-----
Exam[A]                12.0000     0.979     12.258     0.000     9.943    14.057
Exam[B]                12.7500     0.848     15.039     0.000    10.969    14.531
Exam[C]                14.0000     0.906     15.447     0.000    12.096    15.904
=====
Omnibus:                    0.750    Durbin-Watson:              1.388
Prob(Omnibus):              0.687    Jarque-Bera (JB):           0.773
Skew:                      -0.356    Prob(JB):                   0.679
Kurtosis:                   2.386    Cond. No.                   1.15
=====
```

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

## 2 One-way ANOVA

- Context and Example
- Regular model
- **Singular model**
- Predictions, residuals and variance
- Confidence interval
- Test: effect of the factor?

# Singular model

- For interpretation reasons, we may be interested in an other parametrization

$$Y_{ij} = \mu + \alpha_i + \varepsilon_{ij}, \quad \forall i = 1, \dots, I, \quad \forall j = 1, \dots, n_i$$

where

- $\mu$  = average effect
- $\alpha_i = m_i - \mu$  = differential effect of group  $i$ .
- But this model is over-parameterized [ $I+1$  parameters]  
 $\Rightarrow$  one constraint is required to have an identifiable model
  - Orthogonal constraint :  $\sum_{i=1}^I n_i \alpha_i = 0$
  - By default in R:  $\alpha_1 = 0$


# Estimation of $\theta$ - Orthogonal constraints

- Model:

$$Y_{ij} = \mu + \alpha_i + \varepsilon_{ij}, \quad \theta = (\mu, \alpha_1, \dots, \alpha_I)'$$

- The orthogonal constraint  $\sum_{i=1}^I n_i \alpha_i = 0$

- Estimators: 
$$\begin{cases} \hat{\mu} = Y_{..} \\ \hat{\alpha}_i = Y_{i.} - Y_{..} \end{cases}$$






# Estimation of $\theta$ - By default in R

- Model:

$$Y_{ij} = \mu + \alpha_i + \varepsilon_{ij}, \quad \theta = (\mu, \alpha_1, \dots, \alpha_I)'$$

- The constraint by default in R:  $\alpha_1 = 0$

- Estimators: 
$$\begin{cases} \hat{\mu} = Y_{1.} \\ \hat{\alpha}_i = Y_{i.} - Y_{1.} \end{cases}$$



```
anSing <- lm(Notes~Exam,data=Data)
summary(anSing)
```

Call:

```
lm(formula = Notes ~ Exam, data = Data)
```

Residuals:

Min	1Q	Median	3Q	Max
-4.75	-1.00	0.00	2.00	3.25

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	12.0000	0.9789	12.258	3.58e-10 ***
ExamB	0.7500	1.2950	0.579	0.570
ExamC	2.0000	1.3341	1.499	0.151

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.398 on 18 degrees of freedom

Multiple R-squared: 0.115, Adjusted R-squared: 0.01669

F-statistic: 1.17 on 2 and 18 DF, p-value: 0.333

# Example



```
anSingly = ols('Marks ~ Exam', data=Datapy).fit()
anSingly.summary()
```

```
<class 'statsmodels.iolib.summary.Summary'>
"""
```

## OLS Regression Results

```
=====
Dep. Variable:          Marks      R-squared:                0.115
Model:                  OLS        Adj. R-squared:           0.017
Method:                 Least Squares    F-statistic:           1.170
Date:                  Mar, 22 aoû 2023    Prob (F-statistic):     0.333
Time:                  09:30:00          Log-Likelihood:        -46.546
No. Observations:      21              AIC:                  99.09
Df Residuals:          18              BIC:                  102.2
Df Model:               2
Covariance Type:       nonrobust
=====
```

	coef	std err	t	P> t	[0.025	0.975]
Intercept	12.0000	0.979	12.258	0.000	9.943	14.057
Exam[T.B]	0.7500	1.295	0.579	0.570	-1.971	3.471
Exam[T.C]	2.0000	1.334	1.499	0.151	-0.803	4.803

```
=====
Omnibus:                0.750    Durbin-Watson:           1.388
Prob(Omnibus):          0.687    Jarque-Bera (JB):        0.773
Skew:                  -0.356    Prob(JB):                0.679
Kurtosis:              2.386    Cond. No.:               4.00
=====
```

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

```
"""
```

## 2 One-way ANOVA

- Context and Example
- Regular model
- Singular model
- Predictions, residuals and variance
- Confidence interval
- Test: effect of the factor?

# Predictions, residuals and variance

- Predicted values:  $\hat{Y} = X\hat{\theta}$

$$\Leftrightarrow \forall i, \forall j, \quad \hat{Y}_{ij} = \hat{m}_i = \hat{\mu} + \hat{\alpha}_i = Y_i.$$

- Residuals:  $\hat{\varepsilon} = Y - \hat{Y}$

$$\Leftrightarrow \forall i, \forall j, \quad \hat{\varepsilon}_{ij} = Y_{ij} - \hat{Y}_{ij} = Y_{ij} - Y_i.$$

- Estimator of the variance  $\sigma^2$ :

$$\begin{aligned}\hat{\sigma}^2 &= \frac{\|Y - \hat{Y}\|^2}{n - I} = \frac{1}{n - I} \sum_{i=1}^I \sum_{j=1}^{n_i} (Y_{ij} - \hat{Y}_{ij})^2 \\ &= \frac{1}{n - I} \sum_{i=1}^I \sum_{j=1}^{n_i} (\hat{\varepsilon}_{ij})^2 = \frac{SSR}{n - I}\end{aligned}$$

## Proposition

- The mean of residuals per block is null:  $\forall i = 1, \dots, I, \frac{1}{n_i} \sum_{j=1}^{n_i} \hat{\varepsilon}_{ij} = 0$ .
- The mean of residuals is null:  $\frac{1}{n} \sum_{i=1}^I \sum_{j=1}^{n_i} \hat{\varepsilon}_{ij} = 0$ .
- $\frac{1}{n} \sum_{i=1}^I \sum_{j=1}^{n_i} \hat{Y}_{ij} = \frac{1}{n} \sum_{i=1}^I \sum_{j=1}^{n_i} Y_{ij}$ .
- $\text{cov}(\hat{\varepsilon}, \hat{Y}) = 0$ .
- $\text{var}(Y) = \text{var}(\hat{Y}) + \text{var}(\hat{\varepsilon})$ .

*Proof in exercise*

# Decomposition of the variance

- **Between-group variance :**

$$\text{var}(\hat{Y}) = \sum_{i=1}^I \frac{n_i}{n} (Y_{i.} - Y_{..})^2$$

- **Within-group variance (or residual variance):**

$$\text{var}(\hat{\varepsilon}) = \frac{1}{n} \sum_{i=1}^I \sum_{j=1}^{n_i} (Y_{ij} - Y_{i.})^2 = \frac{1}{n} \sum_{i=1}^I n_i \text{var}_i(Y)$$

- The equality  $\text{var}(Y) = \text{var}(\hat{Y}) + \text{var}(\hat{\varepsilon})$ :  
Total Variance = Between Variance + Within Variance.

# Coefficient of determination $R^2$

The coefficient of determination  $R^2$  is the ratio of the between-group variance on the total variance:

$$R^2 = \frac{\text{var}(\hat{Y})}{\text{var}(Y)} = 1 - \frac{\text{var}(\hat{\varepsilon})}{\text{var}(Y)}.$$

It is a measure of connection between a quantitative variable and a qualitative variable.

Remarks:

- ①  $R^2 = 1 \Leftrightarrow \hat{\varepsilon} = 0_n \Leftrightarrow \forall j = 1, \dots, n_i, Y_{ij} = Y_i.$
- ②  $R^2 = 0 \Leftrightarrow \text{var}(\hat{Y}) = 0 \Leftrightarrow \forall i = 1, \dots, I, Y_{i.} = Y_{..}$



## 2 One-way ANOVA

- Context and Example
- Regular model
- Singular model
- Predictions, residuals and variance
- Confidence interval
- Test: effect of the factor?

# Confidence interval for $m_i$

- $m_i$  is estimated by  $\widehat{m}_i = Y_{i.} \sim \mathcal{N}(m_i, \sigma^2/n_i)$

since  $Y_{ij}$  i.i.d  $\mathcal{N}(m_i, \sigma^2)$  ( $j = 1, \dots, n_i$ )

- By Cochran's theorem,  $\frac{(n-l)\widehat{\sigma}^2}{\sigma^2} \sim \chi^2(n-l)$  and  $\widehat{m}_i \perp\!\!\!\perp \widehat{\sigma}^2$

- We deduce that

$$\sqrt{n_i} \frac{\widehat{m}_i - m_i}{\widehat{\sigma}} \sim \mathcal{T}(n-l).$$

- Let  $t_{n-l, 1-\alpha/2}$  be the  $(1 - \alpha/2)$  quantile of  $\mathcal{T}(n-l)$ . Then,

$$\mathbb{P} \left( m_i \in \left[ \widehat{m}_i \pm t_{n-l, 1-\alpha/2} \sqrt{\frac{\widehat{\sigma}^2}{n_i}} \right] \right) = 1 - \alpha.$$



## • With R:

```
anReg<-lm(Marks~Exam -1)
confint(anReg)
```

	2.5 %	97.5 %
ExamA	9.943313	14.05669
ExamB	10.968857	14.53114
ExamC	12.095878	15.90412

## • With python:

```
anRegpy.conf_int(alpha=0.05)
```

	0	1
Exam[A]	9.943313	14.056687
Exam[B]	10.968857	14.531143
Exam[C]	12.095878	15.904122

# Exercise

- Build the following confidence intervals:

```
anSing<-lm(Marks~Exam)
confint(anSing)
```

	2.5 %	97.5 %
(Intercept)	9.9433129	14.056687
ExamB	-1.9707414	3.470741
ExamC	-0.8027921	4.802792

```
anSingpy.conf_int(alpha=0.05)
```

	0	1
Intercept	9.943313	14.056687
Exam[T.B]	-1.970741	3.470741
Exam[T.C]	-0.802792	4.802792

*Indications: determine the law of  $\hat{\mu} = Y_{1.}$  and  $\hat{\alpha}_i = Y_{i.} - Y_{1.}$ .*

## 2 One-way ANOVA

- Context and Example
- Regular model
- Singular model
- Predictions, residuals and variance
- Confidence interval
- Test: effect of the factor?

# Test: effect of the factor?

- Testing procedure:

$$\mathcal{H}_0 : m_1 = m_2 = \dots = m_l = m \iff \forall i = 1, \dots, l, \alpha_i = 0$$

against

$$\mathcal{H}_1 : \exists(i, i') \text{ such that } m_i \neq m_{i'}.$$

- Fisher's test of the sub-model:

$$(M_0) : Y_{ij} = m + \varepsilon_{ij} \text{ with } \hat{m} = Y_{..} = \frac{1}{n} \sum_{i=1}^l \sum_{j=1}^{n_i} Y_{ij}$$

$$(M_1) : Y_{ij} = m_i + \varepsilon_{ij} = \mu + \alpha_i + \varepsilon_{ij}$$

# Test: effect of the factor?

- Fisher's statistics:

$$F = \frac{\sum_{i=1}^I n_i (Y_{i.} - Y_{..})^2 / (I - 1)}{\sum_{i=1}^I \sum_{j=1}^{n_i} (Y_{ij} - Y_{i.})^2 / (n - I)} = \frac{SSE / (I - 1)}{SSR / (n - I)} \underset{\mathcal{H}_0}{\sim} \mathcal{F}(I - 1, n - I),$$

- We reject  $\mathcal{H}_0$  if  $F > f_{1-\alpha, I-1, n-I}$ .



## • With R:

```
anmequal<-lm(Marks~1)
anova(anmequal,anReg)
```

Analysis of Variance Table

Model 1: Marks ~ 1

Model 2: Marks ~ Exam - 1

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	20	116.95				
2	18	103.50	2	13.452	1.1698	0.333

## • With Python:

```
from statsmodels.stats.anova import anova_lm
anmequalpy = ols('Marks ~1', data=Datapy).fit();
anova_lm(anmequalpy,anRegpy)
```

	df_resid	ssr	df_diff	ss_diff	F	Pr(>F)
0	20.0	116.952381	0.0	NaN	NaN	NaN
1	18.0	103.500000	2.0	13.452381	1.169772	0.332952



# Summary table of one-way anova

	df	Sum of squares	Average of squares	$F$	$f_{1-\alpha}$
Factor	$I - 1$	$SSE = \sum_{i=1}^I n_i (Y_{i.} - Y_{..})^2$	$\frac{SSE}{I-1} = MSE$	$\frac{MSE}{\widehat{\sigma}^2}$	$f_{1-\alpha, I-1, n-I}$
Residual	$n - I$	$SSR = \sum_{i=1}^I \sum_{j=1}^{n_i} (Y_{ij} - Y_{i.})^2$	$\frac{SSR}{n-I} = \widehat{\sigma}^2$		
Total	$n - 1$	$SST = \sum_{i=1}^I \sum_{j=1}^{n_i} (Y_{ij} - Y_{..})^2$			

# Outline

- 1 Introduction
- 2 One-way ANOVA
- 3 Two-way ANOVA**
- 4 Conclusion

## 3 Two-way ANOVA

- Introduction
- Models
- Estimation of  $\theta$
- Predicted values, residuals and variance
- Decomposition of the variability
- Interaction plot
- Testing procedures
- Summary table of two-way anova

## Example (Husson et Pagès, 2013)

In a study of factors influencing wheat yield, three varieties of wheat (L, N and NF) and two nitrogen inputs were compared (normal supply = dose 1, intensive supply = dose 2).

Three repetitions for each couple (variety, dose) were performed and the yield (in q / ha) was measured.

We are interested in the differences that could exist from one variety to another, and in the possible interactions of varieties with nitrogen inputs.

```
summary(Ble)
```

Dose	Variety	Yield
1:9	L :6	Min. :55.65
2:9	N :6	1st Qu.:62.82
	NF:6	Median :65.50
		Mean :66.58
		3rd Qu.:69.75
		Max. :79.83

# Notation

- Two factors (qualitative explanatory variables):
  - First factor = factor A [*Dose*] with  $I [=2]$  levels
  - Second factor = factor B [*Variety*] with  $J [=3]$  levels
- $Y$  = quantitative response variable [*wheat yield*]
  - $Y_{ijk}$  = measure of the  $k$ -th individual for level  $i$  of factor A and level  $j$  of factor B
  - $n_{ij}$  = nb of obs. for level  $i$  of factor A and level  $j$  of factor B
  - $Y_{ij.}$  = mean of observations for block  $(i, j)$
- Notation:

$$Y_{i..} = \frac{1}{n_{i+}} \sum_{j=1}^J \sum_{k=1}^{n_{ij}} Y_{ijk} \text{ with } n_{i+} = \sum_{j=1}^J n_{ij}$$

$$Y_{.j.} = \frac{1}{n_{+j}} \sum_{i=1}^I \sum_{k=1}^{n_{ij}} Y_{ijk} \text{ with } n_{+j} = \sum_{i=1}^I n_{ij}$$

$$Y_{...} = \frac{1}{n} \sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^{n_{ij}} Y_{ijk} \text{ with } n = \sum_{i=1}^I n_{i+} = \sum_{j=1}^J n_{+j}$$

# Example

Variety	L ( $j = 1$ )	N ( $j = 2$ )	NF ( $j = 3$ )
Dose 1 ( $i = 1$ )			
( $Y_{1,j,k}$ )	70.35 63.59	62.56 58.89	69.45 64.84
	79.83	55.65	66.12
$n_{1j} = 3$	$Y_{11.} = 71.26$	$Y_{12.} = 59.03$	$Y_{13.} = 66.80$
Dose 2 ( $i = 2$ )			
( $Y_{2,j,k}$ )	74.97 69.12	58.78 64.39	69.85 64.89
	77.18	60.83	67.15
$n_{2j} = 3$	$Y_{21.} = 73.76$	$Y_{22.} = 61.33$	$Y_{23.} = 67.30$

- We are interested in the effect of **Variety** and **Dose** on **wheat yield** with a possible interaction between the two factors.

## 3 Two-way ANOVA

- Introduction
- Models
- Estimation of  $\theta$
- Predicted values, residuals and variance
- Decomposition of the variability
- Interaction plot
- Testing procedures
- Summary table of two-way anova

# Regular model vs Singular model

- **Regular** model:

$$Y_{ijk} = m_{ij} + \varepsilon_{ijk}, \varepsilon_{ijk} \underset{i.i.d}{\sim} \mathcal{N}(0, \sigma^2)$$

but all the effects are included in  $m_{ij}$

- **Singular** model (with interaction):

$$Y_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk}, \varepsilon_{ijk} \underset{i.i.d}{\sim} \mathcal{N}(0, \sigma^2)$$

distinction of effects ... but constraints are required for parameter estimation.

$1 + I + J + IJ$  parameters and  $IJ$  df thus  $1 + I + J$  constraints



# Parameters of the singular model

The  $IJ$  parameters  $m_{ij}$  are thus decomposed into:

- $\mu$  = centering parameter (intercept),
- $\alpha_i$ ,  $I - 1$  parameters (main effect of factor  $A$ ),
- $\beta_j$ ,  $J - 1$  parameters (main effect of factor  $B$ ),
- $\gamma_{ij}$ ,  $(I - 1)(J - 1)$  parameters (interaction effects of factors).

## Proposition

In the two-way anova framework with interaction, there exist some constraints such that the model is orthogonal if and only if

$$n_{ij} = \frac{n_{i+}n_{+j}}{n}.$$

In this case, the constraints (called Type I) are

$$\sum_{i=1}^I n_{i+} \alpha_i = 0; \sum_{j=1}^J n_{+j} \beta_j = 0; \forall i, \sum_{j=1}^J n_{ij} \gamma_{ij} = 0; \forall j, \sum_{i=1}^I n_{ij} \gamma_{ij} = 0.$$

# Other constraints

- In practice, the following constraints (Type III) may be used:

$$\sum_i \alpha_i = 0, \sum_j \beta_j = 0, \forall i, \sum_j \gamma_{ij} = 0 \text{ et } \forall j, \sum_i \gamma_{ij} = 0$$

With these constraints, the model is orthogonal only if  $n_{ij}$  are constant.

- With R, the constraints by default are

$$\alpha_1 = \beta_1 = 0, \gamma_{1j} = 0 \forall j, \gamma_{i1} = 0 \forall i$$

In the sequel, the model is assumed to be orthogonal.

# Additive two-way ANOVA

- The additive two-way ANOVA model = two-way ANOVA model without interaction

$$Y_{ijk} = \mu + \alpha_i + \beta_j + \varepsilon_{ijk}.$$

- Exercise : Determine the constraints such that this model is orthogonal.

## 3 Two-way ANOVA

- Introduction
- Models
- Estimation of  $\theta$
- Predicted values, residuals and variance
- Decomposition of the variability
- Interaction plot
- Testing procedures
- Summary table of two-way anova

# Estimation - Regular model

- Model:

$$\begin{cases} Y_{ijk} = m_{ij} + \varepsilon_{ijk} \\ \varepsilon_{ijk} \text{ i.i.d } \mathcal{N}(0, \sigma^2) \end{cases} \Leftrightarrow Y = X\theta + \varepsilon \text{ with } \varepsilon \sim \mathcal{N}_n(0_n, \sigma^2 I_n)$$

- $\theta = (m_{11}, \dots, m_{IJ})'$  is estimated by  $\hat{\theta} = (X'X)^{-1}X'Y$

## Proposition

$$\hat{m}_{ij} = \frac{1}{n_{ij}} \sum_{k=1}^{n_{ij}} Y_{ijk} = Y_{ij.} \sim \mathcal{N}\left(m_{ij}, \frac{\sigma^2}{n_{ij}}\right).$$

# Estimation - Singular model

- Model:  $Y_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk}$
- Orthogonal constraints: If  $n_{ij} = \frac{n_{i+}n_{+j}}{n}$ , the orthogonal constraints (type I) are

$$\sum_{i=1}^I n_{i+} \alpha_i = 0; \sum_{j=1}^J n_{+j} \beta_j = 0; \forall i, \sum_{j=1}^J n_{ij} \gamma_{ij} = 0; \forall j, \sum_{i=1}^I n_{ij} \gamma_{ij} = 0.$$

## Proposition

Under the orthogonal constraints,

$$\begin{cases} \hat{\mu} = Y_{...} \\ \hat{\alpha}_i = Y_{i..} - Y_{...} \\ \hat{\beta}_j = Y_{.j.} - Y_{...} \\ \hat{\gamma}_{ij} = Y_{ij.} - Y_{i..} - Y_{.j.} + Y_{...} \end{cases}$$

# Estimation - Singular model

- Model:  $Y_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk}$
- By default in R:  $\alpha_1 = \beta_1 = \gamma_{1j} = \gamma_{i1} = 0, \forall i, \forall j$

## Proposition

$$\begin{cases} \hat{\mu} = Y_{11.} \\ \hat{\alpha}_i = Y_{i1.} - Y_{11.} \\ \hat{\beta}_j = Y_{1j.} - Y_{11.} \\ \hat{\gamma}_{ij} = Y_{ij.} - Y_{i1.} - Y_{1j.} + Y_{11.} \end{cases}$$



```
anov2 = lm(Yield~Dose*Variety,data=Ble)
summary(anov2)
```

Call:

```
lm(formula = Yield ~ Dose * Variety, data = Ble)
```

Residuals:

Min	1Q	Median	3Q	Max
-7.667	-2.296	-0.325	2.623	8.573

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	71.257	2.536	28.101	2.55e-12 ***
Dose2	2.500	3.586	0.697	0.49899
VarietyN	-12.223	3.586	-3.409	0.00519 **
VarietyNF	-4.453	3.586	-1.242	0.23801
Dose2:VarietyN	-0.200	5.071	-0.039	0.96919
Dose2:VarietyNF	-2.007	5.071	-0.396	0.69928

---  
 Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.392 on 12 degrees of freedom

Multiple R-squared: 0.6725, Adjusted R-squared: 0.536

F-statistic: 4.928 on 5 and 12 DF, p-value: 0.01105



# Example

```
import statsmodels.api as sm
from statsmodels.formula.api import ols
Blepy=r.Ble
Blepy['Dose']=Blepy['Dose'].astype(str)
Blepy['Variety']=Blepy['Variety'].astype(str)
anov2Singpy = ols('Yield ~ Dose * Variety', data=Blepy).fit();
anov2Singpy.summary()
```

```
<class 'statsmodels.iolib.summary.Summary'>
"""
```

## OLS Regression Results

```
=====
Dep. Variable:          Yield      R-squared:                0.672
Model:                  OLS        Adj. R-squared:           0.536
Method:                 Least Squares   F-statistic:             4.928
Date:                   Mar, 22 aoü 2023   Prob (F-statistic):      0.0111
Time:                   09:30:01         Log-Likelihood:          -48.528
No. Observations:       18             AIC:                    109.1
Df Residuals:           12             BIC:                    114.4
Df Model:                5
Covariance Type:        nonrobust
=====
```

	coef	std err	t	P> t	[0.025	0.975]
Intercept	71.2567	2.536	28.101	0.000	65.732	76.781
Dose[T.2]	2.5000	3.586	0.697	0.499	-5.313	10.313
Variety[T.N]	-12.2233	3.586	-3.409	0.005	-20.037	-4.410
Variety[T.NF]	-4.4533	3.586	-1.242	0.238	-12.267	3.360
Dose[T.2]:Variety[T.N]	-0.2000	5.071	-0.039	0.969	-11.250	10.850
Dose[T.2]:Variety[T.NF]	-2.0067	5.071	-0.396	0.699	-13.056	9.043

```
=====
Omnibus:                 1.202      Durbin-Watson:           2.764
Prob(Omnibus):            0.548      Jarque-Bera (JB):         0.100
```

## 3 Two-way ANOVA

- Introduction
- Models
- Estimation of  $\theta$
- Predicted values, residuals and variance
- Decomposition of the variability
- Interaction plot
- Testing procedures
- Summary table of two-way anova

# Predicted values, residuals and variance

- Predicted values:

$$\hat{Y}_{ijk} = \hat{m}_{ij} = Y_{ij.} = \hat{\mu} + \hat{\alpha}_i + \hat{\beta}_j + \hat{\gamma}_{ij}$$

- Residuals:

$$\hat{\varepsilon}_{ijk} = Y_{ijk} - Y_{ij.}$$

- Estimator of the variance  $\sigma^2$ :

$$\hat{\sigma}^2 = \frac{1}{n - IJ} \sum_{ijk} (\hat{\varepsilon}_{ijk})^2 = \frac{1}{n - IJ} \sum_{ijk} (Y_{ijk} - Y_{ij.})^2 = \frac{SSR}{n - IJ}$$

and

$$\frac{(n - IJ) \hat{\sigma}^2}{\sigma^2} \sim \chi^2(n - IJ)$$

## 3 Two-way ANOVA

- Introduction
- Models
- Estimation of  $\theta$
- Predicted values, residuals and variance
- Decomposition of the variability
- Interaction plot
- Testing procedures
- Summary table of two-way anova

# Decomposition of the variability

- Decomposition of SST:

$$\underbrace{\sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^{n_{ij}} (Y_{ijk} - Y_{...})^2}_{\text{SST}} = \underbrace{\sum_{i=1}^I \sum_{j=1}^J n_{ij} (Y_{ij.} - Y_{...})^2}_{\text{SSE}} + \underbrace{\sum_{i=1}^I \sum_{j=1}^J n_{ij} \text{var}_{ij}(Y)}_{\text{SSR}}$$

with  $\text{var}_{ij}(Y) = \frac{1}{n_{ij}} \sum_{k=1}^{n_{ij}} (Y_{ijk} - Y_{ij.})^2$ .

- Under the orthogonal constraints,  $\text{SSE} = \text{SSA} + \text{SSB} + \text{SSI}$

$$\text{SSA} = \sum_{i=1}^I n_{i+} (Y_{i..} - Y_{...})^2 = \sum_{i=1}^I n_{i+} (\hat{\alpha}_i)^2$$

$$\text{SSB} = \sum_{j=1}^J n_{+j} (Y_{.j.} - Y_{...})^2 = \sum_{j=1}^J n_{+j} (\hat{\beta}_j)^2$$



$$\text{SSI} = \sum_{i=1}^I \sum_{j=1}^J n_{ij} (Y_{ij.} - Y_{i..} - Y_{.j.} + Y_{...})^2 = \sum_{i=1}^I \sum_{j=1}^J n_{ij} (\hat{\gamma}_{ij})^2$$

## 3 Two-way ANOVA

- Introduction
- Models
- Estimation of  $\theta$
- Predicted values, residuals and variance
- Decomposition of the variability
- Interaction plot
- Testing procedures
- Summary table of two-way anova

# Interaction plot


- Graphical control of the possible interaction





```
attach(Ble)
interaction.plot(Variety,Dose,Yield,col=c(2,4),pch=c(18,24),main="Interaction plot",type="b")
```


## Interaction plot



# Example



```
from statsmodels.graphics.factorplots import interaction_plot  
fig=interaction_plot(Blepy['Variety'],Blepy['Dose'],Blepy['Yield']);
```



## 3 Two-way ANOVA

- Introduction
- Models
- Estimation of  $\theta$
- Predicted values, residuals and variance
- Decomposition of the variability
- Interaction plot
- Testing procedures
- Summary table of two-way anova

- Warning: If there is an interaction between two factors, then the principal effect of each factor must be integrated into the model.
- In order to simplify the model, it is necessary to
  - firstly test if there is an interaction effect,
  - if we retain the model without interaction, we can then test the effect of each factor

# Non-interaction testing

- Hypothesis:

$$\mathcal{H}_I : \gamma_{ij} = 0, \forall i = 1, \dots, I, \forall j = 1, \dots, J$$

- Fisher's test of sub-model:

- $[M_1]$   $Y_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk}$  (model with interaction)
- $[M_0]$   $Y_{ijk} = \mu + \alpha_i + \beta_j + \varepsilon_{ijk}$  (additive model)

- Test statistics:

$$F = \frac{SSI/(I-1)(J-1)}{SSR/(n-IJ)} \underset{\mathcal{H}_0}{\sim} \mathcal{F}((I-1)(J-1), n-IJ).$$



## • With R:

```
anov2add = lm(Yield ~ Variety + Dose, data=Ble)
anova(anov2add, anov2)
```

### Analysis of Variance Table

Model 1: Yield ~ Variety + Dose

Model 2: Yield ~ Dose \* Variety

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	14	235.14				
2	12	231.47	2	3.6654	0.095	0.91

## • With python:

```
from statsmodels.stats.anova import anova_lm
anov2addpy = ols('Yield~Dose + Variety', data=Blepy).fit()
anovaResults = anova_lm(anov2addpy, anov2Singpy)
print(anovaResults)
```

	df_resid	ssr	df_diff	ss_diff	F	Pr(>F)
0	14.0	235.137778	0.0	NaN	NaN	NaN
1	12.0	231.472400	2.0	3.665378	0.09501	0.910041

# Test for the effect of factor A

- Hypothesis:  $\mathcal{H}_A : \alpha_i = 0, \forall i = 1, \dots, I$
- Fisher's testing of sub-model:
  - $[M_1] Y_{ijk} = \mu + \alpha_i + \beta_j + \varepsilon_{ijk}$  (additive model)
  - $[M_0] Y_{ijk} = \mu + \beta_j + \varepsilon_{ijk}$  (one-way ANOVA )
- Test statistics:

$$F = \frac{SSA/(I-1)}{SSRAB/(n-(I+J-1))} \underset{\mathcal{H}_0}{\sim} \mathcal{F}(I-1, n-(I+J-1)),$$

where  $SSRAB$  = residual sum of squares for the additive model.

## • With R:

```
anovA = lm(Yield ~ Variety, data=Ble)
anova(anovA, anov2add)
```

Analysis of Variance Table

Model 1: Yield ~ Variety

Model 2: Yield ~ Variety + Dose

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	15	249.15				
2	14	235.14	1	14.01	0.8341	0.3765

## • With python:

```
anovApy = ols('Yield~Variety', data=Blepy).fit()
print(anova_lm(anovApy, anov2addpy))
```

	df_resid	ssr	df_diff	ss_diff	F	Pr(>F)
0	15.0	249.147467	0.0	NaN	NaN	NaN
1	14.0	235.137778	1.0	14.009689	0.834131	0.376541



# Test for the effect of factor $B$

- Hypothesis:  $\mathcal{H}_B : \beta_j = 0, \forall j = 1, \dots, J$
- Fisher's testing of sub-model:
  - $[M_1]$   $Y_{ijk} = \mu + \alpha_i + \beta_j + \varepsilon_{ijk}$  (additive model)
  - $[M_0]$   $Y_{ijk} = \mu + \alpha_i + \varepsilon_{ijk}$  (one-way ANOVA)
- Test statistics:

$$F = \frac{SSB/(J-1)}{SSRAB/(n-(I+J-1))} \underset{\mathcal{H}_0}{\sim} \mathcal{F}(J-1, n-(I+J-1)),$$

where  $SSRAB$  = residual sum of squares for the additive model.



## • With R:

```
anovB = lm(Yield~Dose,data=Ble)
anova(anovB,anov2add)
```

Analysis of Variance Table

Model 1: Yield ~ Dose

Model 2: Yield ~ Variety + Dose

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	16	692.72				
2	14	235.14	2	457.58	13.622	0.0005192 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

## • With python:

```
anovBpy = ols('Yield~Dose', data=Blepy).fit()
print(anova_lm(anovBpy,anov2addpy))
```

	df_resid	ssr	df_diff	ss_diff	F	Pr(>F)
0	16.0	692.719511	0.0	NaN	NaN	NaN
1	14.0	235.137778	2.0	457.581733	13.622108	0.000519

## 3 Two-way ANOVA

- Introduction
- Models
- Estimation of  $\theta$
- Predicted values, residuals and variance
- Decomposition of the variability
- Interaction plot
- Testing procedures
- Summary table of two-way anova

# Summary table

- Two-way Anova with interaction + Orthogonal design
- Decomposition of the variance

$$SST = SSE + SSR = SSA + SSB + SSI + SSR.$$

Source	df	Sum of squares	Average of squares	$F$	$f_{1-\alpha}$
Factor A	$I - 1$	$SSA$	$MSA = \frac{SSA}{I-1}$	$MSA/\hat{\sigma}^2$	$f_{1-\alpha, I-1, n-IJ}$
Factor B	$J - 1$	$SSB$	$MSB = \frac{SSB}{J-1}$	$MSB/\hat{\sigma}^2$	$f_{1-\alpha, J-1, n-IJ}$
Interaction	$(I - 1)(J - 1)$	$SSI$	$MSI = \frac{SSI}{(I-1)(J-1)}$	$MSI/\hat{\sigma}^2$	$f_{1-\alpha, (I-1)(J-1), n-IJ}$
Residual	$n - IJ$	$SSR$	$\hat{\sigma}^2 = \frac{SSR}{n-IJ}$		
Total	$n - 1$	$SST$			

# Outline

- 1 Introduction
- 2 One-way ANOVA
- 3 Two-way ANOVA
- 4 Conclusion**

## Summary

- Know how to write an ANOVA model with one and two factors (individually and matrix), regular and singular
- Know how to distinguish a regular model from a singular model
- Know how to estimate the parameters of the ANOVA model in the regular case and in the singular case (by adapting to the chosen constraint (s))
- Know how to construct a confidence interval for a parameter of the ANOVA model
- Know how to build a procedure to test the effect of a factor, the interaction effect between factors, ... and know how to organize these tests
- Know how to interpret an interaction plot
- Know how to handle SSA, SSB, SSI, SSE, SSR in the case of an orthogonal design.

# References I

- [1] Jean-Marc Azais and Jean-Marc Bardet. *Le modèle linéaire par l'exemple-2e éd.: Régression, analyse de la variance et plans d'expérience illustrés avec R et SAS*. Dunod, 2012.
- [2] Jean-Jacques Daudin. *Le modèle linéaire et ses extensions-Modèle linéaire général, modèle linéaire généralisé, modèle mixte, plans d'expériences (Niveau C)*. 2015.
- [3] Jean-Jacques Droesbeke, Michel Lejeune, and Gilbert Saporta. *Modèles statistiques pour données qualitatives*. Editions Technip, 2005.
- [4] X Guyon. "Modele linéaire et économétrie". In: *Ellipse, Paris* (2001).
- [5] Bernard Prum. *Modèle linéaire: Comparaison de groupes et régression*. INSERM, 1996.
- [6] Nalini Ravishanker, Zhiyi Chi, and Dipak K Dey. *A first course in linear model theory*. CRC Press, 2021.
- [7] Alvin C Rencher and G Bruce Schaalje. *Linear models in statistics*. John Wiley & Sons, 2008.