

Representation Theory and its Applications in Physics

June 5, 2024

Presented by

Max Varverakis (mvarvera@calpoly.edu)



CAL POLY



Outline:

1. Introduction to Representation Theory
2. Examples in Physics
3. The Braid Group
4. Physical Applications of the Braid Group



CAL POLY

1 Introduction to Representation Theory

Definition of a Representation

Definition

Let G be a group. A *representation* of G is a homomorphism from G to a group of operators on a linear vector space V . The dimension of V is the *dimension* or *degree* of the representation.

Definition of a Representation

Definition

Let G be a group. A *representation* of G is a homomorphism from G to a group of operators on a linear vector space V . The dimension of V is the *dimension* or *degree* of the representation.

If X is a representation of G on a vector space V , then X is a map

$$g \in G \xrightarrow{X} X(g),$$

where $X(g)$ is an operator on the V .

Definition of a Representation

Definition

Let G be a group. A *representation* of G is a homomorphism from G to a group of operators on a linear vector space V . The dimension of V is the *dimension* or *degree* of the representation.

If X is a representation of G on a vector space V , then X is a map

$$g \in G \xrightarrow{X} X(g),$$

where $X(g)$ is an operator on the V .

Remark

If V is finite-dimensional with basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$, then X can be realized as an $n \times n$ matrix.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Consequences:

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Consequences:

1. $X(e) = I$, where e is the identity element of the group and I is the identity operator.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Consequences:

1. $X(e) = I$, where e is the identity element of the group and I is the identity operator.
2. In the matrix presentation of X , $X(g)$ is invertible for all $g \in G$.

Example: The Trivial Representation

Trivial Representation of a Group

For any group G , the trivial representation takes $g \mapsto 1$ for all $g \in G$.

Example: The Trivial Representation

Trivial Representation of a Group

For any group G , the trivial representation takes $g \mapsto 1$ for all $g \in G$.

Comments:

- The trivial representation is always one-dimensional.

Example: The Trivial Representation

Trivial Representation of a Group

For any group G , the trivial representation takes $g \mapsto 1$ for all $g \in G$.

Comments:

- ▶ The trivial representation is always one-dimensional.
- ▶ For groups with more than one element, the trivial representation is not injective, so we call it a *degenerate representation*.

Example: The Trivial Representation

Trivial Representation of a Group

For any group G , the trivial representation takes $g \mapsto 1$ for all $g \in G$.

Comments:

- ▶ The trivial representation is always one-dimensional.
- ▶ For groups with more than one element, the trivial representation is not injective, so we call it a *degenerate representation*.
- ▶ If a representation is injective, then it is a *faithful representation*.

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

- The defining representation of S_n is n -dimensional.

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

- ▶ The defining representation of S_n is n -dimensional.
- ▶ This representation is faithful.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2)$.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2)$.

¹**Definition:** Let X be a representation of R on V_2 with

$$X(\phi)\mathbf{e}_1 = \mathbf{e}_1 \cdot \cos \phi + \mathbf{e}_2 \cdot \sin \phi$$

$$X(\phi)\mathbf{e}_2 = -\mathbf{e}_1 \cdot \sin \phi + \mathbf{e}_2 \cdot \cos \phi$$

¹ \mathbf{e}_1 and \mathbf{e}_2 are orthonormal basis vectors of V_2 .

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2)$.

¹**Definition:** Let X be a representation of R on V_2 with

$$\left. \begin{aligned} X(\phi)\mathbf{e}_1 &= \mathbf{e}_1 \cdot \cos \phi + \mathbf{e}_2 \cdot \sin \phi \\ X(\phi)\mathbf{e}_2 &= -\mathbf{e}_1 \cdot \sin \phi + \mathbf{e}_2 \cdot \cos \phi \end{aligned} \right\} \Rightarrow X(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix}$$

¹ \mathbf{e}_1 and \mathbf{e}_2 are orthonormal basis vectors of V_2 .

Thoughts

Thoughts

- Can you think of other ways to represent 2D rotations?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?
- ▶ How many ways can we represent 2D rotations?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?
- ▶ What does it mean for representations to be equivalent? Unique?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?
- ▶ What does it mean for representations to be equivalent? Unique?

Question

How do we classify representations of a group?

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Definition

The *character* of a representation is the trace of the representation matrix.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if $g \in G$ and X is a representation of G , then the character of $X(g)$ is $\chi(g) = \text{tr}(X(g))$.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if $g \in G$ and X is a representation of G , then the character of $X(g)$ is $\chi(g) = \text{tr}(X(g))$.

- ▶ If two representations have the same character for all $g \in G$, then they are equivalent.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if $g \in G$ and X is a representation of G , then the character of $X(g)$ is $\chi(g) = \text{tr}(X(g))$.

- ▶ If two representations have the same character for all $g \in G$, then they are equivalent.
- ▶ We can use characters to classify representations.

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

²Invariant subspace $W \subset V$: $X(g)\mathbf{w} \in W$, $\forall \mathbf{w} \in W$

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

Comments:

²Invariant subspace $W \subset V$: $X(g)\mathbf{w} \in W, \forall \mathbf{w} \in W$

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

Comments:

- Irreducible representations are the building blocks of all representations.

²Invariant subspace $W \subset V$: $X(g)\mathbf{w} \in W$, $\forall \mathbf{w} \in W$

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

Comments:

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ A reducible representation can be decomposed into a direct sum of irreducible representations.

²Invariant subspace $W \subset V$: $X(g)\mathbf{w} \in W, \forall \mathbf{w} \in W$

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

Comments:

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ A reducible representation can be decomposed into a direct sum of irreducible representations.
- ▶ The decomposition of a representation into irreducibles is unique up to equivalence.

²Invariant subspace $W \subset V$: $X(g)\mathbf{w} \in W, \forall \mathbf{w} \in W$

Example: Irreducible Representation of 2D Rotations

Note: The subspace spanned by \mathbf{e}_1 (or \mathbf{e}_2) is *not* invariant under rotations!

Example: Irreducible Representation of 2D Rotations

Note: The subspace spanned by \mathbf{e}_1 (or \mathbf{e}_2) is *not* invariant under rotations!

Invariance of \mathbf{e}_{\pm}

Let $\mathbf{e}_{\pm} = \frac{1}{\sqrt{2}} (\mp \mathbf{e}_1 + i \mathbf{e}_2)$. Then, $X(\phi) \mathbf{e}_{\pm} = e^{\pm i\phi} \mathbf{e}_{\pm}$.

Example: Irreducible Representation of 2D Rotations

Note: The subspace spanned by \mathbf{e}_1 (or \mathbf{e}_2) is *not* invariant under rotations!

Invariance of \mathbf{e}_{\pm}

Let $\mathbf{e}_{\pm} = \frac{1}{\sqrt{2}} (\mp \mathbf{e}_1 + i \mathbf{e}_2)$. Then, $X(\phi) \mathbf{e}_{\pm} = e^{\pm i\phi} \mathbf{e}_{\pm}$.

Decomposition of X

The span of each \mathbf{e}_{\pm} is an X -invariant subspace of V_2 . In this basis, we rewrite X as a direct sum of the 1D irreducible representations³:

$$X(\phi) = \begin{bmatrix} e^{i\phi} & 0 \\ 0 & e^{-i\phi} \end{bmatrix}.$$

³1-dimensional representations are always irreducible!

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

1. The kernel of T is invariant under $X(G)$.

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

1. The kernel of T is invariant under $X(G)$.
2. The image of T is invariant under $Y(G)$.

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

1. The kernel of T is invariant under $X(G)$.
2. The image of T is invariant under $Y(G)$.
3. Since X and Y are irreducible, $\ker(T) = \{\mathbf{0}\}$ and $\text{im}(T) = V$ or $\ker(T) = V$ and $\text{im}(T) = \{0\}$.

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

1. The kernel of T is invariant under $X(G)$.
2. The image of T is invariant under $Y(G)$.
3. Since X and Y are irreducible, $\ker(T) = \{\mathbf{0}\}$ and $\text{im}(T) = V$ or $\ker(T) = V$ and $\text{im}(T) = \{0\}$.
4. By the rank-nullity theorem, conclude that T is either the zero map or invertible.

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

1. Consider λ to be an eigenvalue of T .

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

1. Consider λ to be an eigenvalue of T .
2. Then $T - \lambda I$ is not invertible.

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

1. Consider λ to be an eigenvalue of T .
2. Then $T - \lambda I$ is not invertible.
3. By assumption, $(T - \lambda I)X(g) = X(g)(T - \lambda I)$ for all $g \in G$.

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

1. Consider λ to be an eigenvalue of T .
2. Then $T - \lambda I$ is not invertible.
3. By assumption, $(T - \lambda I)X(g) = X(g)(T - \lambda I)$ for all $g \in G$.
4. By previous lemma, $T - \lambda I = 0 \implies T = \lambda I$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.
3. Schur's second lemma implies $X(h) = \lambda_h I$ for some scalar λ_h .

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.
3. Schur's second lemma implies $X(h) = \lambda_h I$ for some scalar λ_h .
4. The element h was arbitrary, so $X(g) = \lambda_g I$ for all $g \in G$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.
3. Schur's second lemma implies $X(h) = \lambda_h I$ for some scalar λ_h .
4. The element h was arbitrary, so $X(g) = \lambda_g I$ for all $g \in G$.
5. $X(G)$ is equivalent to the representation $g \mapsto \lambda_g$ for all $g \in G$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.
3. Schur's second lemma implies $X(h) = \lambda_h I$ for some scalar λ_h .
4. The element h was arbitrary, so $X(g) = \lambda_g I$ for all $g \in G$.
5. $X(G)$ is equivalent to the representation $g \mapsto \lambda_g$ for all $g \in G$.
6. One-dimensional representations are irreducible.

A Note About Irreducibility

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums
 - ◇ Tensor products

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums
 - ◇ Tensor products
 - ◇ Complex conjugation⁴

⁴If the representation matrices have entries in \mathbb{C} .

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums
 - ◇ Tensor products
 - ◇ Complex conjugation⁴
 - ◇ Similarity transforms

⁴If the representation matrices have entries in \mathbb{C} .

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums
 - ◇ Tensor products
 - ◇ Complex conjugation⁴
 - ◇ Similarity transforms

How does this help in physics?

Irreducible representations can describe symmetries of physical systems with remarkably fundamental implications.

⁴If the representation matrices have entries in \mathbb{C} .



CAL POLY

2 Examples in Physics

Preliminaries

Outline:

1. Dirac notation.
2. Basic quantum mechanics.
3. Quantum Hilbert space.
4. The commutator.

Preliminaries: Dirac Notation

Preliminaries: Basic Quantum Mechanics

2D Rotations and $SO(2)$

1. The group of 2D rotations is $SO(2)$.
2. General properties of $SO(2)$.

Infinitesimal Rotations

- Go through the derivation of the generator of $SO(2)$ in an appropriate level of detail.

Recovering the Rotation Matrix from J

- Do Taylor expansion thing to get the rotation matrix from J (looks familiar phys majors?)

Irreducible Representations of $SO(2)$

- ▶ Rep generated by J is unitary, J is Hermitian.
- ▶ $SO(2)$ abelian implies 1D irreps (reference previous thm's).
- ▶ Construct 1D invariant subspaces, obtain 1D irreps.
- ▶ Get result about $m \in \mathbb{Z}$ for irrep label.
- ▶ Mention ortho/completeness relations?
- ▶ State vector decomposition. Probably don't have time to delve into detailed derivations but would be great to show part of the argument for getting explicit differential form of J .

Conservation of Angular Momentum

- ▶ Do commutator example with Hamiltonian and J .
- ▶ Discuss implications.
- ▶ We can do the same thing for translation group which gives us the familiar \hat{p} operator and conservation of linear momentum!

Generalization to 3 Spatial Dimensions

- ▶ Show but don't derive $R_n(\theta)$ decomposition into \mathbf{J} components.
- ▶ We have basis from the components of \mathbf{J} .
- ▶ Ladies and gentlemen, we got $SO(3)$...
- ▶ \mathbf{J} component differential forms?
- ▶ Commutation relations, in some form talk about J_{\pm} , J^2 and final eigenvalue results.

Connection to Quantum Mechanics

- ▶ Discuss connection between generators and quantum operators, eigenvalues and classical observables, discretization (!), etc.
- ▶ This is the kicker. I will get very excited here probably.

Multi-valued Irreducible Representations and Spinors

Not sure where to put this. . .

- ▶ Let's come back to $SO(2)$ for a second. . .
- ▶ Show $m = 1/2$ irreps.
- ▶ Discuss implications, spinors, etc. . .



CAL POLY

3 The Braid Group

Basic Definitions

- ▶ Formal definitions.
- ▶ Physical/intuitive visualization and interpretation.
- ▶ Standard generators.
- ▶ Automorphisms of $\pi_1(\mathbb{D}_n)$.
- ▶ Braid relations in this picture.
- ▶ 1D Reps.
- ▶ Burau representation.
- ▶ Note on faithfulness.
- ▶ Unitary representation from reduced Burau.



CAL POLY

4 Physical Applications of the Braid Group

Rotations of Quantum Hilbert Space

- ▶ 1D action on Hilbert space, permuting particles, compare/contrast to bosons/fermions.
- ▶ Talk about nontrivial braiding effects.
- ▶ Example of unitary braid rep acting on Hilbert space.

Anyons: A Consequence of Braiding

- ▶ Introduce anyons.
- ▶ Discuss how anyons are described by the braid group.
- ▶ Fusion rules, abelian vs nonabelian anyons.
- ▶ Non-interacting anyons.
- ▶ Non-interacting anyons in harmonic potential.
- ▶ Nontrivial braiding effects anyone?
- ▶ Applications of anyons! (quantum computing, topological quantum field theory, FQHE, etc.)

Summary/Conclusion

Acknowledgements, questions, references (?)