

Representation Theory and its Applications in Physics

June 5, 2024

CPTitlePage169

Presented by

Max Varverakis (mvarvera@calpoly.edu)

Outline:

1. Introduction to Representation Theory
2. Examples in Physics
3. The Braid Group
4. Physical Applications of the Braid Group

Definition of a Representation

Definition

Let G be a group. A *representation* of G is a homomorphism from G to a group of operators on a linear vector space V . The dimension of V is the *dimension* or *degree* of the representation.

Definition of a Representation

Definition

Let G be a group. A *representation* of G is a homomorphism from G to a group of operators on a linear vector space V . The dimension of V is the *dimension* or *degree* of the representation.

If X is a representation of G on a vector space V , then X is a map

$$g \in G \xrightarrow{X} X(g),$$

where $X(g)$ is an operator on the V .

Definition of a Representation

Definition

Let G be a group. A *representation* of G is a homomorphism from G to a group of operators on a linear vector space V . The dimension of V is the *dimension* or *degree* of the representation.

If X is a representation of G on a vector space V , then X is a map

$$g \in G \xrightarrow{X} X(g),$$

where $X(g)$ is an operator on the V .

Remark

If V is finite-dimensional with basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$, then X can be realized as an $n \times n$ matrix.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Consequences:

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Consequences:

1. $X(e) = I$, where e is the identity element of the group and I is the identity operator.

Properties of Representations

Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

Invertibility

If X is a representation of G , then $X(g)^{-1} = X(g^{-1})$, $\forall g \in G$.

Consequences:

1. $X(e) = I$, where e is the identity element of the group and I is the identity operator.
2. In the matrix presentation of X , $X(g)$ is invertible for all $g \in G$.

Example: The Trivial Representation

Trivial Representation of a Group

For any group G , the trivial representation takes $g \mapsto 1$ for all $g \in G$.

Example: The Trivial Representation

Trivial Representation of a Group

For any group G , the trivial representation takes $g \mapsto 1$ for all $g \in G$.

Comments:

- The trivial representation is always one-dimensional.

Example: The Trivial Representation

Trivial Representation of a Group

For any group G , the trivial representation takes $g \mapsto 1$ for all $g \in G$.

Comments:

- ▶ The trivial representation is always one-dimensional.
- ▶ For groups with more than one element, the trivial representation is not injective, so we call it a *degenerate representation*.

Example: The Trivial Representation

Trivial Representation of a Group

For any group G , the trivial representation takes $g \mapsto 1$ for all $g \in G$.

Comments:

- ▶ The trivial representation is always one-dimensional.
- ▶ For groups with more than one element, the trivial representation is not injective, so we call it a *degenerate representation*.
- ▶ If a representation is injective, then it is a *faithful representation*.

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

- The defining representation of S_n is n -dimensional.

Example: A Faithful Representation of S_n

Defining representation of S_n

The defining representation D of S_n encodes the action of the symmetric group on the standard basis of \mathbb{R}^n . If a permutation sends i to j , then place a 1 the i -th column and j -th row of the representation matrix.

E.g., in S_3 :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

- ▶ The defining representation of S_n is n -dimensional.
- ▶ This representation is faithful.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$.

Identity Element: $R(0) = I$.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$.

Identity Element: $R(0) = I$.

Inverses: $R(\phi)^{-1} = R(-\phi) = R(2\pi - \phi)$.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$.

Identity Element: $R(0) = I$.

Inverses: $R(\phi)^{-1} = R(-\phi) = R(2\pi - \phi)$.

Periodicity Condition: $R(\phi \pm 2\pi) = R(\phi)$.

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$.

Identity Element: $R(0) = I$.

Inverses: $R(\phi)^{-1} = R(-\phi) = R(2\pi - \phi)$.

Periodicity Condition: $R(\phi \pm 2\pi) = R(\phi)$.

Representation: Let X be a representation of R on V_2 with¹

$$X(\phi)\mathbf{e}_1 = \mathbf{e}_1 \cdot \cos \phi + \mathbf{e}_2 \cdot \sin \phi$$

$$X(\phi)\mathbf{e}_2 = -\mathbf{e}_1 \cdot \sin \phi + \mathbf{e}_2 \cdot \cos \phi$$

¹ \mathbf{e}_1 and \mathbf{e}_2 are orthonormal basis vectors of V_2 .

Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let $G = \{R(\phi), 0 \leq \phi < 2\pi\}$ be the group of continuous rotations in the xy -plane (V_2) about the origin.

Group operation: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$.

Identity Element: $R(0) = I$.

Inverses: $R(\phi)^{-1} = R(-\phi) = R(2\pi - \phi)$.

Periodicity Condition: $R(\phi \pm 2\pi) = R(\phi)$.

Representation: Let X be a representation of R on V_2 with¹

$$\left. \begin{aligned} X(\phi)\mathbf{e}_1 &= \mathbf{e}_1 \cdot \cos \phi + \mathbf{e}_2 \cdot \sin \phi \\ X(\phi)\mathbf{e}_2 &= -\mathbf{e}_1 \cdot \sin \phi + \mathbf{e}_2 \cdot \cos \phi \end{aligned} \right\} \Rightarrow \boxed{X(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix}}$$

¹ \mathbf{e}_1 and \mathbf{e}_2 are orthonormal basis vectors of V_2 .

Thoughts

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?
- ▶ How many ways can we represent 2D rotations?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?
- ▶ What does it mean for representations to be equivalent? Unique?

Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about $e^{i\phi}$ parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?
- ▶ What does it mean for representations to be equivalent? Unique?

Question

How do we classify representations of a group?

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Definition

The *character* of a representation is the trace of the representation matrix.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if $g \in G$ and X is a representation of G , then the character of $X(g)$ is $\chi(g) = \text{tr}(X(g))$.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if $g \in G$ and X is a representation of G , then the character of $X(g)$ is $\chi(g) = \text{tr}(X(g))$.

- ▶ If two representations have the same character for all $g \in G$, then they are equivalent.

Equivalent Representations

Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if $g \in G$ and X is a representation of G , then the character of $X(g)$ is $\chi(g) = \text{tr}(X(g))$.

- ▶ If two representations have the same character for all $g \in G$, then they are equivalent.
- ▶ We can use characters to classify representations.

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

²Invariant subspace $W \subset V$: $X(a)\mathbf{w} \in W$. $\forall \mathbf{w} \in W$

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

Comments:

²Invariant subspace $W \subset V$: $X(a)\mathbf{w} \in W$. $\forall \mathbf{w} \in W$

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

Comments:

- ▶ Irreducible representations are the building blocks of all representations.

²Invariant subspace $W \subset V: X(a)\mathbf{w} \in W, \forall \mathbf{w} \in W$

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

Comments:

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ A reducible representation can be decomposed into a direct sum of irreducible representations.

²Invariant subspace $W \subset V$: $X(a)\mathbf{w} \in W$. $\forall \mathbf{w} \in W$

Decomposing Representations

Definition

A representation $X(G)$ on V is *irreducible* if there is no non-trivial invariant subspace² in V with respect to $X(G)$. Otherwise, $X(G)$ is *reducible*.

Comments:

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ A reducible representation can be decomposed into a direct sum of irreducible representations.
- ▶ The decomposition of a representation into irreducibles is unique up to equivalence.

²Invariant subspace $W \subset V$: $X(a)\mathbf{w} \in W$. $\forall \mathbf{w} \in W$

Example: Irreducible Representation of 2D Rotations

Note: The subspace spanned by \mathbf{e}_1 (or \mathbf{e}_2) is *not* invariant under rotations!

Example: Irreducible Representation of 2D Rotations

Note: The subspace spanned by \mathbf{e}_1 (or \mathbf{e}_2) is *not* invariant under rotations!

Invariance of \mathbf{e}_{\pm}

Let $\mathbf{e}_{\pm} = \frac{1}{\sqrt{2}} (\mp \mathbf{e}_1 + i \mathbf{e}_2)$. Then, $X(\phi) \mathbf{e}_{\pm} = e^{\pm i \phi} \mathbf{e}_{\pm}$.

Example: Irreducible Representation of 2D Rotations

Note: The subspace spanned by \mathbf{e}_1 (or \mathbf{e}_2) is *not* invariant under rotations!

Invariance of \mathbf{e}_{\pm}

Let $\mathbf{e}_{\pm} = \frac{1}{\sqrt{2}} (\mp \mathbf{e}_1 + i \mathbf{e}_2)$. Then, $X(\phi) \mathbf{e}_{\pm} = e^{\pm i\phi} \mathbf{e}_{\pm}$.

Decomposition of X

The span of each \mathbf{e}_{\pm} is an X -invariant subspace of V_2 . In this basis, we rewrite X as a direct sum of the 1D irreducible representations³:

$$X(\phi) = \begin{bmatrix} e^{i\phi} & 0 \\ 0 & e^{-i\phi} \end{bmatrix}.$$

³1-dimensional representations are always irreducible!

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

1. The kernel of T is invariant under $X(G)$.

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

1. The kernel of T is invariant under $X(G)$.
2. The image of T is invariant under $Y(G)$.

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

1. The kernel of T is invariant under $X(G)$.
2. The image of T is invariant under $Y(G)$.
3. Since X and Y are irreducible, $\ker(T) = \{\mathbf{0}\}$ and $\text{im}(T) = V$ or $\ker(T) = V$ and $\text{im}(T) = \{0\}$.

Schur's Lemmas (pt. 1)

Lemma

Let $X : G \rightarrow V$ and $Y : G \rightarrow W$ be irreducible representations of a group G . If there exists a fixed linear transformation $T : V \rightarrow W$ such that $TX(g) = Y(g)T$ for all $g \in G$, then T is either the zero map or invertible.

Proof (sketch)

1. The kernel of T is invariant under $X(G)$.
2. The image of T is invariant under $Y(G)$.
3. Since X and Y are irreducible, $\ker(T) = \{\mathbf{0}\}$ and $\text{im}(T) = V$ or $\ker(T) = V$ and $\text{im}(T) = \{0\}$.
4. By the rank-nullity theorem, conclude that T is either the zero map or invertible.

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

1. Consider λ to be an eigenvalue of T .

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

1. Consider λ to be an eigenvalue of T .
2. Then $T - \lambda I$ is not invertible.

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

1. Consider λ to be an eigenvalue of T .
2. Then $T - \lambda I$ is not invertible.
3. By assumption, $(T - \lambda I)X(g) = X(g)(T - \lambda I)$ for all $g \in G$.

Schur's Lemma's (pt. 2)

Lemma

Let X be an irreducible representation of a group G and T a linear operator that commutes with all $X(g)$ for $g \in G$. Then T is a scalar multiple of the identity operator.

Proof (sketch)

1. Consider λ to be an eigenvalue of T .
2. Then $T - \lambda I$ is not invertible.
3. By assumption, $(T - \lambda I)X(g) = X(g)(T - \lambda I)$ for all $g \in G$.
4. By previous lemma, $T - \lambda I = 0 \implies T = \lambda I$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.
3. Schur's second lemma implies $X(h) = \lambda_h I$ for some scalar λ_h .

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.
3. Schur's second lemma implies $X(h) = \lambda_h I$ for some scalar λ_h .
4. The element h was arbitrary, so $X(g) = \lambda_g I$ for all $g \in G$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.
3. Schur's second lemma implies $X(h) = \lambda_h I$ for some scalar λ_h .
4. The element h was arbitrary, so $X(g) = \lambda_g I$ for all $g \in G$.
5. $X(G)$ is equivalent to the representation $g \mapsto \lambda_g$ for all $g \in G$.

Consequence of Schur's Lemmas

Corollary

If G is a finite abelian group, then the irreducible representations of G are one-dimensional.

Proof (sketch)

1. Fix $h \in G$.
2. Since G is abelian, $X(h)X(g) = X(g)X(h)$ for all $g \in G$.
3. Schur's second lemma implies $X(h) = \lambda_h I$ for some scalar λ_h .
4. The element h was arbitrary, so $X(g) = \lambda_g I$ for all $g \in G$.
5. $X(G)$ is equivalent to the representation $g \mapsto \lambda_g$ for all $g \in G$.
6. One-dimensional representations are irreducible.

A Note About Irreducibility

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
 - ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums
 - ◇ Tensor products
-

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums
 - ◇ Tensor products
 - ◇ Complex conjugation⁴

⁴If the representation matrices have entries in \mathbb{C} .

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums
 - ◇ Tensor products
 - ◇ Complex conjugation⁴
 - ◇ Similarity transforms

⁴If the representation matrices have entries in \mathbb{C} .

A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
 - ◇ Direct sums
 - ◇ Tensor products
 - ◇ Complex conjugation⁴
 - ◇ Similarity transforms

How does this help in physics?

Irreducible representations can describe symmetries of physical systems with remarkably fundamental implications.

⁴If the representation matrices have entries in \mathbb{C} .

CPSectionPage169

Preliminaries

Skip preliminaries?

Preliminaries: Physics Conventions

Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.

Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.

Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \qquad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \qquad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \qquad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \qquad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

4. The *Hermitian conjugate* or *adjoint* of an operator A is denoted A^\dagger , and is thought of as complex conjugation and transposition in matrix form.

Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \qquad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

4. The *Hermitian conjugate* or *adjoint* of an operator A is denoted A^\dagger , and is thought of as complex conjugation and transposition in matrix form.
5. Operators that are self-adjoint are called *Hermitian*.

Preliminaries: Dirac notation

Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.

Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted $|\psi\rangle$.

Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted $|\psi\rangle$.
- ▶ A *bra* is a row vector, $\langle\psi|$. This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted $|\psi\rangle$.
- ▶ A *bra* is a row vector, $\langle\psi|$. This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

- ▶ **Inner product:** $\langle\phi|\psi\rangle$

Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted $|\psi\rangle$.
- ▶ A *bra* is a row vector, $\langle\psi|$. This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

- ▶ **Inner product:** $\langle\phi|\psi\rangle$
- ▶ **Outer product:** $|\phi\rangle\langle\psi|$

Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted $|\psi\rangle$.
- ▶ A *bra* is a row vector, $\langle\psi|$. This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

- ▶ **Inner product:** $\langle\phi|\psi\rangle$
- ▶ **Outer product:** $|\phi\rangle\langle\psi|$
- ▶ The action of an operator A on a vector $|\psi\rangle$ is written as $|A\psi\rangle = A|\psi\rangle$.

Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted $|\psi\rangle$.
- ▶ A *bra* is a row vector, $\langle\psi|$. This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

- ▶ **Inner product:** $\langle\phi|\psi\rangle$
- ▶ **Outer product:** $|\phi\rangle\langle\psi|$
- ▶ The action of an operator A on a vector $|\psi\rangle$ is written as $|A\psi\rangle = A|\psi\rangle$.
- ▶ Equivalent ways to write the same thing:

$$\langle A^\dagger \phi | \psi \rangle = \langle \phi | A | \psi \rangle = \langle \phi | A \psi \rangle .$$

Orthonormality, Completeness, and Wavefunctions

Definition

Let $\{|1\rangle, |2\rangle, |3\rangle, \dots\}$ be an orthonormal basis for some quantum Hilbert space. In the context of physics, the **orthonormality** and **completeness** relations of the basis vectors allow any state vector $|\psi\rangle$ to be written as a linear combination of the basis vectors:

$$|\psi\rangle = \left(\sum_n |n\rangle \langle n| \right) |\psi\rangle = \sum_n |n\rangle \langle n|\psi\rangle,$$

where $\sum_n |n\rangle \langle n|$ is the identity operator.

Orthonormality, Completeness, and Wavefunctions

Definition

Let $\{|1\rangle, |2\rangle, |3\rangle, \dots\}$ be an orthonormal basis for some quantum Hilbert space. In the context of physics, the **orthonormality** and **completeness** relations of the basis vectors allow any state vector $|\psi\rangle$ to be written as a linear combination of the basis vectors:

$$|\psi\rangle = \left(\sum_n |n\rangle \langle n| \right) |\psi\rangle = \sum_n |n\rangle \langle n|\psi\rangle,$$

where $\sum_n |n\rangle \langle n|$ is the identity operator.

This is just a fancy change of basis!

Orthonormality, Completeness, and Wavefunctions

Definition

Let $\{|1\rangle, |2\rangle, |3\rangle, \dots\}$ be an orthonormal basis for some quantum Hilbert space. In the context of physics, the **orthonormality** and **completeness** relations of the basis vectors allow any state vector $|\psi\rangle$ to be written as a linear combination of the basis vectors:

$$|\psi\rangle = \left(\sum_n |n\rangle \langle n| \right) |\psi\rangle = \sum_n |n\rangle \langle n|\psi\rangle,$$

where $\sum_n |n\rangle \langle n|$ is the identity operator.

This is just a fancy change of basis!

Definition

For a continuous basis labelled by $|x\rangle$ where x is a continuous parameter, the **wavefunction** $\psi(x)$ is the projection: $\langle x|\psi\rangle = \psi(x)$.

Preliminaries: Basic Quantum Mechanics

- Talk about probabilities and whatnot? Eigenvalues = observables? Or just mention when connecting later stuff to physics?

Properties of 2D Rotations

Let R denote the familiar rotation matrix representation from before.

Properties of 2D Rotations

Let R denote the familiar rotation matrix representation from before.

Definition

An *orthogonal matrix* O satisfies $O^{\top} = O^{-1}$.

Properties of 2D Rotations

Let R denote the familiar rotation matrix representation from before.

Definition

An *orthogonal matrix* O satisfies $O^T = O^{-1}$.

Rotation matrices are orthogonal:

$$R(\phi)R^T(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

Properties of 2D Rotations

Let R denote the familiar rotation matrix representation from before.

Definition

An *orthogonal matrix* O satisfies $O^T = O^{-1}$.

Rotation matrices are orthogonal:

$$R(\phi)R^T(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

Rotations preserve vector lengths:

$$R(\phi)\mathbf{x} = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \phi - x_2 \sin \phi \\ x_1 \sin \phi + x_2 \cos \phi \end{bmatrix} \implies |R(\phi)\mathbf{x}|^2 = |\mathbf{x}|^2.$$

Properties of 2D Rotations

Let R denote the familiar rotation matrix representation from before.

Definition

An *orthogonal matrix* O satisfies $O^\top = O^{-1}$.

Rotation matrices are orthogonal:

$$R(\phi)R^\top(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

Rotations preserve vector lengths:

$$R(\phi)\mathbf{x} = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \phi - x_2 \sin \phi \\ x_1 \sin \phi + x_2 \cos \phi \end{bmatrix} \implies |R(\phi)\mathbf{x}|^2 = |\mathbf{x}|^2.$$

This *special* property is summarized by noting $\det R(\phi) = 1$ for all $\phi \in [0, 2\pi)$.

The $SO(2)$ Group

Definition

The *special orthogonal group* in two dimensions, denoted $SO(2)$, is the group of all 2×2 orthogonal matrices with determinant equal to $+1$.⁵

⁵For all intents and purposes. $SO(2)$ is R from before.

The $SO(2)$ Group

Definition

The *special orthogonal group* in two dimensions, denoted $SO(2)$, is the group of all 2×2 orthogonal matrices with determinant equal to $+1$.⁵

Properties of $SO(2)$:

⁵For all intents and purposes. $SO(2)$ is R from before.

The $SO(2)$ Group

Definition

The *special orthogonal group* in two dimensions, denoted $SO(2)$, is the group of all 2×2 orthogonal matrices with determinant equal to $+1$.⁵

Properties of $SO(2)$:

- The *periodicity condition* $R(\phi + 2\pi) = R(\phi)$ is satisfied.

⁵For all intents and purposes. $SO(2)$ is R from before.

The $SO(2)$ Group

Definition

The *special orthogonal group* in two dimensions, denoted $SO(2)$, is the group of all 2×2 orthogonal matrices with determinant equal to $+1$.⁵

Properties of $SO(2)$:

- ▶ The *periodicity condition* $R(\phi + 2\pi) = R(\phi)$ is satisfied.
- ▶ The *identity element* is $R(0) = I$.

⁵For all intents and purposes. $SO(2)$ is R from before.

The SO(2) Group

Definition

The *special orthogonal group* in two dimensions, denoted $\text{SO}(2)$, is the group of all 2×2 orthogonal matrices with determinant equal to $+1$.⁵

Properties of SO(2):

- ▶ The *periodicity condition* $R(\phi + 2\pi) = R(\phi)$ is satisfied.
- ▶ The *identity element* is $R(0) = I$.
- ▶ $\text{SO}(2)$ is *abelian*: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$.

⁵For all intents and purposes. $\text{SO}(2)$ is R from before.

The SO(2) Group

Definition

The *special orthogonal group* in two dimensions, denoted $\text{SO}(2)$, is the group of all 2×2 orthogonal matrices with determinant equal to $+1$.⁵

Properties of SO(2):

- ▶ The *periodicity condition* $R(\phi + 2\pi) = R(\phi)$ is satisfied.
- ▶ The *identity element* is $R(0) = I$.
- ▶ $\text{SO}(2)$ is *abelian*: $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$.
- ▶ $\text{SO}(2)$ is *reducible* (earlier example with \mathbf{e}_{\pm}).

⁵For all intents and purposes. $\text{SO}(2)$ is R from before.

Infinitesimal Rotations

- Consider an *infinitesimal rotation* labelled by some infinitesimal angle $d\phi$.

Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle $d\phi$.
- ▶ This is equivalent to the identity plus some small rotation, which can be written as⁶

$$R(d\phi) = I - i d\phi J$$

⁶The constant $-i$ is introduced for later convenience. and J is a quantity independent of ϕ .

Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle $d\phi$.
- ▶ This is equivalent to the identity plus some small rotation, which can be written as⁶

$$R(d\phi) = I - i d\phi J$$

- ▶ There are two ways to interpret $R(\phi + d\phi)$:

$$R(\phi + d\phi) = R(\phi)R(d\phi) = R(\phi)(I - id\phi J) = R(\phi) - id\phi R(\phi)J,$$

$$R(\phi + d\phi) = R(\phi) + d\phi \frac{dR(\phi)}{d\phi}$$

⁶The constant $-i$ is introduced for later convenience. and J is a quantity independent of ϕ .

Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle $d\phi$.
- ▶ This is equivalent to the identity plus some small rotation, which can be written as⁶

$$R(d\phi) = I - i d\phi J$$

- ▶ There are two ways to interpret $R(\phi + d\phi)$:

$$R(\phi + d\phi) = R(\phi)R(d\phi) = R(\phi)(I - id\phi J) = R(\phi) - id\phi R(\phi)J,$$

$$R(\phi + d\phi) = R(\phi) + d\phi \frac{dR(\phi)}{d\phi}$$

- ▶ Equating the two expressions gives the differential equation $dR(\phi) = -id\phi R(\phi)J$.

⁶The constant $-i$ is introduced for later convenience. and J is a quantity independent of ϕ .

Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle $d\phi$.
- ▶ This is equivalent to the identity plus some small rotation, which can be written as⁶

$$R(d\phi) = I - i d\phi J$$

- ▶ There are two ways to interpret $R(\phi + d\phi)$:

$$R(\phi + d\phi) = R(\phi)R(d\phi) = R(\phi)(I - id\phi J) = R(\phi) - id\phi R(\phi)J,$$

$$R(\phi + d\phi) = R(\phi) + d\phi \frac{dR(\phi)}{d\phi}$$

- ▶ Equating the two expressions gives the differential equation $dR(\phi) = -id\phi R(\phi)J$.
- ▶ With $R(0) = I$ boundary condition: $R(\phi) = e^{-i\phi J}$.

⁶The constant $-i$ is introduced for later convenience. and J is a quantity independent of ϕ .

Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle $d\phi$.
- ▶ This is equivalent to the identity plus some small rotation, which can be written as⁶

$$R(d\phi) = I - i d\phi J$$

- ▶ There are two ways to interpret $R(\phi + d\phi)$:

$$R(\phi + d\phi) = R(\phi)R(d\phi) = R(\phi)(I - id\phi J) = R(\phi) - id\phi R(\phi)J,$$

$$R(\phi + d\phi) = R(\phi) + d\phi \frac{dR(\phi)}{d\phi}$$

- ▶ Equating the two expressions gives the differential equation $dR(\phi) = -id\phi R(\phi)J$.
- ▶ With $R(0) = I$ boundary condition: $R(\phi) = e^{-i\phi J}$.
- ▶ We call J the *generator* of SO(2) rotations.

⁶The constant $-i$ is introduced for later convenience. and J is a quantity independent of ϕ .

Recovering the Rotation Matrix from J

To first order in $d\phi$: $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

Recovering the Rotation Matrix from J

To first order in $d\phi$: $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before: $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

Recovering the Rotation Matrix from J

To first order in $d\phi$: $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before: $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = -I$

Taylor expand:

$$R(\phi) = e^{-iJ\phi} = I - iJ\phi - I\frac{\phi^2}{2!} + iJ\frac{\phi^3}{3!} + \dots$$

Recovering the Rotation Matrix from J

To first order in $d\phi$: $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before: $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

Taylor expand:

$$\begin{aligned} R(\phi) &= e^{-iJ\phi} = I - iJ\phi - I\frac{\phi^2}{2!} + iJ\frac{\phi^3}{3!} + \dots \\ &= I \left(\sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n}}{(2n)!} \right) - iJ \left(\sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n+1}}{(2n+1)!} \right) \end{aligned}$$

Recovering the Rotation Matrix from J

To first order in $d\phi$: $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before: $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

Taylor expand:

$$\begin{aligned} R(\phi) &= e^{-iJ\phi} = I - iJ\phi - I\frac{\phi^2}{2!} + iJ\frac{\phi^3}{3!} + \dots \\ &= I \left(\sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n}}{(2n)!} \right) - iJ \left(\sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n+1}}{(2n+1)!} \right) \\ &= I \cos \phi - iJ \sin \phi \end{aligned}$$

Recovering the Rotation Matrix from J

To first order in $d\phi$: $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before: $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

Taylor expand:

$$\begin{aligned} R(\phi) &= e^{-iJ\phi} = I - iJ\phi - I\frac{\phi^2}{2!} + iJ\frac{\phi^3}{3!} + \dots \\ &= I \left(\sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n}}{(2n)!} \right) - iJ \left(\sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n+1}}{(2n+1)!} \right) \\ &= I \cos \phi - iJ \sin \phi \\ &= \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix}. \end{aligned}$$

Irreducible Representations of $SO(2)$

Process to obtaining irreducibles:

1. Let U be any representation of $SO(2)$.

Irreducible Representations of $SO(2)$

Process to obtaining irreducibles:

1. Let U be any representation of $SO(2)$.
2. Same argument as before: $U(\phi) = e^{-iJ\phi}$, where J is not necessarily the same as before.

Irreducible Representations of $SO(2)$

Process to obtaining irreducibles:

1. Let U be any representation of $SO(2)$.
2. Same argument as before: $U(\phi) = e^{-iJ\phi}$, where J is not necessarily the same as before.
3. $SO(2)$ is abelian: Schur's Lemmas \implies all irreducible representations are 1D.

Irreducible Representations of SO(2)

Process to obtaining irreducibles:

1. Let U be any representation of $SO(2)$.
2. Same argument as before: $U(\phi) = e^{-iJ\phi}$, where J is not necessarily the same as before.
3. $SO(2)$ is abelian: Schur's Lemmas \implies all irreducible representations are 1D.
4. Each invariant subspace is spanned by an eigenvector of J :

$$J|m\rangle = m|m\rangle ,$$
$$U(\phi)|m\rangle = e^{-iJ\phi}|m\rangle = e^{-im\phi}|m\rangle .$$

Irreducible Representations of $SO(2)$

Process to obtaining irreducibles:

1. Let U be any representation of $SO(2)$.
2. Same argument as before: $U(\phi) = e^{-iJ\phi}$, where J is not necessarily the same as before.
3. $SO(2)$ is abelian: Schur's Lemmas \implies all irreducible representations are 1D.
4. Each invariant subspace is spanned by an eigenvector of J :

$$J|m\rangle = m|m\rangle ,$$
$$U(\phi)|m\rangle = e^{-iJ\phi}|m\rangle = e^{-im\phi}|m\rangle .$$

5. Periodicity of $SO(2) \implies e^{-i2\pi m} = 1 \implies m \in \mathbb{Z}$.

Irreducible Representations of SO(2)

Process to obtaining irreducibles:

1. Let U be any representation of $SO(2)$.
2. Same argument as before: $U(\phi) = e^{-iJ\phi}$, where J is not necessarily the same as before.
3. $SO(2)$ is abelian: Schur's Lemmas \implies all irreducible representations are 1D.
4. Each invariant subspace is spanned by an eigenvector of J :

$$J|m\rangle = m|m\rangle ,$$
$$U(\phi)|m\rangle = e^{-iJ\phi}|m\rangle = e^{-im\phi}|m\rangle .$$

5. Periodicity of $SO(2) \implies e^{-i2\pi m} = 1 \implies m \in \mathbb{Z}$.

Theorem

The single-valued irreducible representations of $SO(2)$ are defined as

$$U^m(\phi) = e^{-im\phi}, \forall m \in \mathbb{Z}.$$

Generalization to 3 Spatial Dimensions

- In 3 spatial dimensions, every rotation can be thought of a rotation in a plane with some perpendicular axis of rotation \mathbf{n} : $R_{\mathbf{n}}(\theta)$.

Generalization to 3 Spatial Dimensions

- ▶ In 3 spatial dimensions, every rotation can be thought of a rotation in a plane with some perpendicular axis of rotation \mathbf{n} : $R_{\mathbf{n}}(\theta)$.
- ▶ Rotations in a plane are isomorphic to $SO(2)$: $R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}}$ for some generator $J_{\mathbf{n}}$.

Generalization to 3 Spatial Dimensions

- ▶ In 3 spatial dimensions, every rotation can be thought of a rotation in a plane with some perpendicular axis of rotation \mathbf{n} : $R_{\mathbf{n}}(\theta)$.
- ▶ Rotations in a plane are isomorphic to $SO(2)$: $R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}}$ for some generator $J_{\mathbf{n}}$.
- ▶ The standard generators along each axis $\{J_x, J_y, J_z\}$ form a basis for all rotation generators: $J_{\mathbf{n}} = n_x J_x + n_y J_y + n_z J_z$.

Generalization to 3 Spatial Dimensions

- ▶ In 3 spatial dimensions, every rotation can be thought of a rotation in a plane with some perpendicular axis of rotation \mathbf{n} : $R_{\mathbf{n}}(\theta)$.
- ▶ Rotations in a plane are isomorphic to $SO(2)$: $R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}}$ for some generator $J_{\mathbf{n}}$.
- ▶ The standard generators along each axis $\{J_x, J_y, J_z\}$ form a basis for all rotation generators: $J_{\mathbf{n}} = n_x J_x + n_y J_y + n_z J_z$.

Consequence: Any rotation in Euclidean 3-space can be written in terms of the generators:

$$R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}} = e^{-i\theta(n_x J_x + n_y J_y + n_z J_z)} = e^{-i\theta \mathbf{n} \cdot \mathbf{J}}.$$

Generalization to 3 Spatial Dimensions

- ▶ In 3 spatial dimensions, every rotation can be thought of a rotation in a plane with some perpendicular axis of rotation \mathbf{n} : $R_{\mathbf{n}}(\theta)$.
- ▶ Rotations in a plane are isomorphic to $SO(2)$: $R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}}$ for some generator $J_{\mathbf{n}}$.
- ▶ The standard generators along each axis $\{J_x, J_y, J_z\}$ form a basis for all rotation generators: $J_{\mathbf{n}} = n_x J_x + n_y J_y + n_z J_z$.

Consequence: Any rotation in Euclidean 3-space can be written in terms of the generators:

$$R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}} = e^{-i\theta(n_x J_x + n_y J_y + n_z J_z)} = e^{-i\theta \mathbf{n} \cdot \mathbf{J}}.$$

Definition

The *special orthogonal group* in three dimensions, denoted $SO(3)$, is the group of all 3×3 orthogonal matrices with determinant equal to $+1$. $SO(3)$ rotations are generated by the components of the Hermitian generator $\mathbf{J} = [J_x, J_y, J_z]^T$.

Connection to Quantum Mechanics

Using a similar process to generate $SO(3)$ invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

Connection to Quantum Mechanics

Using a similar process to generate $SO(3)$ invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

Theorem

*The irreducible representations of $SO(3)$ are labeled by $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$, and the $2j + 1$ eigenvectors spanning an invariant subspace are labelled by their eigenvalues:
 $m = -j, -j + 1, \dots, j - 1, j$.*

Connection to Quantum Mechanics

Using a similar process to generate $SO(3)$ invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

Theorem

*The irreducible representations of $SO(3)$ are labeled by $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$, and the $2j + 1$ eigenvectors spanning an invariant subspace are labelled by their eigenvalues:
 $m = -j, -j + 1, \dots, j - 1, j$.*

Consequences:

Connection to Quantum Mechanics

Using a similar process to generate $SO(3)$ invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

Theorem

*The irreducible representations of $SO(3)$ are labeled by $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$, and the $2j + 1$ eigenvectors spanning an invariant subspace are labelled by their eigenvalues:
 $m = -j, -j + 1, \dots, j - 1, j$.*

Consequences:

- One can obtain the explicit form of \mathbf{J} and subsequently its components J_x, J_y, J_z . These are precisely the angular momentum operators in quantum mechanics.
-

Connection to Quantum Mechanics

Using a similar process to generate $SO(3)$ invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

Theorem

The irreducible representations of $SO(3)$ are labeled by $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$, and the $2j + 1$ eigenvectors spanning an invariant subspace are labelled by their eigenvalues: $m = -j, -j + 1, \dots, j - 1, j$.

Consequences:

- ▶ One can obtain the explicit form of \mathbf{J} and subsequently its components J_x, J_y, J_z . These are precisely the angular momentum operators in quantum mechanics.
- ▶ The eigenvalues of \mathbf{J}^2 and J_z are $j(j + 1)$ and m , respectively⁷. In quantum physics, these eigenvalues correspond to the observable total angular momentum and its z -component.

⁷Typically, the z -axis is chosen as the standard axis.

Connection to Quantum Mechanics

Using a similar process to generate $SO(3)$ invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

Theorem

The irreducible representations of $SO(3)$ are labeled by $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$, and the $2j + 1$ eigenvectors spanning an invariant subspace are labelled by their eigenvalues: $m = -j, -j + 1, \dots, j - 1, j$.

Consequences:

- ▶ One can obtain the explicit form of \mathbf{J} and subsequently its components J_x, J_y, J_z . These are precisely the angular momentum operators in quantum mechanics.
- ▶ The eigenvalues of \mathbf{J}^2 and J_z are $j(j + 1)$ and m , respectively⁷. In quantum physics, these eigenvalues correspond to the observable total angular momentum and its z -component.
- ▶ This generalizes to other types of angular momentum, such as *spin angular momentum*!

⁷Typically, the z -axis is chosen as the standard axis.

Connection to Quantum Mechanics: Punchline

Connection to Quantum Mechanics: Punchline

Discretization of Angular Momentum for Free

Arguably the most defining characteristic of quantum mechanics is that classically measurable quantities become discretized (quantized) when observed on the quantum scale. Without any physical motivation, the irreducible representations of $SO(3)$ gave it to us for free!

Connection to Quantum Mechanics: Punchline

Discretization of Angular Momentum for Free

Arguably the most defining characteristic of quantum mechanics is that classically measurable quantities become discretized (quantized) when observed on the quantum scale. Without any physical motivation, the irreducible representations of $SO(3)$ gave it to us for free!

Now is an appropriate time to let some tears out.

Connection to Quantum Mechanics: Punchline

Discretization of Angular Momentum for Free

Arguably the most defining characteristic of quantum mechanics is that classically measurable quantities become discretized (quantized) when observed on the quantum scale. Without any physical motivation, the irreducible representations of $SO(3)$ gave it to us for free!

Now is an appropriate time to let some tears out.

But that's not all folks!

Conservation of Angular Momentum

1. The *commutator* of two operators A and B is defined as $[A, B] = AB - BA$.

Conservation of Angular Momentum

1. The *commutator* of two operators A and B is defined as $[A, B] = AB - BA$.
2. The *Hamiltonian* operator \hat{H} is the quantum mechanical operator corresponding to the total energy of a system.

Conservation of Angular Momentum

1. The *commutator* of two operators A and B is defined as $[A, B] = AB - BA$.
2. The *Hamiltonian* operator \hat{H} is the quantum mechanical operator corresponding to the total energy of a system.

Theorem (Ehrenfest)

If a time-independent Hermitian operator commutes with the Hamiltonian, then the physical observable corresponding to the operator is conserved.

Conservation of Angular Momentum

1. The *commutator* of two operators A and B is defined as $[A, B] = AB - BA$.
2. The *Hamiltonian* operator \hat{H} is the quantum mechanical operator corresponding to the total energy of a system.

Theorem (Ehrenfest)

If a time-independent Hermitian operator commutes with the Hamiltonian, then the physical observable corresponding to the operator is conserved.

Consequences

1. Any system with radial symmetry is invariant under $SO(3)$ rotations, so $[\hat{H}, \mathbf{J}] = 0$.

Conservation of Angular Momentum

1. The *commutator* of two operators A and B is defined as $[A, B] = AB - BA$.
2. The *Hamiltonian* operator \hat{H} is the quantum mechanical operator corresponding to the total energy of a system.

Theorem (Ehrenfest)

If a time-independent Hermitian operator commutes with the Hamiltonian, then the physical observable corresponding to the operator is conserved.

Consequences

1. Any system with radial symmetry is invariant under $SO(3)$ rotations, so $[\hat{H}, \mathbf{J}] = 0$.
2. Conservation of angular momentum is a direct result of the radial symmetry of the system.

Conservation of Angular Momentum

1. The *commutator* of two operators A and B is defined as $[A, B] = AB - BA$.
2. The *Hamiltonian* operator \hat{H} is the quantum mechanical operator corresponding to the total energy of a system.

Theorem (Ehrenfest)

If a time-independent Hermitian operator commutes with the Hamiltonian, then the physical observable corresponding to the operator is conserved.

Consequences

1. Any system with radial symmetry is invariant under $SO(3)$ rotations, so $[\hat{H}, \mathbf{J}] = 0$.
2. Conservation of angular momentum is a direct result of the radial symmetry of the system.
3. Similar arguments can be made for the continuous group of translations in space, leading to the conservation of linear momentum for translationally invariant systems.

Additional Applications

Additional Applications

1. The $j = 1/2$ irreducible representation of $SO(3)$ describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!

Additional Applications

1. The $j = 1/2$ irreducible representation of $SO(3)$ describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of $SO(3)$ to obtain multi-particle states, arriving at results such as:

Additional Applications

1. The $j = 1/2$ irreducible representation of $SO(3)$ describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of $SO(3)$ to obtain multi-particle states, arriving at results such as:
 - ◇ Clebsch-Gordan coefficients

Additional Applications

1. The $j = 1/2$ irreducible representation of $SO(3)$ describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of $SO(3)$ to obtain multi-particle states, arriving at results such as:
 - ◇ Clebsch-Gordan coefficients
 - ◇ singlet versus triplet states

Additional Applications

1. The $j = 1/2$ irreducible representation of $SO(3)$ describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of $SO(3)$ to obtain multi-particle states, arriving at results such as:
 - ◇ Clebsch-Gordan coefficients
 - ◇ singlet versus triplet states
 - ◇ the Pauli exclusion principle

Additional Applications

1. The $j = 1/2$ irreducible representation of $SO(3)$ describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of $SO(3)$ to obtain multi-particle states, arriving at results such as:
 - ◇ Clebsch-Gordan coefficients
 - ◇ singlet versus triplet states
 - ◇ the Pauli exclusion principle

This is the tip of the iceberg!

CPSectionPage169

The Braid Group

- ▶ Definition: config space and standard visualization

Standard Generators

- ▶ σ_i generators.
- ▶ Define *degree*?
- ▶ Braid relations.
- ▶ Skip YBE verification?

Automorphisms of the Free Group

- ▶ Automorphisms of $\pi_1(\mathbb{D}_n)$.
- ▶ Braid relations in this picture.

One-Dimensional Representations of the Braid Group

One-Dimensional Representations of the Braid Group

For $\theta \in \mathbb{R}$ and $j = 1, 2, \dots, n - 1$, we define some *one-dimensional representations* of B_n :

$$\begin{aligned} p_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

One-Dimensional Representations of the Braid Group

For $\theta \in \mathbb{R}$ and $j = 1, 2, \dots, n - 1$, we define some *one-dimensional representations* of B_n :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*

One-Dimensional Representations of the Braid Group

For $\theta \in \mathbb{R}$ and $j = 1, 2, \dots, n - 1$, we define some *one-dimensional representations* of B_n :

$$\begin{aligned} p_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$p_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) = p_\theta(\sigma_1) p_\theta(\sigma_2) p_\theta(\sigma_1^{-1}) p_\theta(\sigma_2)$$

One-Dimensional Representations of the Braid Group

For $\theta \in \mathbb{R}$ and $j = 1, 2, \dots, n - 1$, we define some *one-dimensional representations* of B_n :

$$\begin{aligned} p_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} p_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= p_\theta(\sigma_1) p_\theta(\sigma_2) p_\theta(\sigma_1^{-1}) p_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \end{aligned}$$

One-Dimensional Representations of the Braid Group

For $\theta \in \mathbb{R}$ and $j = 1, 2, \dots, n - 1$, we define some *one-dimensional representations* of B_n :

$$\begin{aligned} p_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} p_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= p_\theta(\sigma_1) p_\theta(\sigma_2) p_\theta(\sigma_1^{-1}) p_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \\ &= e^{i(\theta_1 - \theta_1 + \theta_2 + \theta_2)} \end{aligned}$$

One-Dimensional Representations of the Braid Group

For $\theta \in \mathbb{R}$ and $j = 1, 2, \dots, n-1$, we define some *one-dimensional representations* of B_n :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} \rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \\ &= e^{i(\theta_1 - \theta_1 + \theta_2 + \theta_2)} \\ &= e^{i \cdot 2\theta_2} = \rho_\theta(\sigma_2^2) \end{aligned}$$

One-Dimensional Representations of the Braid Group

For $\theta \in \mathbb{R}$ and $j = 1, 2, \dots, n - 1$, we define some *one-dimensional representations* of B_n :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} \rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \\ &= e^{i(\theta_1 - \theta_1 + \theta_2 + \theta_2)} \\ &= e^{i \cdot 2\theta_2} = \rho_\theta(\sigma_2^2) \end{aligned}$$

Hence, for any $\beta \in B_n$ with degree k :

One-Dimensional Representations of the Braid Group

For $\theta \in \mathbb{R}$ and $j = 1, 2, \dots, n-1$, we define some *one-dimensional representations* of B_n :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} \rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \\ &= e^{i(\theta_1 - \theta_1 + \theta_2 + \theta_2)} \\ &= e^{i \cdot 2\theta_2} = \rho_\theta(\sigma_2^2) \end{aligned}$$

Hence, for any $\beta \in B_n$ with degree k :

$$\rho_\theta(\beta) = \rho_\theta(\sigma_1^{m_1} \sigma_2^{m_2} \cdots \sigma_{n-1}^{m_{n-1}}) = e^{i\theta(m_1 + m_2 + \cdots + m_{n-1})} = e^{ik\theta}.$$

The Baur Representation

- ▶ Go through arguments/motivation for Baur?
- ▶ Show covering space picture/diagrams?
- ▶ Define Baur representation.
- ▶ Note on faithfulness!
- ▶ Quickly show it's reducible with the $\mathbf{1}$ eigenvector?

Unitary Representation of the Braid Group

- ▶ Define reduced Burau representation.
- ▶ Obtain unitary representation from reduced Burau. (Not sure how much detail to go into here.)
- ▶ Maybe just jump right to defining the unitary reps in the 2×2 case?
- ▶ Comment on why we want a unitary rep!

Nonabelian Characteristics of the Unitary Representation

- ▶ Compare and contrast $\mathcal{U}(\sigma_{1,2})$ to their inverses.
- ▶ Note that $[\mathcal{U}(\sigma_{1,2}), \mathcal{U}(\sigma_{2,1})] \neq 0$ to highlight nonabelian-ness.

Question

What are the physical implications of this nonabelian representation?

CPSectionPage169

(Abelian) Braiding Action on a Quantum System

.

(Abelian) Braiding Action on a Quantum System

1D Representation: Let $p_\theta : B_n \rightarrow \mathbb{C}$ be defined by $\sigma_j \mapsto e^{i\theta}$ for some θ , for all j .

.

(Abelian) Braiding Action on a Quantum System

1D Representation: Let $p_\theta : B_n \rightarrow \mathbb{C}$ be defined by $\sigma_j \mapsto e^{i\theta}$ for some θ , for all j .

Quantum system: Some wavefunction $\psi(r_1, \dots, r_n)$ describing the identical particles fixed at nondegenerate positions r_1, r_2, \dots, r_n .

(Abelian) Braiding Action on a Quantum System

1D Representation: Let $p_\theta : B_n \rightarrow \mathbb{C}$ be defined by $\sigma_j \mapsto e^{i\theta}$ for some θ , for all j .

Quantum system: Some wavefunction $\psi(r_1, \dots, r_n)$ describing the identical particles fixed at nondegenerate positions r_1, r_2, \dots, r_n .

Braiding action: For any degree- k braid $\beta \in B_n$, we have

$$\psi(r_{1'}, r_{2'}, \dots, r_{n'}) = p_\theta(\beta) \psi(r_1, r_2, \dots, r_n) = \underbrace{e^{ik\theta}}_{\text{phase shift}} \psi(r_1, r_2, \dots, r_n),$$

(Nonabelian) Braiding Action on a Quantum System

(Nonabelian) Braiding Action on a Quantum System

2D Representation: Consider the 2×2 unitary representation \mathcal{U} from before.

(Nonabelian) Braiding Action on a Quantum System

2D Representation: Consider the 2×2 unitary representation \mathcal{U} from before.

Quantum system: A degenerate set of two quantum states with orthonormal basis $\psi_1(r_1, r_2, r_3)$ and $\psi_2(r_1, r_2, r_3)$. Shorthand: $|1\rangle$ and $|2\rangle$.

(Nonabelian) Braiding Action on a Quantum System

2D Representation: Consider the 2×2 unitary representation \mathcal{U} from before.

Quantum system: A degenerate set of two quantum states with orthonormal basis $\psi_1(r_1, r_2, r_3)$ and $\psi_2(r_1, r_2, r_3)$. Shorthand: $|1\rangle$ and $|2\rangle$.

Braiding action: The transformed basis states due to the action of σ_1 are

$$\begin{aligned} |1'\rangle &= \mathcal{U}(\sigma_1)_{1,1} |1\rangle + \mathcal{U}(\sigma_1)_{1,2} |2\rangle = \frac{1}{2} e^{-i\frac{\pi}{6}} \left(\sqrt{3} e^{i\arctan\left(\frac{1}{\sqrt{2}}\right)} |1\rangle + |2\rangle \right), \\ |2'\rangle &= \mathcal{U}(\sigma_1)_{2,1} |1\rangle + \mathcal{U}(\sigma_1)_{2,2} |2\rangle = \frac{1}{2} e^{-i\frac{\pi}{6}} \left(|1\rangle - \sqrt{3} e^{-i\arctan\left(\frac{1}{\sqrt{2}}\right)} |2\rangle \right). \end{aligned}$$

(Nonabelian) Braiding Action on a Quantum System

2D Representation: Consider the 2×2 unitary representation \mathcal{U} from before.

Quantum system: A degenerate set of two quantum states with orthonormal basis $\psi_1(r_1, r_2, r_3)$ and $\psi_2(r_1, r_2, r_3)$. Shorthand: $|1\rangle$ and $|2\rangle$.

Braiding action: The transformed basis states due to the action of σ_1 are

$$\begin{aligned} |1'\rangle &= \mathcal{U}(\sigma_1)_{1,1} |1\rangle + \mathcal{U}(\sigma_1)_{1,2} |2\rangle = \frac{1}{2} e^{-i\frac{\pi}{6}} \left(\sqrt{3} e^{i\arctan\left(\frac{1}{\sqrt{2}}\right)} |1\rangle + |2\rangle \right), \\ |2'\rangle &= \mathcal{U}(\sigma_1)_{2,1} |1\rangle + \mathcal{U}(\sigma_1)_{2,2} |2\rangle = \frac{1}{2} e^{-i\frac{\pi}{6}} \left(|1\rangle - \sqrt{3} e^{-i\arctan\left(\frac{1}{\sqrt{2}}\right)} |2\rangle \right). \end{aligned}$$

Remark

The action of a nonabelian braid group representation on a quantum system leads to nontrivial rotations in the many-particle Hilbert space that describes the quantum system.

Anyons: A Consequence of Braiding

Anyons: A Consequence of Braiding

Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- The statistical behavior of anyons is governed by the corresponding braid group representation acting on the system.

Anyons: A Consequence of Braiding

Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- ▶ The statistical behavior of anyons is governed by the corresponding braid group representation acting on the system.
- ▶ Broadly speaking, there are two types of anyons:

Anyons: A Consequence of Braiding

Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- ▶ The statistical behavior of anyons is governed by the corresponding braid group representation acting on the system.
- ▶ Broadly speaking, there are two types of anyons:
 1. *Abelian anyons*: The braid group representation is abelian.

Anyons: A Consequence of Braiding

Definition

Particles that obey the braid group permutation rules are known as *anyons*.


- ▶ The statistical behavior of anyons is governed by the corresponding braid group representation acting on the system.
- ▶ Broadly speaking, there are two types of anyons:
 1. *Abelian anyons*: The braid group representation is abelian.
 2. *Nonabelian anyons*: The braid group representation is nonabelian.

Nontrivial Braiding Effects in 1D Representations

- ▶ Introduce the idea (define it).
- ▶ Show diagram to illustrate nontrivial braiding effects qualitatively.
- ▶ Nontrivial braiding in 1D rep corresponding to diagram.
- ▶ Hint at a greater conclusion but first need to look into the physics perspective. . .

Nontrivial Braiding Effects in 1D Representations


Trajectory A




- ▶ Introduce the idea (define it).
- ▶ Show diagram to illustrate nontrivial braiding effects qualitatively.
- ▶ Nontrivial braiding in 1D rep corresponding to diagram.
- ▶ Hint at a greater conclusion but first need to look into the physics perspective. . .

Nontrivial Braiding Effects in 1D Representations

Trajectory A



Trajectory B





- ▶ Introduce the idea (define it).
- ▶ Show diagram to illustrate nontrivial braiding effects qualitatively.
- ▶ Nontrivial braiding in 1D rep corresponding to diagram.
- ▶ Hint at a greater conclusion but first need to look into the physics perspective. . .



Nontrivial Braiding Effects in 1D Representations

- ▶ Introduce the idea (define it).
- ▶ Show diagram to illustrate nontrivial braiding effects qualitatively.
- ▶ Nontrivial braiding in 1D rep corresponding to diagram.
- ▶ Hint at a greater conclusion but first need to look into the physics perspective...

Trajectory A



Trajectory B



A Physicists Approach to Anyons (pt. 1)

- ▶ Non-interacting anyons.

A Physicists Approach to Anyons (pt. 2)

- ▶ Non-interacting anyons in harmonic potential.
- ▶ Arrive at N -anyon Hamiltonian.

Nontrivial Braiding Effects in the Hamiltonian

- ▶ Compare $N = 2$ to $N = 3$ Hamiltonian from previous slide.
- ▶ Highlight nontrivial braiding effects in the $N = 3$ case.
- ▶ How does this compare to bosons/fermions? (maybe redundant depending on the depth of the previous discussion)
- ▶ Question the physical implications.

Physical Implications of Nontrivial Braiding Effects

- ▶ FQHE
- ▶ Fault-tolerant quantum computing

Summary/Conclusion

- ▶ Outline the talk: what did we talk about?
- ▶ What are the takeaways?
- ▶ Acknowledgements, questions, references (?)