

# Representation Theory and its Applications in Physics

June 5, 2024

**Presented by**

Max Varverakis ([mvarvera@calpoly.edu](mailto:mvarvera@calpoly.edu))



CAL POLY



## Outline:

1. Introduction to Representation Theory
2. Examples in Physics
3. The Braid Group
4. Physical Applications of the Braid Group



**CAL POLY**

## **1 Introduction to Representation Theory**

# Definition of a Representation

## Definition

Let  $G$  be a group. A *representation* of  $G$  is a homomorphism from  $G$  to a group of operators on a linear vector space  $V$ . The dimension of  $V$  is the *dimension* or *degree* of the representation.

# Definition of a Representation

## Definition

Let  $G$  be a group. A *representation* of  $G$  is a homomorphism from  $G$  to a group of operators on a linear vector space  $V$ . The dimension of  $V$  is the *dimension* or *degree* of the representation.

If  $X$  is a representation of  $G$  on a vector space  $V$ , then  $X$  is a map

$$g \in G \xrightarrow{X} X(g),$$

where  $X(g)$  is an operator on the  $V$ .

# Definition of a Representation

## Definition

Let  $G$  be a group. A *representation* of  $G$  is a homomorphism from  $G$  to a group of operators on a linear vector space  $V$ . The dimension of  $V$  is the *dimension* or *degree* of the representation.

If  $X$  is a representation of  $G$  on a vector space  $V$ , then  $X$  is a map

$$g \in G \xrightarrow{X} X(g),$$

where  $X(g)$  is an operator on the  $V$ .

## Remark

If  $V$  is finite-dimensional with basis  $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ , then  $X$  can be realized as an  $n \times n$  matrix.

# Properties of Representations

## Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

## Invertibility

If  $X$  is a representation of  $G$ , then  $X(g)^{-1} = X(g^{-1})$ ,  $\forall g \in G$ .

# Properties of Representations

## Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

## Invertibility

If  $X$  is a representation of  $G$ , then  $X(g)^{-1} = X(g^{-1})$ ,  $\forall g \in G$ .



# Properties of Representations

## Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

## Invertibility

If  $X$  is a representation of  $G$ , then  $X(g)^{-1} = X(g^{-1})$ ,  $\forall g \in G$ .

# Properties of Representations

## Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

## Invertibility

If  $X$  is a representation of  $G$ , then  $X(g)^{-1} = X(g^{-1})$ ,  $\forall g \in G$ .

## Consequences:

# Properties of Representations

## Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

## Invertibility

If  $X$  is a representation of  $G$ , then  $X(g)^{-1} = X(g^{-1})$ ,  $\forall g \in G$ .

## Consequences:

1.  $X(e) = I$ , where  $e$  is the identity element of the group and  $I$  is the identity operator.

# Properties of Representations

## Group Multiplication

Representations are group morphisms, so they satisfy the group multiplication rule:

$$X(gh) = X(g)X(h), \quad \forall g, h \in G.$$

## Invertibility

If  $X$  is a representation of  $G$ , then  $X(g)^{-1} = X(g^{-1})$ ,  $\forall g \in G$ .

## Consequences:

1.  $X(e) = I$ , where  $e$  is the identity element of the group and  $I$  is the identity operator.
2. In the matrix presentation of  $X$ ,  $X(g)$  is invertible for all  $g \in G$ .

# Example: The Trivial Representation

## Trivial Representation of a Group

For any group  $G$ , the trivial representation takes  $g \mapsto 1$  for all  $g \in G$ .

# Example: The Trivial Representation

## Trivial Representation of a Group

For any group  $G$ , the trivial representation takes  $g \mapsto 1$  for all  $g \in G$ .

### Comments:

- ▶ The trivial representation is always one-dimensional.

# Example: The Trivial Representation

## Trivial Representation of a Group

For any group  $G$ , the trivial representation takes  $g \mapsto 1$  for all  $g \in G$ .

### Comments:

- ▶ The trivial representation is always one-dimensional.
- ▶ For groups with more than one element, the trivial representation is not injective, so we call it a *degenerate representation*.

# Example: The Trivial Representation

## Trivial Representation of a Group

For any group  $G$ , the trivial representation takes  $g \mapsto 1$  for all  $g \in G$ .

### Comments:

- ▶ The trivial representation is always one-dimensional.
- ▶ For groups with more than one element, the trivial representation is not injective, so we call it a *degenerate representation*.
- ▶ If a representation is injective, then it is a *faithful representation*.



# Example: A Faithful Representation of $S_n$

## Defining representation of $S_n$

The defining representation  $D$  of  $S_n$  encodes the action of the symmetric group on the standard basis of  $\mathbb{R}^n$ . If a permutation sends  $i$  to  $j$ , then place a 1 the  $i$ -th column and  $j$ -th row of the representation matrix.

# Example: A Faithful Representation of $S_n$

## Defining representation of $S_n$

The defining representation  $D$  of  $S_n$  encodes the action of the symmetric group on the standard basis of  $\mathbb{R}^n$ . If a permutation sends  $i$  to  $j$ , then place a 1 the  $i$ -th column and  $j$ -th row of the representation matrix.

E.g., in  $S_3$ :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

# Example: A Faithful Representation of $S_n$

## Defining representation of $S_n$

The defining representation  $D$  of  $S_n$  encodes the action of the symmetric group on the standard basis of  $\mathbb{R}^n$ . If a permutation sends  $i$  to  $j$ , then place a 1 the  $i$ -th column and  $j$ -th row of the representation matrix.

E.g., in  $S_3$ :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

# Example: A Faithful Representation of $S_n$

## Defining representation of $S_n$

The defining representation  $D$  of  $S_n$  encodes the action of the symmetric group on the standard basis of  $\mathbb{R}^n$ . If a permutation sends  $i$  to  $j$ , then place a 1 the  $i$ -th column and  $j$ -th row of the representation matrix.

E.g., in  $S_3$ :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

# Example: A Faithful Representation of $S_n$

## Defining representation of $S_n$

The defining representation  $D$  of  $S_n$  encodes the action of the symmetric group on the standard basis of  $\mathbb{R}^n$ . If a permutation sends  $i$  to  $j$ , then place a 1 the  $i$ -th column and  $j$ -th row of the representation matrix.

E.g., in  $S_3$ :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

- The defining representation of  $S_n$  is  $n$ -dimensional.

# Example: A Faithful Representation of $S_n$

## Defining representation of $S_n$

The defining representation  $D$  of  $S_n$  encodes the action of the symmetric group on the standard basis of  $\mathbb{R}^n$ . If a permutation sends  $i$  to  $j$ , then place a 1 the  $i$ -th column and  $j$ -th row of the representation matrix.

E.g., in  $S_3$ :

$$D((23)) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad D((123)) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

- ▶ The defining representation of  $S_n$  is  $n$ -dimensional.
- ▶ This representation is faithful.

# Example: Representation of Continuous Rotation Group

*Representations also work for continuous groups!*

# Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let  $G = \{R(\phi), 0 \leq \phi < 2\pi\}$  be the group of continuous rotations in the  $xy$ -plane ( $V_2$ ) about the origin.



# Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let  $G = \{R(\phi), 0 \leq \phi < 2\pi\}$  be the group of continuous rotations in the  $xy$ -plane ( $V_2$ ) about the origin.

**Group operation:**  $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$ .

# Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let  $G = \{R(\phi), 0 \leq \phi < 2\pi\}$  be the group of continuous rotations in the  $xy$ -plane ( $V_2$ ) about the origin.

**Group operation:**  $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$ .

**Identity Element:**  $R(0) = I$ .

# Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let  $G = \{R(\phi), 0 \leq \phi < 2\pi\}$  be the group of continuous rotations in the  $xy$ -plane ( $V_2$ ) about the origin.

**Group operation:**  $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$ .

**Identity Element:**  $R(0) = I$ .

**Inverses:**  $R(\phi)^{-1} = R(-\phi) = R(2\pi - \phi)$ .

# Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let  $G = \{R(\phi), 0 \leq \phi < 2\pi\}$  be the group of continuous rotations in the  $xy$ -plane ( $V_2$ ) about the origin.

**Group operation:**  $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$ .

**Identity Element:**  $R(0) = I$ .

**Inverses:**  $R(\phi)^{-1} = R(-\phi) = R(2\pi - \phi)$ .

**Periodicity Condition:**  $R(\phi \pm 2\pi) = R(\phi)$ .

# Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let  $G = \{R(\phi), 0 \leq \phi < 2\pi\}$  be the group of continuous rotations in the  $xy$ -plane ( $V_2$ ) about the origin.

**Group operation:**  $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$ .

**Identity Element:**  $R(0) = I$ .

**Inverses:**  $R(\phi)^{-1} = R(-\phi) = R(2\pi - \phi)$ .

**Periodicity Condition:**  $R(\phi \pm 2\pi) = R(\phi)$ .

**Representation:** Let  $X$  be a representation of  $G$  on  $V_2$  with<sup>1</sup>

$$X(\phi)\mathbf{e}_1 = \mathbf{e}_1 \cdot \cos \phi + \mathbf{e}_2 \cdot \sin \phi$$

$$X(\phi)\mathbf{e}_2 = -\mathbf{e}_1 \cdot \sin \phi + \mathbf{e}_2 \cdot \cos \phi$$

---

<sup>1</sup> $\mathbf{e}_1$  and  $\mathbf{e}_2$  are orthonormal basis vectors of  $V_2$ .

# Example: Representation of Continuous Rotation Group

Representations also work for continuous groups!

Let  $G = \{R(\phi), 0 \leq \phi < 2\pi\}$  be the group of continuous rotations in the  $xy$ -plane ( $V_2$ ) about the origin.

**Group operation:**  $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$ .

**Identity Element:**  $R(0) = I$ .

**Inverses:**  $R(\phi)^{-1} = R(-\phi) = R(2\pi - \phi)$ .

**Periodicity Condition:**  $R(\phi \pm 2\pi) = R(\phi)$ .

**Representation:** Let  $X$  be a representation of  $G$  on  $V_2$  with<sup>1</sup>

$$\left. \begin{aligned} X(\phi)\mathbf{e}_1 &= \mathbf{e}_1 \cdot \cos \phi + \mathbf{e}_2 \cdot \sin \phi \\ X(\phi)\mathbf{e}_2 &= -\mathbf{e}_1 \cdot \sin \phi + \mathbf{e}_2 \cdot \cos \phi \end{aligned} \right\} \Rightarrow X(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix}$$

---

<sup>1</sup> $\mathbf{e}_1$  and  $\mathbf{e}_2$  are orthonormal basis vectors of  $V_2$ .

# Thoughts

# Thoughts

- Can you think of other ways to represent 2D rotations?



# Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about  $e^{i\phi}$  parameterization?

# Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about  $e^{i\phi}$  parameterization?
- ▶ How many ways can we represent 2D rotations?

# Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about  $e^{i\phi}$  parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?

# Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about  $e^{i\phi}$  parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?
- ▶ What does it mean for representations to be equivalent? Unique?

# Thoughts

- ▶ Can you think of other ways to represent 2D rotations?
- ▶ What about  $e^{i\phi}$  parameterization?
- ▶ How many ways can we represent 2D rotations?
- ▶ Are certain representations equivalent?
- ▶ What does it mean for representations to be equivalent? Unique?

## Question

How do we classify representations of a group?

# Equivalent Representations

## Definition

Two representations are *equivalent* if they are related by a similarity transformation.

# Equivalent Representations

## Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.

# Equivalent Representations

## Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.



# Equivalent Representations

## Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

## Definition

The *character* of a representation is the trace of the representation matrix.

# Equivalent Representations

## Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

## Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if  $g \in G$  and  $X$  is a representation of  $G$ , then the character of  $X(g)$  is  $\chi(g) = \text{tr}(X(g))$ .

# Equivalent Representations

## Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

## Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if  $g \in G$  and  $X$  is a representation of  $G$ , then the character of  $X(g)$  is  $\chi(g) = \text{tr}(X(g))$ .

- ▶ If two representations have the same character for all  $g \in G$ , then they are equivalent.

# Equivalent Representations

## Definition

Two representations are *equivalent* if they are related by a similarity transformation.

- ▶ If two representations are equivalent, then their matrix forms have the same *trace*.
- ▶ Equivalent representations form an equivalence class.

## Definition

The *character* of a representation is the trace of the representation matrix.

E.g., if  $g \in G$  and  $X$  is a representation of  $G$ , then the character of  $X(g)$  is  $\chi(g) = \text{tr}(X(g))$ .

- ▶ If two representations have the same character for all  $g \in G$ , then they are equivalent.
- ▶ We can use characters to classify representations.

# Decomposing Representations

## Definition

A representation  $X(G)$  on  $V$  is *irreducible* if there is no non-trivial invariant subspace<sup>2</sup> in  $V$  with respect to  $X(G)$ . Otherwise,  $X(G)$  is *reducible*.

---

<sup>2</sup>Invariant subspace  $W \subset V$ :  $X(g)\mathbf{w} \in W$ ,  $\forall \mathbf{w} \in W$

# Decomposing Representations

## Definition

A representation  $X(G)$  on  $V$  is *irreducible* if there is no non-trivial invariant subspace<sup>2</sup> in  $V$  with respect to  $X(G)$ . Otherwise,  $X(G)$  is *reducible*.

## Comments:

---

<sup>2</sup>Invariant subspace  $W \subset V$ :  $X(g)\mathbf{w} \in W$ ,  $\forall \mathbf{w} \in W$

# Decomposing Representations

## Definition

A representation  $X(G)$  on  $V$  is *irreducible* if there is no non-trivial invariant subspace<sup>2</sup> in  $V$  with respect to  $X(G)$ . Otherwise,  $X(G)$  is *reducible*.

## Comments:

- Irreducible representations are the building blocks of all representations.

---

<sup>2</sup>Invariant subspace  $W \subset V$ :  $X(g)\mathbf{w} \in W$ ,  $\forall \mathbf{w} \in W$

# Decomposing Representations

## Definition

A representation  $X(G)$  on  $V$  is *irreducible* if there is no non-trivial invariant subspace<sup>2</sup> in  $V$  with respect to  $X(G)$ . Otherwise,  $X(G)$  is *reducible*.

## Comments:

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ A reducible representation can be decomposed into a direct sum of irreducible representations.

---

<sup>2</sup>Invariant subspace  $W \subset V$ :  $X(g)\mathbf{w} \in W, \forall \mathbf{w} \in W$



# Decomposing Representations

## Definition

A representation  $X(G)$  on  $V$  is *irreducible* if there is no non-trivial invariant subspace<sup>2</sup> in  $V$  with respect to  $X(G)$ . Otherwise,  $X(G)$  is *reducible*.

## Comments:

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ A reducible representation can be decomposed into a direct sum of irreducible representations.
- ▶ The decomposition of a representation into irreducibles is unique up to equivalence.

---

<sup>2</sup>Invariant subspace  $W \subset V$ :  $X(g)\mathbf{w} \in W$ ,  $\forall \mathbf{w} \in W$

# Example: Irreducible Representation of 2D Rotations

**Note:** The subspace spanned by  $\mathbf{e}_1$  (or  $\mathbf{e}_2$ ) is *not* invariant under rotations!

# Example: Irreducible Representation of 2D Rotations

**Note:** The subspace spanned by  $\mathbf{e}_1$  (or  $\mathbf{e}_2$ ) is *not* invariant under rotations!

## Invariance of $\mathbf{e}_{\pm}$

Let  $\mathbf{e}_{\pm} = \frac{1}{\sqrt{2}} (\mp \mathbf{e}_1 + i \mathbf{e}_2)$ . Then,  $X(\phi) \mathbf{e}_{\pm} = e^{\pm i \phi} \mathbf{e}_{\pm}$ .

# Example: Irreducible Representation of 2D Rotations

**Note:** The subspace spanned by  $\mathbf{e}_1$  (or  $\mathbf{e}_2$ ) is *not* invariant under rotations!

## Invariance of $\mathbf{e}_{\pm}$

Let  $\mathbf{e}_{\pm} = \frac{1}{\sqrt{2}} (\mp \mathbf{e}_1 + i \mathbf{e}_2)$ . Then,  $X(\phi) \mathbf{e}_{\pm} = e^{\pm i\phi} \mathbf{e}_{\pm}$ .

## Decomposition of $X$

The span of each  $\mathbf{e}_{\pm}$  is an  $X$ -invariant subspace of  $V_2$ . In this basis, we rewrite  $X$  as a direct sum of the 1D irreducible representations<sup>3</sup>:

$$X(\phi) = \begin{bmatrix} e^{i\phi} & 0 \\ 0 & e^{-i\phi} \end{bmatrix}.$$

---

<sup>3</sup>1-dimensional representations are always irreducible!

# Schur's Lemmas (pt. 1)

## Lemma

*Let  $X : G \rightarrow V$  and  $Y : G \rightarrow W$  be irreducible representations of a group  $G$ . If there exists a fixed linear transformation  $T : V \rightarrow W$  such that  $TX(g) = Y(g)T$  for all  $g \in G$ , then  $T$  is either the zero map or invertible.*

# Schur's Lemmas (pt. 1)

## Lemma

*Let  $X : G \rightarrow V$  and  $Y : G \rightarrow W$  be irreducible representations of a group  $G$ . If there exists a fixed linear transformation  $T : V \rightarrow W$  such that  $TX(g) = Y(g)T$  for all  $g \in G$ , then  $T$  is either the zero map or invertible.*

## Proof (sketch)

# Schur's Lemmas (pt. 1)

## Lemma

*Let  $X : G \rightarrow V$  and  $Y : G \rightarrow W$  be irreducible representations of a group  $G$ . If there exists a fixed linear transformation  $T : V \rightarrow W$  such that  $TX(g) = Y(g)T$  for all  $g \in G$ , then  $T$  is either the zero map or invertible.*

## Proof (sketch)

1. The kernel of  $T$  is invariant under  $X(G)$ .

# Schur's Lemmas (pt. 1)

## Lemma

*Let  $X : G \rightarrow V$  and  $Y : G \rightarrow W$  be irreducible representations of a group  $G$ . If there exists a fixed linear transformation  $T : V \rightarrow W$  such that  $TX(g) = Y(g)T$  for all  $g \in G$ , then  $T$  is either the zero map or invertible.*

## Proof (sketch)

1. The kernel of  $T$  is invariant under  $X(G)$ .
2. The image of  $T$  is invariant under  $Y(G)$ .



# Schur's Lemmas (pt. 1)

## Lemma

*Let  $X : G \rightarrow V$  and  $Y : G \rightarrow W$  be irreducible representations of a group  $G$ . If there exists a fixed linear transformation  $T : V \rightarrow W$  such that  $TX(g) = Y(g)T$  for all  $g \in G$ , then  $T$  is either the zero map or invertible.*

## Proof (sketch)

1. The kernel of  $T$  is invariant under  $X(G)$ .
2. The image of  $T$  is invariant under  $Y(G)$ .
3. Since  $X$  and  $Y$  are irreducible,  $\ker(T) = \{\mathbf{0}\}$  and  $\text{im}(T) = V$  or  $\ker(T) = V$  and  $\text{im}(T) = \{0\}$ .

# Schur's Lemmas (pt. 1)

## Lemma

*Let  $X : G \rightarrow V$  and  $Y : G \rightarrow W$  be irreducible representations of a group  $G$ . If there exists a fixed linear transformation  $T : V \rightarrow W$  such that  $TX(g) = Y(g)T$  for all  $g \in G$ , then  $T$  is either the zero map or invertible.*

## Proof (sketch)

1. The kernel of  $T$  is invariant under  $X(G)$ .
2. The image of  $T$  is invariant under  $Y(G)$ .
3. Since  $X$  and  $Y$  are irreducible,  $\ker(T) = \{\mathbf{0}\}$  and  $\text{im}(T) = V$  or  $\ker(T) = V$  and  $\text{im}(T) = \{0\}$ .
4. By the rank-nullity theorem, conclude that  $T$  is either the zero map or invertible.

# Schur's Lemma's (pt. 2)

## Lemma

*Let  $X$  be an irreducible representation of a group  $G$  and  $T$  a linear operator that commutes with all  $X(g)$  for  $g \in G$ . Then  $T$  is a scalar multiple of the identity operator.*

# Schur's Lemma's (pt. 2)

## Lemma

*Let  $X$  be an irreducible representation of a group  $G$  and  $T$  a linear operator that commutes with all  $X(g)$  for  $g \in G$ . Then  $T$  is a scalar multiple of the identity operator.*

## Proof (sketch)

# Schur's Lemma's (pt. 2)

## Lemma

*Let  $X$  be an irreducible representation of a group  $G$  and  $T$  a linear operator that commutes with all  $X(g)$  for  $g \in G$ . Then  $T$  is a scalar multiple of the identity operator.*

## Proof (sketch)

1. Consider  $\lambda$  to be an eigenvalue of  $T$ .

# Schur's Lemma's (pt. 2)

## Lemma

*Let  $X$  be an irreducible representation of a group  $G$  and  $T$  a linear operator that commutes with all  $X(g)$  for  $g \in G$ . Then  $T$  is a scalar multiple of the identity operator.*

## Proof (sketch)

1. Consider  $\lambda$  to be an eigenvalue of  $T$ .
2. Then  $T - \lambda I$  is not invertible.

# Schur's Lemma's (pt. 2)

## Lemma

*Let  $X$  be an irreducible representation of a group  $G$  and  $T$  a linear operator that commutes with all  $X(g)$  for  $g \in G$ . Then  $T$  is a scalar multiple of the identity operator.*

## Proof (sketch)

1. Consider  $\lambda$  to be an eigenvalue of  $T$ .
2. Then  $T - \lambda I$  is not invertible.
3. By assumption,  $(T - \lambda I)X(g) = X(g)(T - \lambda I)$  for all  $g \in G$ .

# Schur's Lemma's (pt. 2)

## Lemma

*Let  $X$  be an irreducible representation of a group  $G$  and  $T$  a linear operator that commutes with all  $X(g)$  for  $g \in G$ . Then  $T$  is a scalar multiple of the identity operator.*

## Proof (sketch)

1. Consider  $\lambda$  to be an eigenvalue of  $T$ .
2. Then  $T - \lambda I$  is not invertible.
3. By assumption,  $(T - \lambda I)X(g) = X(g)(T - \lambda I)$  for all  $g \in G$ .
4. By previous lemma,  $T - \lambda I = 0 \implies T = \lambda I$ .



# Consequence of Schur's Lemmas

## Corollary

*If  $G$  is a finite abelian group, then the irreducible representations of  $G$  are one-dimensional.*

# Consequence of Schur's Lemmas

## Corollary

*If  $G$  is a finite abelian group, then the irreducible representations of  $G$  are one-dimensional.*

## Proof (sketch)

# Consequence of Schur's Lemmas

## Corollary

*If  $G$  is a finite abelian group, then the irreducible representations of  $G$  are one-dimensional.*

## Proof (sketch)

1. Fix  $h \in G$ .

# Consequence of Schur's Lemmas

## Corollary

*If  $G$  is a finite abelian group, then the irreducible representations of  $G$  are one-dimensional.*

## Proof (sketch)

1. Fix  $h \in G$ .
2. Since  $G$  is abelian,  $X(h)X(g) = X(g)X(h)$  for all  $g \in G$ .

# Consequence of Schur's Lemmas

## Corollary

*If  $G$  is a finite abelian group, then the irreducible representations of  $G$  are one-dimensional.*

## Proof (sketch)

1. Fix  $h \in G$ .
2. Since  $G$  is abelian,  $X(h)X(g) = X(g)X(h)$  for all  $g \in G$ .
3. Schur's second lemma implies  $X(h) = \lambda_h I$  for some scalar  $\lambda_h$ .

# Consequence of Schur's Lemmas

## Corollary

*If  $G$  is a finite abelian group, then the irreducible representations of  $G$  are one-dimensional.*

## Proof (sketch)

1. Fix  $h \in G$ .
2. Since  $G$  is abelian,  $X(h)X(g) = X(g)X(h)$  for all  $g \in G$ .
3. Schur's second lemma implies  $X(h) = \lambda_h I$  for some scalar  $\lambda_h$ .
4. The element  $h$  was arbitrary, so  $X(g) = \lambda_g I$  for all  $g \in G$ .

# Consequence of Schur's Lemmas

## Corollary

*If  $G$  is a finite abelian group, then the irreducible representations of  $G$  are one-dimensional.*

## Proof (sketch)

1. Fix  $h \in G$ .
2. Since  $G$  is abelian,  $X(h)X(g) = X(g)X(h)$  for all  $g \in G$ .
3. Schur's second lemma implies  $X(h) = \lambda_h I$  for some scalar  $\lambda_h$ .
4. The element  $h$  was arbitrary, so  $X(g) = \lambda_g I$  for all  $g \in G$ .
5.  $X(G)$  is equivalent to the representation  $g \mapsto \lambda_g$  for all  $g \in G$ .

# Consequence of Schur's Lemmas

## Corollary

*If  $G$  is a finite abelian group, then the irreducible representations of  $G$  are one-dimensional.*

## Proof (sketch)

1. Fix  $h \in G$ .
2. Since  $G$  is abelian,  $X(h)X(g) = X(g)X(h)$  for all  $g \in G$ .
3. Schur's second lemma implies  $X(h) = \lambda_h I$  for some scalar  $\lambda_h$ .
4. The element  $h$  was arbitrary, so  $X(g) = \lambda_g I$  for all  $g \in G$ .
5.  $X(G)$  is equivalent to the representation  $g \mapsto \lambda_g$  for all  $g \in G$ .
6. One-dimensional representations are irreducible.



# A Note About Irreducibility

# A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.

# A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:

# A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
  - ◇ Direct sums

# A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
  - ◇ Direct sums
  - ◇ Tensor products

# A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
  - ◇ Direct sums
  - ◇ Tensor products
  - ◇ Complex conjugation<sup>4</sup>

---

<sup>4</sup>If the representation matrices have entries in  $\mathbb{C}$ .

# A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
  - ◇ Direct sums
  - ◇ Tensor products
  - ◇ Complex conjugation<sup>4</sup>
  - ◇ Similarity transforms

---

<sup>4</sup>If the representation matrices have entries in  $\mathbb{C}$ .

# A Note About Irreducibility

- ▶ Irreducible representations are the building blocks of all representations.
- ▶ Irreducible representations can be combined/modified to create new representations, such as:
  - ◇ Direct sums
  - ◇ Tensor products
  - ◇ Complex conjugation<sup>4</sup>
  - ◇ Similarity transforms

## How does this help in physics?

Irreducible representations can describe symmetries of physical systems with remarkably fundamental implications.

---

<sup>4</sup>If the representation matrices have entries in  $\mathbb{C}$ .





**CAL POLY**

## **2 Examples in Physics**

# Preliminaries: Physics Conventions

# Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.

# Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.

# Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \qquad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

# Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \qquad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

# Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \qquad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

# Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \quad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

4. The *Hermitian conjugate* or *adjoint* of an operator  $A$  is denoted  $A^\dagger$ , and is thought of as complex conjugation and transposition in matrix form.



# Preliminaries: Physics Conventions

1. The quantum state of a system is described by a vector in a complex Hilbert space.
2. The corresponding vectors are often called *state vectors*.
3. The inner product defined on the Hilbert space is linear in the second argument:

$$(1) \quad \langle \phi, \lambda \psi \rangle = \lambda \langle \phi, \psi \rangle \quad (2) \quad \langle \alpha \phi, \psi \rangle = \bar{\alpha} \langle \phi, \psi \rangle$$

4. The *Hermitian conjugate* or *adjoint* of an operator  $A$  is denoted  $A^\dagger$ , and is thought of as complex conjugation and transposition in matrix form.
5. Operators that are self-adjoint are called *Hermitian*.

# Preliminaries: Dirac notation

## Preliminaries: Dirac notation

- *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.

# Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted  $|\psi\rangle$ .

# Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted  $|\psi\rangle$ .
- ▶ A *bra* is a row vector,  $\langle\psi|$ . This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

# Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted  $|\psi\rangle$ .
- ▶ A *bra* is a row vector,  $\langle\psi|$ . This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

- ▶ **Inner product:**  $\langle\phi|\psi\rangle$

# Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted  $|\psi\rangle$ .
- ▶ A *bra* is a row vector,  $\langle\psi|$ . This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

- ▶ **Inner product:**  $\langle\phi|\psi\rangle$
- ▶ **Outer product:**  $|\phi\rangle\langle\psi|$

# Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted  $|\psi\rangle$ .
- ▶ A *bra* is a row vector,  $\langle\psi|$ . This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

- ▶ **Inner product:**  $\langle\phi|\psi\rangle$
- ▶ **Outer product:**  $|\phi\rangle\langle\psi|$
- ▶ The action of an operator  $A$  on a vector  $|\psi\rangle$  is written as  $|A\psi\rangle = A|\psi\rangle$ .



# Preliminaries: Dirac notation

- ▶ *Dirac* or *bra-ket* notation is a convenient way to represent vectors and operators in quantum mechanics.
- ▶ A *ket* is a column (state) vector, denoted  $|\psi\rangle$ .
- ▶ A *bra* is a row vector,  $\langle\psi|$ . This can be thought of as a linear functional on the relevant Hilbert space:

$$\langle\phi|(\psi) = \langle\phi, \psi\rangle.$$

- ▶ **Inner product:**  $\langle\phi|\psi\rangle$
- ▶ **Outer product:**  $|\phi\rangle\langle\psi|$
- ▶ The action of an operator  $A$  on a vector  $|\psi\rangle$  is written as  $|A\psi\rangle = A|\psi\rangle$ .
- ▶ Equivalent ways to write the same thing:

$$\langle A^\dagger \phi | \psi \rangle = \langle \phi | A | \psi \rangle = \langle \phi | A \psi \rangle .$$

# Orthonormality, Completeness, and Wavefunctions

## Definition

Let  $\{|1\rangle, |2\rangle, |3\rangle, \dots\}$  be an orthonormal basis for some quantum Hilbert space. In the context of physics, the **orthonormality** and **completeness** relations of the basis vectors allow any state vector  $|\psi\rangle$  to be written as a linear combination of the basis vectors:

$$|\psi\rangle = \left( \sum_n |n\rangle \langle n| \right) |\psi\rangle = \sum_n |n\rangle \langle n|\psi\rangle,$$

where  $\sum_n |n\rangle \langle n|$  is the identity operator.

# Orthonormality, Completeness, and Wavefunctions

## Definition

Let  $\{|1\rangle, |2\rangle, |3\rangle, \dots\}$  be an orthonormal basis for some quantum Hilbert space. In the context of physics, the **orthonormality** and **completeness** relations of the basis vectors allow any state vector  $|\psi\rangle$  to be written as a linear combination of the basis vectors:

$$|\psi\rangle = \left( \sum_n |n\rangle \langle n| \right) |\psi\rangle = \sum_n |n\rangle \langle n|\psi\rangle,$$

where  $\sum_n |n\rangle \langle n|$  is the identity operator.

*This is just a fancy change of basis!*

# Orthonormality, Completeness, and Wavefunctions

## Definition

Let  $\{|1\rangle, |2\rangle, |3\rangle, \dots\}$  be an orthonormal basis for some quantum Hilbert space. In the context of physics, the **orthonormality** and **completeness** relations of the basis vectors allow any state vector  $|\psi\rangle$  to be written as a linear combination of the basis vectors:

$$|\psi\rangle = \left( \sum_n |n\rangle \langle n| \right) |\psi\rangle = \sum_n |n\rangle \langle n|\psi\rangle,$$

where  $\sum_n |n\rangle \langle n|$  is the identity operator.

*This is just a fancy change of basis!*

## Definition

For a continuous basis labelled by  $|x\rangle$  where  $x$  is a continuous parameter, the **wavefunction**  $\psi(x)$  is the projection:  $\langle x|\psi\rangle = \psi(x)$ .

# Properties of 2D Rotations

Let  $R$  denote the familiar rotation matrix representation from before.

# Properties of 2D Rotations

Let  $R$  denote the familiar rotation matrix representation from before.

## Definition

An *orthogonal matrix*  $O$  satisfies  $O^T = O^{-1}$ .

# Properties of 2D Rotations

Let  $R$  denote the familiar rotation matrix representation from before.

## Definition

An *orthogonal matrix*  $O$  satisfies  $O^\top = O^{-1}$ .

**Rotation matrices are orthogonal:**

$$R(\phi)R^\top(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

# Properties of 2D Rotations

Let  $R$  denote the familiar rotation matrix representation from before.

## Definition

An *orthogonal matrix*  $O$  satisfies  $O^\top = O^{-1}$ .

**Rotation matrices are orthogonal:**

$$R(\phi)R^\top(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

**Rotations preserve vector lengths:**

$$R(\phi)\mathbf{x} = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \phi - x_2 \sin \phi \\ x_1 \sin \phi + x_2 \cos \phi \end{bmatrix} \implies |R(\phi)\mathbf{x}|^2 = |\mathbf{x}|^2.$$



# Properties of 2D Rotations

Let  $R$  denote the familiar rotation matrix representation from before.

## Definition

An *orthogonal matrix*  $O$  satisfies  $O^\top = O^{-1}$ .

**Rotation matrices are orthogonal:**

$$R(\phi)R^\top(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

**Rotations preserve vector lengths:**

$$R(\phi)\mathbf{x} = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \phi - x_2 \sin \phi \\ x_1 \sin \phi + x_2 \cos \phi \end{bmatrix} \implies |R(\phi)\mathbf{x}|^2 = |\mathbf{x}|^2.$$

This *special* property is summarized by noting  $\det R(\phi) = 1$  for all  $\phi \in [0, 2\pi)$ .

# The $SO(2)$ Group

## Definition

The *special orthogonal group* in two dimensions, denoted  $SO(2)$ , is the group of all  $2 \times 2$  orthogonal matrices with determinant equal to  $+1$ .<sup>5</sup>

---

<sup>5</sup>For all intents and purposes,  $SO(2)$  is  $R$  from before.

# The $SO(2)$ Group

## Definition

The *special orthogonal group* in two dimensions, denoted  $SO(2)$ , is the group of all  $2 \times 2$  orthogonal matrices with determinant equal to  $+1$ .<sup>5</sup>

## Properties of $SO(2)$ :

---

<sup>5</sup>For all intents and purposes,  $SO(2)$  is  $R$  from before.

# The $SO(2)$ Group

## Definition

The *special orthogonal group* in two dimensions, denoted  $SO(2)$ , is the group of all  $2 \times 2$  orthogonal matrices with determinant equal to  $+1$ .<sup>5</sup>

## Properties of $SO(2)$ :

- ▶ The *periodicity condition*  $R(\phi + 2\pi) = R(\phi)$  is satisfied.

---

<sup>5</sup>For all intents and purposes,  $SO(2)$  is  $R$  from before.

# The $SO(2)$ Group

## Definition

The *special orthogonal group* in two dimensions, denoted  $SO(2)$ , is the group of all  $2 \times 2$  orthogonal matrices with determinant equal to  $+1$ .<sup>5</sup>

## Properties of $SO(2)$ :

- ▶ The *periodicity condition*  $R(\phi + 2\pi) = R(\phi)$  is satisfied.
- ▶ The *identity element* is  $R(0) = I$ .

---

<sup>5</sup>For all intents and purposes,  $SO(2)$  is  $R$  from before.

# The $SO(2)$ Group

## Definition

The *special orthogonal group* in two dimensions, denoted  $SO(2)$ , is the group of all  $2 \times 2$  orthogonal matrices with determinant equal to  $+1$ .<sup>5</sup>

## Properties of $SO(2)$ :

- ▶ The *periodicity condition*  $R(\phi + 2\pi) = R(\phi)$  is satisfied.
- ▶ The *identity element* is  $R(0) = I$ .
- ▶  $SO(2)$  is *abelian*:  $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$ .

---

<sup>5</sup>For all intents and purposes,  $SO(2)$  is  $R$  from before.

# The $SO(2)$ Group

## Definition

The *special orthogonal group* in two dimensions, denoted  $SO(2)$ , is the group of all  $2 \times 2$  orthogonal matrices with determinant equal to  $+1$ .<sup>5</sup>

## Properties of $SO(2)$ :

- ▶ The *periodicity condition*  $R(\phi + 2\pi) = R(\phi)$  is satisfied.
- ▶ The *identity element* is  $R(0) = I$ .
- ▶  $SO(2)$  is *abelian*:  $R(\phi_1)R(\phi_2) = R(\phi_1 + \phi_2) = R(\phi_2)R(\phi_1)$ .
- ▶  $SO(2)$  is *reducible* (earlier example with  $\mathbf{e}_{\pm}$ ).

---

<sup>5</sup>For all intents and purposes,  $SO(2)$  is  $R$  from before.

# Infinitesimal Rotations

- Consider an *infinitesimal rotation* labelled by some infinitesimal angle  $d\phi$ .



# Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle  $d\phi$ .
- ▶ This is equivalent to the identity plus some small rotation, which can be written as<sup>6</sup>

$$R(d\phi) = I - i d\phi J$$

---

<sup>6</sup>The constant  $-i$  is introduced for later convenience, and  $J$  is a quantity independent of  $\phi$ .

# Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle  $d\phi$ .
- ▶ This is equivalent to the identity plus some small rotation, which can be written as<sup>6</sup>

$$R(d\phi) = I - i d\phi J$$

- ▶ There are two ways to interpret  $R(\phi + d\phi)$ :

$$R(\phi + d\phi) = R(\phi)R(d\phi) = R(\phi)(I - id\phi J) = R(\phi) - id\phi R(\phi)J,$$

$$R(\phi + d\phi) = R(\phi) + d\phi \frac{dR(\phi)}{d\phi}$$

---

<sup>6</sup>The constant  $-i$  is introduced for later convenience, and  $J$  is a quantity independent of  $\phi$ .

# Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle  $d\phi$ .
- ▶ This is equivalent to the identity plus some small rotation, which can be written as<sup>6</sup>

$$R(d\phi) = I - i d\phi J$$

- ▶ There are two ways to interpret  $R(\phi + d\phi)$ :

$$R(\phi + d\phi) = R(\phi)R(d\phi) = R(\phi)(I - id\phi J) = R(\phi) - id\phi R(\phi)J,$$

$$R(\phi + d\phi) = R(\phi) + d\phi \frac{dR(\phi)}{d\phi}$$

- ▶ Equating the two expressions gives the differential equation  $dR(\phi) = -id\phi R(\phi)J$ .

---

<sup>6</sup>The constant  $-i$  is introduced for later convenience, and  $J$  is a quantity independent of  $\phi$ .

# Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle  $d\phi$ .
- ▶ This is equivalent to the identity plus some small rotation, which can be written as<sup>6</sup>

$$R(d\phi) = I - i d\phi J$$

- ▶ There are two ways to interpret  $R(\phi + d\phi)$ :

$$R(\phi + d\phi) = R(\phi)R(d\phi) = R(\phi)(I - id\phi J) = R(\phi) - id\phi R(\phi)J,$$

$$R(\phi + d\phi) = R(\phi) + d\phi \frac{dR(\phi)}{d\phi}$$

- ▶ Equating the two expressions gives the differential equation  $dR(\phi) = -id\phi R(\phi)J$ .
- ▶ With  $R(0) = I$  boundary condition:  $R(\phi) = e^{-i\phi J}$ .

<sup>6</sup>The constant  $-i$  is introduced for later convenience, and  $J$  is a quantity independent of  $\phi$ .

# Infinitesimal Rotations

- ▶ Consider an *infinitesimal rotation* labelled by some infinitesimal angle  $d\phi$ .
- ▶ This is equivalent to the identity plus some small rotation, which can be written as<sup>6</sup>

$$R(d\phi) = I - i d\phi J$$

- ▶ There are two ways to interpret  $R(\phi + d\phi)$ :

$$R(\phi + d\phi) = R(\phi)R(d\phi) = R(\phi)(I - id\phi J) = R(\phi) - id\phi R(\phi)J,$$

$$R(\phi + d\phi) = R(\phi) + d\phi \frac{dR(\phi)}{d\phi}$$

- ▶ Equating the two expressions gives the differential equation  $dR(\phi) = -id\phi R(\phi)J$ .
- ▶ With  $R(0) = I$  boundary condition:  $R(\phi) = e^{-i\phi J}$ .
- ▶ We call  $J$  the *generator* of SO(2) rotations.

<sup>6</sup>The constant  $-i$  is introduced for later convenience, and  $J$  is a quantity independent of  $\phi$ .

# Recovering the Rotation Matrix from $J$

To first order in  $d\phi$ :  $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

# Recovering the Rotation Matrix from $J$

To first order in  $d\phi$ :  $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before:  $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

# Recovering the Rotation Matrix from $J$

To first order in  $d\phi$ :  $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before:  $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

Taylor expand:

$$R(\phi) = e^{-iJ\phi} = I - iJ\phi - I\frac{\phi^2}{2!} + iJ\frac{\phi^3}{3!} + \dots$$



# Recovering the Rotation Matrix from $J$

To first order in  $d\phi$ :  $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before:  $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

Taylor expand:

$$\begin{aligned} R(\phi) &= e^{-iJ\phi} = I - iJ\phi - I\frac{\phi^2}{2!} + iJ\frac{\phi^3}{3!} + \dots \\ &= I \left( \sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n}}{(2n)!} \right) - iJ \left( \sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n+1}}{(2n+1)!} \right) \end{aligned}$$

# Recovering the Rotation Matrix from $J$

To first order in  $d\phi$ :  $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before:  $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

Taylor expand:

$$\begin{aligned} R(\phi) &= e^{-iJ\phi} = I - iJ\phi - I\frac{\phi^2}{2!} + iJ\frac{\phi^3}{3!} + \dots \\ &= I \left( \sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n}}{(2n)!} \right) - iJ \left( \sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n+1}}{(2n+1)!} \right) \\ &= I \cos \phi - iJ \sin \phi \end{aligned}$$

# Recovering the Rotation Matrix from $J$

To first order in  $d\phi$ :  $R(d\phi) = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix}$

From before:  $I - id\phi J = \begin{bmatrix} 1 & -d\phi \\ d\phi & 1 \end{bmatrix} \implies J = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \implies J^2 = I$

Taylor expand:

$$\begin{aligned} R(\phi) &= e^{-iJ\phi} = I - iJ\phi - I\frac{\phi^2}{2!} + iJ\frac{\phi^3}{3!} + \dots \\ &= I \left( \sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n}}{(2n)!} \right) - iJ \left( \sum_{n=0}^{\infty} (-1)^n \frac{\phi^{2n+1}}{(2n+1)!} \right) \\ &= I \cos \phi - iJ \sin \phi \\ &= \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix}. \end{aligned}$$

# Irreducible Representations of $SO(2)$

**Process to obtaining irreducibles:**

# Irreducible Representations of $SO(2)$

**Process to obtaining irreducibles:**

1. Let  $U$  be any representation of  $SO(2)$ .

# Irreducible Representations of $SO(2)$

## Process to obtaining irreducibles:

1. Let  $U$  be any representation of  $SO(2)$ .
2. Same argument as before:  $U(\phi) = e^{-iJ\phi}$ , where  $J$  is not necessarily the same as before.

# Irreducible Representations of $SO(2)$

## Process to obtaining irreducibles:

1. Let  $U$  be any representation of  $SO(2)$ .
2. Same argument as before:  $U(\phi) = e^{-iJ\phi}$ , where  $J$  is not necessarily the same as before.
3.  $SO(2)$  is abelian: Schur's Lemmas  $\implies$  all irreducible representations are 1D.

# Irreducible Representations of $SO(2)$

## Process to obtaining irreducibles:

1. Let  $U$  be any representation of  $SO(2)$ .
2. Same argument as before:  $U(\phi) = e^{-iJ\phi}$ , where  $J$  is not necessarily the same as before.
3.  $SO(2)$  is abelian: Schur's Lemmas  $\implies$  all irreducible representations are 1D.
4. Each invariant subspace is spanned by an eigenvector of  $J$ :

$$J|m\rangle = m|m\rangle ,$$
$$U(\phi)|m\rangle = e^{-iJ\phi}|m\rangle = e^{-im\phi}|m\rangle .$$



# Irreducible Representations of $SO(2)$

## Process to obtaining irreducibles:

1. Let  $U$  be any representation of  $SO(2)$ .
2. Same argument as before:  $U(\phi) = e^{-iJ\phi}$ , where  $J$  is not necessarily the same as before.
3.  $SO(2)$  is abelian: Schur's Lemmas  $\implies$  all irreducible representations are 1D.
4. Each invariant subspace is spanned by an eigenvector of  $J$ :

$$J|m\rangle = m|m\rangle,$$
$$U(\phi)|m\rangle = e^{-iJ\phi}|m\rangle = e^{-im\phi}|m\rangle.$$

5. Periodicity of  $SO(2) \implies e^{-i2\pi m} = 1 \implies m \in \mathbb{Z}$ .

# Irreducible Representations of $SO(2)$

## Process to obtaining irreducibles:

1. Let  $U$  be any representation of  $SO(2)$ .
2. Same argument as before:  $U(\phi) = e^{-iJ\phi}$ , where  $J$  is not necessarily the same as before.
3.  $SO(2)$  is abelian: Schur's Lemmas  $\implies$  all irreducible representations are 1D.
4. Each invariant subspace is spanned by an eigenvector of  $J$ :

$$J|m\rangle = m|m\rangle,$$
$$U(\phi)|m\rangle = e^{-iJ\phi}|m\rangle = e^{-im\phi}|m\rangle.$$

5. Periodicity of  $SO(2) \implies e^{-i2\pi m} = 1 \implies m \in \mathbb{Z}$ .

## Theorem

*The single-valued irreducible representations of  $SO(2)$  are defined as*

$$U^m(\phi) = e^{-im\phi}, \quad \forall m \in \mathbb{Z}.$$

# Generalization to 3 Spatial Dimensions

# Generalization to 3 Spatial Dimensions

- In 3 spatial dimensions, every rotation can be thought of as a rotation in a plane with some perpendicular axis of rotation  $\mathbf{n}$ :  $R_{\mathbf{n}}(\theta)$ .

# Generalization to 3 Spatial Dimensions

- ▶ In 3 spatial dimensions, every rotation can be thought of as a rotation in a plane with some perpendicular axis of rotation  $\mathbf{n}$ :  $R_{\mathbf{n}}(\theta)$ .
- ▶ Rotations in a plane are isomorphic to  $\text{SO}(2)$ :  $R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}}$  for some generator  $J_{\mathbf{n}}$ .

# Generalization to 3 Spatial Dimensions

- ▶ In 3 spatial dimensions, every rotation can be thought of as a rotation in a plane with some perpendicular axis of rotation  $\mathbf{n}$ :  $R_{\mathbf{n}}(\theta)$ .
- ▶ Rotations in a plane are isomorphic to  $\text{SO}(2)$ :  $R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}}$  for some generator  $J_{\mathbf{n}}$ .
- ▶ The standard generators along each axis  $\{J_x, J_y, J_z\}$  form a basis for all rotation generators:  $J_{\mathbf{n}} = n_x J_x + n_y J_y + n_z J_z$ .

# Generalization to 3 Spatial Dimensions

- ▶ In 3 spatial dimensions, every rotation can be thought of as a rotation in a plane with some perpendicular axis of rotation  $\mathbf{n}$ :  $R_{\mathbf{n}}(\theta)$ .
- ▶ Rotations in a plane are isomorphic to  $SO(2)$ :  $R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}}$  for some generator  $J_{\mathbf{n}}$ .
- ▶ The standard generators along each axis  $\{J_x, J_y, J_z\}$  form a basis for all rotation generators:  $J_{\mathbf{n}} = n_x J_x + n_y J_y + n_z J_z$ .

**Consequence:** Any rotation in Euclidean 3-space can be written in terms of the generators:

$$R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}} = e^{-i\theta(n_x J_x + n_y J_y + n_z J_z)} = e^{-i\theta \mathbf{n} \cdot \mathbf{J}}.$$

# Generalization to 3 Spatial Dimensions

- ▶ In 3 spatial dimensions, every rotation can be thought of as a rotation in a plane with some perpendicular axis of rotation  $\mathbf{n}$ :  $R_{\mathbf{n}}(\theta)$ .
- ▶ Rotations in a plane are isomorphic to  $\text{SO}(2)$ :  $R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}}$  for some generator  $J_{\mathbf{n}}$ .
- ▶ The standard generators along each axis  $\{J_x, J_y, J_z\}$  form a basis for all rotation generators:  $J_{\mathbf{n}} = n_x J_x + n_y J_y + n_z J_z$ .

**Consequence:** Any rotation in Euclidean 3-space can be written in terms of the generators:

$$R_{\mathbf{n}}(\theta) = e^{-i\theta J_{\mathbf{n}}} = e^{-i\theta(n_x J_x + n_y J_y + n_z J_z)} = e^{-i\theta \mathbf{n} \cdot \mathbf{J}}.$$

## Definition

The *special orthogonal group* in three dimensions, denoted  $\text{SO}(3)$ , is the group of all  $3 \times 3$  orthogonal matrices with determinant equal to  $+1$ .  $\text{SO}(3)$  rotations are generated by the components of the Hermitian generator  $\mathbf{J} = [J_x, J_y, J_z]^T$ .



# Connection to Quantum Mechanics

Using a similar process to generate  $SO(3)$  invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

# Connection to Quantum Mechanics

Using a similar process to generate  $SO(3)$  invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

## Theorem

*The irreducible representations of  $SO(3)$  are labeled by  $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$ , and the  $2j + 1$  eigenvectors spanning an invariant subspace are labelled by their eigenvalues:  
 $m = -j, -j + 1, \dots, j - 1, j$ .*

# Connection to Quantum Mechanics

Using a similar process to generate  $SO(3)$  invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

## Theorem

*The irreducible representations of  $SO(3)$  are labeled by  $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$ , and the  $2j + 1$  eigenvectors spanning an invariant subspace are labelled by their eigenvalues:  
 $m = -j, -j + 1, \dots, j - 1, j$ .*

**Consequences:**

# Connection to Quantum Mechanics

Using a similar process to generate  $SO(3)$  invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

## Theorem

*The irreducible representations of  $SO(3)$  are labeled by  $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$ , and the  $2j + 1$  eigenvectors spanning an invariant subspace are labelled by their eigenvalues:  
 $m = -j, -j + 1, \dots, j - 1, j$ .*

## Consequences:

- One can obtain the explicit form of  $\mathbf{J}$  and subsequently its components  $J_x, J_y, J_z$ . These are precisely the angular momentum operators in quantum mechanics.

# Connection to Quantum Mechanics

Using a similar process to generate  $SO(3)$  invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

## Theorem

*The irreducible representations of  $SO(3)$  are labeled by  $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$ , and the  $2j + 1$  eigenvectors spanning an invariant subspace are labelled by their eigenvalues:  $m = -j, -j + 1, \dots, j - 1, j$ .*

## Consequences:

- ▶ One can obtain the explicit form of  $\mathbf{J}$  and subsequently its components  $J_x, J_y, J_z$ . These are precisely the angular momentum operators in quantum mechanics.
- ▶ The eigenvalues of  $J^2 = \mathbf{J} \cdot \mathbf{J}$  and  $J_z$  are  $j(j + 1)$  and  $m$ , respectively<sup>7</sup>. In quantum physics, these eigenvalues correspond to the observable total angular momentum and its z-component.

---

<sup>7</sup>Typically, the z-axis is chosen as the standard axis.

# Connection to Quantum Mechanics

Using a similar process to generate  $SO(3)$  invariant subspaces that correspond to irreducible representations, we summarize the results in a theorem:

## Theorem

*The irreducible representations of  $SO(3)$  are labeled by  $j = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$ , and the  $2j + 1$  eigenvectors spanning an invariant subspace are labelled by their eigenvalues:  
 $m = -j, -j + 1, \dots, j - 1, j$ .*

## Consequences:

- ▶ One can obtain the explicit form of  $\mathbf{J}$  and subsequently its components  $J_x, J_y, J_z$ . These are precisely the angular momentum operators in quantum mechanics.
- ▶ The eigenvalues of  $J^2 = \mathbf{J} \cdot \mathbf{J}$  and  $J_z$  are  $j(j + 1)$  and  $m$ , respectively<sup>7</sup>. In quantum physics, these eigenvalues correspond to the observable total angular momentum and its z-component.
- ▶ This generalizes to other types of angular momentum, such as *spin angular momentum*!

---

<sup>7</sup>Typically, the z-axis is chosen as the standard axis.

# Connection to Quantum Mechanics: Punchline

# Connection to Quantum Mechanics: Punchline

## Discretization of Angular Momentum for Free

Arguably the most defining characteristic of quantum mechanics is that classically measurable quantities become discretized (quantized) when observed on the quantum scale. Without any physical motivation, the irreducible representations of  $SO(3)$  gave it to us for free!



# Connection to Quantum Mechanics: Punchline

## Discretization of Angular Momentum for Free

Arguably the most defining characteristic of quantum mechanics is that classically measurable quantities become discretized (quantized) when observed on the quantum scale. Without any physical motivation, the irreducible representations of  $SO(3)$  gave it to us for free!

But that's not all folks!

# Conservation of Angular Momentum

1. The *commutator* of two operators  $A$  and  $B$  is defined as  $[A, B] = AB - BA$ .

# Conservation of Angular Momentum

1. The *commutator* of two operators  $A$  and  $B$  is defined as  $[A, B] = AB - BA$ .
2. The *Hamiltonian* operator  $\hat{H}$  is the quantum mechanical operator corresponding to the total energy of a system.

# Conservation of Angular Momentum

1. The *commutator* of two operators  $A$  and  $B$  is defined as  $[A, B] = AB - BA$ .
2. The *Hamiltonian* operator  $\hat{H}$  is the quantum mechanical operator corresponding to the total energy of a system.

## Theorem (Ehrenfest)

*If a time-independent Hermitian operator commutes with the Hamiltonian, then the physical observable corresponding to the operator is conserved.*

# Conservation of Angular Momentum

1. The *commutator* of two operators  $A$  and  $B$  is defined as  $[A, B] = AB - BA$ .
2. The *Hamiltonian* operator  $\hat{H}$  is the quantum mechanical operator corresponding to the total energy of a system.

## Theorem (Ehrenfest)

*If a time-independent Hermitian operator commutes with the Hamiltonian, then the physical observable corresponding to the operator is conserved.*

## Consequences

1. Any system with radial symmetry is invariant under  $SO(3)$  rotations, so  $[\hat{H}, \mathbf{J}] = 0$ .

# Conservation of Angular Momentum

1. The *commutator* of two operators  $A$  and  $B$  is defined as  $[A, B] = AB - BA$ .
2. The *Hamiltonian* operator  $\hat{H}$  is the quantum mechanical operator corresponding to the total energy of a system.

## Theorem (Ehrenfest)

*If a time-independent Hermitian operator commutes with the Hamiltonian, then the physical observable corresponding to the operator is conserved.*

## Consequences

1. Any system with radial symmetry is invariant under  $SO(3)$  rotations, so  $[\hat{H}, \mathbf{J}] = 0$ .
2. Conservation of angular momentum is a direct result of the radial symmetry of the system.

# Conservation of Angular Momentum

1. The *commutator* of two operators  $A$  and  $B$  is defined as  $[A, B] = AB - BA$ .
2. The *Hamiltonian* operator  $\hat{H}$  is the quantum mechanical operator corresponding to the total energy of a system.

## Theorem (Ehrenfest)

*If a time-independent Hermitian operator commutes with the Hamiltonian, then the physical observable corresponding to the operator is conserved.*

## Consequences

1. Any system with radial symmetry is invariant under  $SO(3)$  rotations, so  $[\hat{H}, \mathbf{J}] = 0$ .
2. Conservation of angular momentum is a direct result of the radial symmetry of the system.
3. Similar arguments can be made for the continuous group of translations in space, leading to the conservation of linear momentum for translationally invariant systems.

# Additional Applications



# Additional Applications

1. The  $j = 1/2$  irreducible representation of  $SO(3)$  describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!

## Additional Applications

1. The  $j = 1/2$  irreducible representation of  $SO(3)$  describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of  $SO(3)$  to obtain multi-particle states, arriving at results such as:

## Additional Applications

1. The  $j = 1/2$  irreducible representation of  $SO(3)$  describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of  $SO(3)$  to obtain multi-particle states, arriving at results such as:
  - ◇ Clebsch-Gordan coefficients

# Additional Applications

1. The  $j = 1/2$  irreducible representation of  $SO(3)$  describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of  $SO(3)$  to obtain multi-particle states, arriving at results such as:
  - ◇ Clebsch-Gordan coefficients
  - ◇ singlet versus triplet states

# Additional Applications

1. The  $j = 1/2$  irreducible representation of  $SO(3)$  describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of  $SO(3)$  to obtain multi-particle states, arriving at results such as:
  - ◇ Clebsch-Gordan coefficients
  - ◇ singlet versus triplet states
  - ◇ the Pauli exclusion principle

# Additional Applications

1. The  $j = 1/2$  irreducible representation of  $SO(3)$  describes *fermions*. A modified periodicity condition due to the half-integer representation leads to *spinors*!
2. We can take tensor products of the irreducibles of  $SO(3)$  to obtain multi-particle states, arriving at results such as:
  - ◇ Clebsch-Gordan coefficients
  - ◇ singlet versus triplet states
  - ◇ the Pauli exclusion principle

*This is the tip of the iceberg!*



**CAL POLY**

### **3 The Braid Group**

# The Braid Group



# The Braid Group

## Definition

The *configuration space* of  $n$  ordered distinct points in the complex plane  $\mathbb{C}$  is defined as  $M_n = \{(z_1, \dots, z_n) \in \mathbb{C}^n; z_i \neq z_j, \forall i \neq j\}$ .

# The Braid Group

## Definition

The *configuration space* of  $n$  ordered distinct points in the complex plane  $\mathbb{C}$  is defined as  $M_n = \{(z_1, \dots, z_n) \in \mathbb{C}^n; z_i \neq z_j, \forall i \neq j\}$ .

- Note that  $(z_1, z_2, z_3, \dots, z_n)$  and  $(z_2, z_1, z_3, \dots, z_n)$  are distinct configurations in  $M_n$ .

# The Braid Group

## Definition

The *configuration space* of  $n$  ordered distinct points in the complex plane  $\mathbb{C}$  is defined as  $M_n = \{(z_1, \dots, z_n) \in \mathbb{C}^n; z_i \neq z_j, \forall i \neq j\}$ .

- ▶ Note that  $(z_1, z_2, z_3, \dots, z_n)$  and  $(z_2, z_1, z_3, \dots, z_n)$  are distinct configurations in  $M_n$ .
- ▶ A *braid*  $\beta$  is a *loop*<sup>8</sup> in  $M_n$  and can be thought of as a configuration that evolves over time:

$$\begin{aligned}\beta &: [0, 1] \rightarrow M_n \\ t &\mapsto \beta(t) = (\beta_1(t), \beta_2(t), \dots, \beta_n(t)),\end{aligned}$$

---

<sup>8</sup>The topological formalisms that define the braid group are omitted for times sake.

# The Braid Group

## Definition

The *configuration space* of  $n$  ordered distinct points in the complex plane  $\mathbb{C}$  is defined as  $M_n = \{(z_1, \dots, z_n) \in \mathbb{C}^n; z_i \neq z_j, \forall i \neq j\}$ .

- ▶ Note that  $(z_1, z_2, z_3, \dots, z_n)$  and  $(z_2, z_1, z_3, \dots, z_n)$  are distinct configurations in  $M_n$ .
- ▶ A *braid*  $\beta$  is a *loop*<sup>8</sup> in  $M_n$  and can be thought of as a configuration that evolves over time:

$$\begin{aligned}\beta &: [0, 1] \rightarrow M_n \\ t &\mapsto \beta(t) = (\beta_1(t), \beta_2(t), \dots, \beta_n(t)),\end{aligned}$$

## Definition

The *braid group*  $B_n$  is the (fundamental) group of all complex-valued  $n$ -tuples  $(M_n)$  up to *homotopy*.

---

<sup>8</sup>The topological formalisms that define the braid group are omitted for times sake.

# Visualization of Braids

# Visualization of Braids

- Each path traced out by a point in the configuration space is a *strand*.

# Visualization of Braids

- ▶ Each path traced out by a point in the configuration space is a *strand*.
- ▶ The number of strands of a braid is equal to the number of points in each tuple in the configuration space.

# Visualization of Braids

- ▶ Each path traced out by a point in the configuration space is a *strand*.
- ▶ The number of strands of a braid is equal to the number of points in each tuple in the configuration space.
- ▶ We can think of a braid on  $n$  strands as the motion of  $n$  distinct points in the complex plane over a normalized time interval.



# Visualization of Braids

- ▶ Each path traced out by a point in the configuration space is a *strand*.
- ▶ The number of strands of a braid is equal to the number of points in each tuple in the configuration space.
- ▶ We can think of a braid on  $n$  strands as the motion of  $n$  distinct points in the complex plane over a normalized time interval.
- ▶ Each trajectory is a strand, and the braid is the collection of all strands.

# Visualization of Braids


- ▶ Each path traced out by a point in the configuration space is a *strand*.
- ▶ The number of strands of a braid is equal to the number of points in each tuple in the configuration space.
- ▶ We can think of a braid on  $n$  strands as the motion of  $n$  distinct points in the complex plane over a normalized time interval.
- ▶ Each trajectory is a strand, and the braid is the collection of all strands.
- ▶ A braid is defined up to *homotopy*.

# Visualization of Braids

- ▶ Each path traced out by a point in the configuration space is a *strand*.
- ▶ The number of strands of a braid is equal to the number of points in each tuple in the configuration space.
- ▶ We can think of a braid on  $n$  strands as the motion of  $n$  distinct points in the complex plane over a normalized time interval.
- ▶ Each trajectory is a strand, and the braid is the collection of all strands.
- ▶ A braid is defined up to *homotopy*.
- ▶ Visualized in  $\mathbb{C} \times [0, 1]$ .

# Visualization of Braids

- ▶ Each path traced out by a point in the configuration space is a *strand*.
- ▶ The number of strands of a braid is equal to the number of points in each tuple in the configuration space.
- ▶ We can think of a braid on  $n$  strands as the motion of  $n$  distinct points in the complex plane over a normalized time interval.
- ▶ Each trajectory is a strand, and the braid is the collection of all strands.
- ▶ A braid is defined up to *homotopy*.
- ▶ Visualized in  $\mathbb{C} \times [0, 1]$ .



# Standard Generators

# Standard Generators



- Every braid can be decomposed into a finite product of *standard generators* that permute adjacent points.

# Standard Generators

- ▶ Every braid can be decomposed into a finite product of *standard generators* that permute adjacent points.
- ▶ The standard generators of  $B_n$  are defined as  $\{\sigma_1, \sigma_2, \dots, \sigma_{n-1}\}$ , in which:

# Standard Generators



- ▶ Every braid can be decomposed into a finite product of *standard generators* that permute adjacent points.
- ▶ The standard generators of  $B_n$  are defined as  $\{\sigma_1, \sigma_2, \dots, \sigma_{n-1}\}$ , in which:





# Standard Generators

- ▶ Every braid can be decomposed into a finite product of *standard generators* that permute adjacent points.
- ▶ The standard generators of  $B_n$  are defined as  $\{\sigma_1, \sigma_2, \dots, \sigma_{n-1}\}$ , in which:



- ▶ The *degree* of a braid  $\beta \in B_n$  is the sum of the powers of the standard generators in the decomposition of  $\beta$ .

# Alternative Description of $B_n$

# Alternative Description of $B_n$

## Definition

The braid group on  $n$  strands, denoted  $B_n$ , is generated by the standard generators that follow the *braid relations*, summarized below:

$$B_n = \left\langle \sigma_1, \dots, \sigma_{n-1} \left| \begin{array}{l} \sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i - j| > 1 \\ \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \end{array} \right. \right\rangle.$$


# Alternative Description of $B_n$

## Definition

The braid group on  $n$  strands, denoted  $B_n$ , is generated by the standard generators that follow the *braid relations*, summarized below:

$$B_n = \left\langle \sigma_1, \dots, \sigma_{n-1} \left| \begin{array}{l} \sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i - j| > 1 \\ \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \end{array} \right. \right\rangle.$$

**Comment:**  $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$  is known as the *Yang-Baxter equation*, visualized below:



# One-Dimensional Representations of the Braid Group

# One-Dimensional Representations of the Braid Group

For  $\theta \in \mathbb{R}$  and  $j = 1, 2, \dots, n-1$ , we define some *one-dimensional representations* of  $B_n$ :

$$\begin{aligned} p_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

# One-Dimensional Representations of the Braid Group

For  $\theta \in \mathbb{R}$  and  $j = 1, 2, \dots, n-1$ , we define some *one-dimensional representations* of  $B_n$ :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*

# One-Dimensional Representations of the Braid Group

For  $\theta \in \mathbb{R}$  and  $j = 1, 2, \dots, n-1$ , we define some *one-dimensional representations* of  $B_n$ :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) = \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2)$$



# One-Dimensional Representations of the Braid Group

For  $\theta \in \mathbb{R}$  and  $j = 1, 2, \dots, n-1$ , we define some *one-dimensional representations* of  $B_n$ :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} \rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \end{aligned}$$

# One-Dimensional Representations of the Braid Group

For  $\theta \in \mathbb{R}$  and  $j = 1, 2, \dots, n-1$ , we define some *one-dimensional representations* of  $B_n$ :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} \rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \\ &= e^{i(\theta_1 - \theta_1 + \theta_2 + \theta_2)} \end{aligned}$$

# One-Dimensional Representations of the Braid Group

For  $\theta \in \mathbb{R}$  and  $j = 1, 2, \dots, n-1$ , we define some *one-dimensional representations* of  $B_n$ :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} \rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \\ &= e^{i(\theta_1 - \theta_1 + \theta_2 + \theta_2)} \\ &= e^{i \cdot 2\theta_2} = \rho_\theta(\sigma_2^2) \end{aligned}$$

# One-Dimensional Representations of the Braid Group

For  $\theta \in \mathbb{R}$  and  $j = 1, 2, \dots, n-1$ , we define some *one-dimensional representations* of  $B_n$ :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} \rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \\ &= e^{i(\theta_1 - \theta_1 + \theta_2 + \theta_2)} \\ &= e^{i \cdot 2\theta_2} = \rho_\theta(\sigma_2^2) \end{aligned}$$

Hence, for any  $\beta \in B_n$  with degree  $k$ :

# One-Dimensional Representations of the Braid Group

For  $\theta \in \mathbb{R}$  and  $j = 1, 2, \dots, n-1$ , we define some *one-dimensional representations* of  $B_n$ :

$$\begin{aligned} \rho_\theta : B_n &\rightarrow \mathbb{C}_{|z|=1} \\ \sigma_j &\mapsto e^{i\theta}. \end{aligned}$$

These representations are *abelian*:

$$\begin{aligned} \rho_\theta(\sigma_1 \sigma_2 \sigma_1^{-1} \sigma_2) &= \rho_\theta(\sigma_1) \rho_\theta(\sigma_2) \rho_\theta(\sigma_1^{-1}) \rho_\theta(\sigma_2) \\ &= e^{i\theta_1} e^{i\theta_2} e^{-i\theta_1} e^{i\theta_2} \\ &= e^{i(\theta_1 - \theta_1 + \theta_2 + \theta_2)} \\ &= e^{i \cdot 2\theta_2} = \rho_\theta(\sigma_2^2) \end{aligned}$$

Hence, for any  $\beta \in B_n$  with degree  $k$ :

$$\rho_\theta(\beta) = \rho_\theta(\sigma_1^{m_1} \sigma_2^{m_2} \cdots \sigma_{n-1}^{m_{n-1}}) = e^{i\theta(m_1 + m_2 + \cdots + m_{n-1})} = e^{ik\theta}.$$

# The Burau Representation

# The Burau Representation

Define the matrix  $U = \begin{bmatrix} 1-t & t \\ 1 & 0 \end{bmatrix}$ , where  $t$  is a free parameter.

# The Burau Representation

Define the matrix  $U = \begin{bmatrix} 1-t & t \\ 1 & 0 \end{bmatrix}$ , where  $t$  is a free parameter.

## Definition

The *Burau representation* of the braid group  $B_n$  is defined on the standard generators:

$$\psi_n : B_n \rightarrow \mathrm{GL}_n(\mathbb{Z}[t, t^{-1}])$$
$$\sigma_i \mapsto \begin{bmatrix} I_{i-1} & 0 & 0 \\ 0 & U & 0 \\ 0 & 0 & I_{n-i-1} \end{bmatrix}.$$



# The Burau Representation

Define the matrix  $U = \begin{bmatrix} 1-t & t \\ 1 & 0 \end{bmatrix}$ , where  $t$  is a free parameter.

## Definition

The *Burau representation* of the braid group  $B_n$  is defined on the standard generators:

$$\begin{aligned} \psi_n : B_n &\rightarrow \mathrm{GL}_n(\mathbb{Z}[t, t^{-1}]) \\ \sigma_i &\mapsto \begin{bmatrix} I_{i-1} & 0 & 0 \\ 0 & U & 0 \\ 0 & 0 & I_{n-i-1} \end{bmatrix}. \end{aligned}$$

The Burau representation satisfies the braid relations:

$$\begin{aligned} \psi_n(\sigma_i)\psi_n(\sigma_j) &= \psi_n(\sigma_j)\psi_n(\sigma_i) \text{ for } |i-j| > 1, \\ \psi_n(\sigma_i)\psi_n(\sigma_{i+1})\psi_n(\sigma_i) &= \psi_n(\sigma_{i+1})\psi_n(\sigma_i)\psi_n(\sigma_{i+1}) \text{ for } i \in \{1, \dots, n-2\}. \end{aligned}$$

# The Burau Representation is Reducible

# The Burau Representation is Reducible

**Notice:**  $U \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1-t & t \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1-t+t \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

# The Burau Representation is Reducible

**Notice:**  $U \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1-t & t \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1-t+t \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

**Block structure of  $\psi_n(\sigma_i) \implies \mathbf{1} = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}$  is invariant under  $\psi_n(\sigma_i) \forall i = 1, 2, \dots, n-1$**

# The Burau Representation is Reducible

**Notice:**  $U \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1-t & t \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1-t+t \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

**Block structure of  $\psi_n(\sigma_i)$**   $\implies \mathbf{1} = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}$  is invariant under  $\psi_n(\sigma_i) \forall i = 1, 2, \dots, n-1$

$\implies \psi_n(\beta)\mathbf{1} = \mathbf{1} \forall \beta \in B_n$  (span $\{\mathbf{1}\}$  is  $\psi_n$ -invariant)

# The Burau Representation is Reducible

**Notice:**  $U \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1-t & t \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1-t+t \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

**Block structure of  $\psi_n(\sigma_i)$**   $\implies \mathbf{1} = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}$  is invariant under  $\psi_n(\sigma_i) \forall i = 1, 2, \dots, n-1$

$$\implies \psi_n(\beta)\mathbf{1} = \mathbf{1} \forall \beta \in B_n \quad (\text{span}\{\mathbf{1}\} \text{ is } \psi_n\text{-invariant})$$

$$\implies \text{Burau representation is reducible!}$$

# Unitary Representation of the Braid Group

# Unitary Representation of the Braid Group

## Definition

A matrix  $M \in \mathrm{GL}_n(\mathbb{C})$  is *unitary* if  $M^\dagger = M^{-1}$ .



# Unitary Representation of the Braid Group

## Definition

A matrix  $M \in \text{GL}_n(\mathbb{C})$  is *unitary* if  $M^\dagger = M^{-1}$ .

- ▶ The *reduced Burau representation* on  $B_n$  is an  $(n - 1)$ -dimensional representation of the braid group.

# Unitary Representation of the Braid Group

## Definition

A matrix  $M \in \text{GL}_n(\mathbb{C})$  is *unitary* if  $M^\dagger = M^{-1}$ .

- ▶ The *reduced Burau representation* on  $B_n$  is an  $(n - 1)$ -dimensional representation of the braid group.
- ▶ *Unitary representations* of  $B_n$  can be constructed from the reduced Burau representation.

# Unitary Representation of the Braid Group

## Definition

A matrix  $M \in \text{GL}_n(\mathbb{C})$  is *unitary* if  $M^\dagger = M^{-1}$ .

- ▶ The *reduced Burau representation* on  $B_n$  is an  $(n-1)$ -dimensional representation of the braid group.
- ▶ *Unitary representations* of  $B_n$  can be constructed from the reduced Burau representation.

## Definition

Define the unitary representation  $\mathcal{U} : B_3 \rightarrow U(2)$  by

$$\mathcal{U}(\sigma_1) = \frac{1}{2}e^{-i\frac{\pi}{6}} \begin{bmatrix} \sqrt{3}e^{i\arctan\left(\frac{1}{\sqrt{2}}\right)} & 1 \\ 1 & -\sqrt{3}e^{-i\arctan\left(\frac{1}{\sqrt{2}}\right)} \end{bmatrix}$$

$$\mathcal{U}(\sigma_2) = \frac{1}{2}e^{-i\frac{\pi}{6}} \begin{bmatrix} -\sqrt{3}e^{-i\arctan\left(\frac{1}{\sqrt{2}}\right)} & 1 \\ 1 & \sqrt{3}e^{i\arctan\left(\frac{1}{\sqrt{2}}\right)} \end{bmatrix}$$

# Nonabelian Characteristics of the Unitary Representation

# Nonabelian Characteristics of the Unitary Representation

**Observations:**

# Nonabelian Characteristics of the Unitary Representation

## Observations:

1.  $[\mathcal{U}(\sigma_1), \mathcal{U}(\sigma_2)] \neq 0 \implies \mathcal{U} \text{ nonabelian}.$

# Nonabelian Characteristics of the Unitary Representation

## Observations:

1.  $[\mathcal{U}(\sigma_1), \mathcal{U}(\sigma_2)] \neq 0 \implies \mathcal{U} \text{ nonabelian}.$
2.  $\mathcal{U}(\sigma_i)^{-1} = \mathcal{U}(\sigma_i)^\dagger \neq \mathcal{U}(\sigma_i)$  for  $i = 1, 2.$

# Nonabelian Characteristics of the Unitary Representation

## Observations:

1.  $[\mathcal{U}(\sigma_1), \mathcal{U}(\sigma_2)] \neq 0 \implies \mathcal{U} \text{ nonabelian.}$
2.  $\mathcal{U}(\sigma_i)^{-1} = \mathcal{U}(\sigma_i)^\dagger \neq \mathcal{U}(\sigma_i)$  for  $i = 1, 2$ .

**Consequence:**  $\sigma_1^2$  and  $\sigma_2^2$  are not the identity braid, which is in contrast to the permutation group where transpositions are involutory.



# Nonabelian Characteristics of the Unitary Representation

## Observations:

1.  $[\mathcal{U}(\sigma_1), \mathcal{U}(\sigma_2)] \neq 0 \implies \mathcal{U} \text{ nonabelian.}$
2.  $\mathcal{U}(\sigma_i)^{-1} = \mathcal{U}(\sigma_i)^\dagger \neq \mathcal{U}(\sigma_i)$  for  $i = 1, 2$ .

**Consequence:**  $\sigma_1^2$  and  $\sigma_2^2$  are not the identity braid, which is in contrast to the permutation group where transpositions are involutory.

## Question

What are the physical implications of this nonabelian unitary representation?

# Nonabelian Characteristics of the Unitary Representation

## Observations:

1.  $[\mathcal{U}(\sigma_1), \mathcal{U}(\sigma_2)] \neq 0 \implies \mathcal{U} \text{ nonabelian.}$
2.  $\mathcal{U}(\sigma_i)^{-1} = \mathcal{U}(\sigma_i)^\dagger \neq \mathcal{U}(\sigma_i)$  for  $i = 1, 2$ .

**Consequence:**  $\sigma_1^2$  and  $\sigma_2^2$  are not the identity braid, which is in contrast to the permutation group where transpositions are involutory.

## Question

What are the physical implications of this nonabelian unitary representation?

**Answer:** Unitary matrices preserve inner products, so the unitary representations of the braid group can act on a quantum system by braiding particles!



**CAL POLY**

## **4 Physical Applications of the Braid Group**

# (Abelian) Braiding Action on a Quantum System

# (Abelian) Braiding Action on a Quantum System

**1D Representation:** Let  $p_\theta : B_n \rightarrow \mathbb{C}$  be defined by  $\sigma_j \mapsto e^{i\theta}$  for some  $\theta$ , for all  $j$ .

# (Abelian) Braiding Action on a Quantum System

**1D Representation:** Let  $p_\theta : B_n \rightarrow \mathbb{C}$  be defined by  $\sigma_j \mapsto e^{i\theta}$  for some  $\theta$ , for all  $j$ .

**Quantum system:** Some wavefunction  $\psi(r_1, \dots, r_n)$  describing the identical particles fixed at nondegenerate positions  $r_1, r_2, \dots, r_n$ .

# (Abelian) Braiding Action on a Quantum System

**1D Representation:** Let  $\rho_\theta : B_n \rightarrow \mathbb{C}$  be defined by  $\sigma_j \mapsto e^{i\theta}$  for some  $\theta$ , for all  $j$ .

**Quantum system:** Some wavefunction  $\psi(r_1, \dots, r_n)$  describing the identical particles fixed at nondegenerate positions  $r_1, r_2, \dots, r_n$ .

**Braiding action:** For any degree- $k$  braid  $\beta \in B_n$ , we have

$$\psi(r_{1'}, r_{2'}, \dots, r_{n'}) = \rho_\theta(\beta) \psi(r_1, r_2, \dots, r_n) = \underbrace{e^{ik\theta}}_{\substack{\text{phase} \\ \text{shift}}} \psi(r_1, r_2, \dots, r_n),$$

# (Nonabelian) Braiding Action on a Quantum System



# (Nonabelian) Braiding Action on a Quantum System

**2D Representation:** Consider the  $2 \times 2$  unitary representation  $\mathcal{U}$  from before.

# (Nonabelian) Braiding Action on a Quantum System

**2D Representation:** Consider the  $2 \times 2$  unitary representation  $\mathcal{U}$  from before.

**Quantum system:** A degenerate set of two quantum states with orthonormal basis  $\psi_1(r_1, r_2, r_3)$  and  $\psi_2(r_1, r_2, r_3)$ . Shorthand:  $|1\rangle$  and  $|2\rangle$ .

# (Nonabelian) Braiding Action on a Quantum System

**2D Representation:** Consider the  $2 \times 2$  unitary representation  $\mathcal{U}$  from before.

**Quantum system:** A degenerate set of two quantum states with orthonormal basis  $\psi_1(r_1, r_2, r_3)$  and  $\psi_2(r_1, r_2, r_3)$ . Shorthand:  $|1\rangle$  and  $|2\rangle$ .

**Braiding action:** The transformed basis states due to the action of  $\sigma_1$  are

$$|1'\rangle = \mathcal{U}(\sigma_1)_{1,1} |1\rangle + \mathcal{U}(\sigma_1)_{1,2} |2\rangle = \frac{1}{2} e^{-i\frac{\pi}{6}} \left( \sqrt{3} e^{i\arctan\left(\frac{1}{\sqrt{2}}\right)} |1\rangle + |2\rangle \right),$$

$$|2'\rangle = \mathcal{U}(\sigma_1)_{2,1} |1\rangle + \mathcal{U}(\sigma_1)_{2,2} |2\rangle = \frac{1}{2} e^{-i\frac{\pi}{6}} \left( |1\rangle - \sqrt{3} e^{-i\arctan\left(\frac{1}{\sqrt{2}}\right)} |2\rangle \right).$$

# (Nonabelian) Braiding Action on a Quantum System

**2D Representation:** Consider the  $2 \times 2$  unitary representation  $\mathcal{U}$  from before.

**Quantum system:** A degenerate set of two quantum states with orthonormal basis  $\psi_1(r_1, r_2, r_3)$  and  $\psi_2(r_1, r_2, r_3)$ . Shorthand:  $|1\rangle$  and  $|2\rangle$ .

**Braiding action:** The transformed basis states due to the action of  $\sigma_1$  are

$$|1'\rangle = \mathcal{U}(\sigma_1)_{1,1} |1\rangle + \mathcal{U}(\sigma_1)_{1,2} |2\rangle = \frac{1}{2} e^{-i\frac{\pi}{6}} \left( \sqrt{3} e^{i\arctan\left(\frac{1}{\sqrt{2}}\right)} |1\rangle + |2\rangle \right),$$

$$|2'\rangle = \mathcal{U}(\sigma_1)_{2,1} |1\rangle + \mathcal{U}(\sigma_1)_{2,2} |2\rangle = \frac{1}{2} e^{-i\frac{\pi}{6}} \left( |1\rangle - \sqrt{3} e^{-i\arctan\left(\frac{1}{\sqrt{2}}\right)} |2\rangle \right).$$

## Remark

The action of a nonabelian braid group representation on a quantum system leads to *nontrivial rotations* in the many-particle Hilbert space that describes the quantum system<sup>9</sup>.

<sup>9</sup>Nayak et al., 2008, Non-abelian anyons and topological quantum computation, *Reviews of Modern Physics*

# Anyons: A Consequence of Braiding

## Definition

Particles that obey the braid group permutation rules are known as *anyons*.

# Anyons: A Consequence of Braiding

## Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- Anyons are  $(2 + 1)$ -*dimensional* quasi-particles (2D space + 1D time).

# Anyons: A Consequence of Braiding

## Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- ▶ Anyons are  $(2 + 1)$ -*dimensional* quasi-particles (2D space + 1D time).
- ▶ Anyon statistics are governed by the specific braid group representation acting on the system.

# Anyons: A Consequence of Braiding

## Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- ▶ Anyons are  $(2 + 1)$ -*dimensional* quasi-particles (2D space + 1D time).
- ▶ Anyon statistics are governed by the specific braid group representation acting on the system.
- ▶ Two types of anyons:



# Anyons: A Consequence of Braiding

## Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- ▶ Anyons are  $(2 + 1)$ -*dimensional* quasi-particles (2D space + 1D time).
- ▶ Anyon statistics are governed by the specific braid group representation acting on the system.
- ▶ Two types of anyons:
  1. *Abelian anyons*: The braid group representation is abelian.

# Anyons: A Consequence of Braiding

## Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- ▶ Anyons are  $(2 + 1)$ -*dimensional* quasi-particles (2D space + 1D time).
- ▶ Anyon statistics are governed by the specific braid group representation acting on the system.
- ▶ Two types of anyons:
  1. *Abelian anyons*: The braid group representation is abelian.
  2. *Nonabelian anyons*: The braid group representation is nonabelian.

# Anyons: A Consequence of Braiding

## Definition

Particles that obey the braid group permutation rules are known as *anyons*.

- ▶ Anyons are  $(2 + 1)$ -*dimensional* quasi-particles (2D space + 1D time).
- ▶ Anyon statistics are governed by the specific braid group representation acting on the system.
- ▶ Two types of anyons:
  1. *Abelian anyons*: The braid group representation is abelian.
  2. *Nonabelian anyons*: The braid group representation is nonabelian.
- ▶ Edge cases: *bosons* and *fermions*.

# Nontrivial Braiding Effects in 1D Representations

# Nontrivial Braiding Effects in 1D Representations

**Recall:** A braid is only well-defined if all particle trajectories are known.

# Nontrivial Braiding Effects in 1D Representations

**Recall:** A braid is only well-defined if all particle trajectories are known.

**Consequences:**

1. A permutation of two anyons requires the knowledge of the positions of all other anyons in the system.

# Nontrivial Braiding Effects in 1D Representations

**Recall:** A braid is only well-defined if all particle trajectories are known.

**Consequences:**

1. A permutation of two anyons requires the knowledge of the positions of all other anyons in the system.
2. This is a consequence of the so-called *nontrivial braiding effects* of the braid group.


# Nontrivial Braiding Effects in 1D Representations

**Recall:** A braid is only well-defined if all particle trajectories are known.

**Consequences:**

1. A permutation of two anyons requires the knowledge of the positions of all other anyons in the system.
2. This is a consequence of the so-called *nontrivial braiding effects* of the braid group.

Trajectory A






# Nontrivial Braiding Effects in 1D Representations

**Recall:** A braid is only well-defined if all particle trajectories are known.


**Consequences:**

1. A permutation of two anyons requires the knowledge of the positions of all other anyons in the system.
2. This is a consequence of the so-called *nontrivial braiding effects* of the braid group.

Trajectory A



Trajectory B





# Nontrivial Braiding Effects in 1D Representations

**Recall:** A braid is only well-defined if all particle trajectories are known.



**Consequences:**

1. A permutation of two anyons requires the knowledge of the positions of all other anyons in the system.
2. This is a consequence of the so-called *nontrivial braiding effects* of the braid group.

Trajectory A



Trajectory B



# Nontrivial Braiding Effects in 1D Representations

**Recall:** A braid is only well-defined if all particle trajectories are known.

## Consequences:



1. A permutation of two anyons requires the knowledge of the positions of all other anyons in the system.
2. This is a consequence of the so-called *nontrivial braiding effects* of the braid group.

## 1D representation:



$$\sigma_1 \mapsto e^{i\theta}$$

$$\sigma_2 \sigma_1 \sigma_2 \mapsto e^{3i\theta}$$

### Trajectory A



### Trajectory B



# Nontrivial Braiding Effects in 1D Representations

**Recall:** A braid is only well-defined if all particle trajectories are known.



## Consequences:

1. A permutation of two anyons requires the knowledge of the positions of all other anyons in the system.
2. This is a consequence of the so-called *nontrivial braiding effects* of the braid group.



## 1D representation:

$$\left. \begin{aligned} \sigma_1 &\mapsto e^{i\theta} \\ \sigma_2\sigma_1\sigma_2 &\mapsto e^{3i\theta} \end{aligned} \right\} \neq \text{if } \theta \notin \pi\mathbb{Z}$$

### Trajectory A



### Trajectory B



# A Physicists Approach to Anyons (Lagrangian)

Consider two identical non-interacting anyons with positions  $\mathbf{r}_1 = (x_1, y_1)$  and  $\mathbf{r}_2 = (x_2, y_2)$  in a harmonic potential. Let  $\phi = \arctan\left(\frac{y_2 - y_1}{x_2 - x_1}\right)$  be the relative angle between the two anyons and  $\dot{\phi} = \frac{d\phi}{dt}$ .

# A Physicists Approach to Anyons (Lagrangian)

Consider two identical non-interacting anyons with positions  $\mathbf{r}_1 = (x_1, y_1)$  and  $\mathbf{r}_2 = (x_2, y_2)$  in a harmonic potential. Let  $\phi = \arctan\left(\frac{y_2 - y_1}{x_2 - x_1}\right)$  be the relative angle between the two anyons and  $\dot{\phi} = \frac{d\phi}{dt}$ .

*Potential:*  $V(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{2}m\omega^2 (\mathbf{r}_1^2 + \mathbf{r}_2^2)$

# A Physicists Approach to Anyons (Lagrangian)

Consider two identical non-interacting anyons with positions  $\mathbf{r}_1 = (x_1, y_1)$  and  $\mathbf{r}_2 = (x_2, y_2)$  in a harmonic potential. Let  $\phi = \arctan\left(\frac{y_2 - y_1}{x_2 - x_1}\right)$  be the relative angle between the two anyons and  $\dot{\phi} = \frac{d\phi}{dt}$ .

*Potential:*  $V(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{2}m\omega^2 (\mathbf{r}_1^2 + \mathbf{r}_2^2)$

*Statistical interaction due to braiding:*  $\mathcal{L}_{\text{int}} = \hbar\alpha\dot{\phi}, \quad \alpha \in [0, 1]$

# A Physicists Approach to Anyons (Lagrangian)

Consider two identical non-interacting anyons with positions  $\mathbf{r}_1 = (x_1, y_1)$  and  $\mathbf{r}_2 = (x_2, y_2)$  in a harmonic potential. Let  $\phi = \arctan\left(\frac{y_2 - y_1}{x_2 - x_1}\right)$  be the relative angle between the two anyons and  $\dot{\phi} = \frac{d\phi}{dt}$ .

*Potential:*  $V(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{2}m\omega^2 (\mathbf{r}_1^2 + \mathbf{r}_2^2)$

*Statistical interaction due to braiding:*  $\mathcal{L}_{\text{int}} = \hbar\alpha\dot{\phi}, \quad \alpha \in [0, 1]$

*Classical Kinetic Energy:*  $T = \frac{1}{2}m (\dot{\mathbf{r}}_1^2 + \dot{\mathbf{r}}_2^2)$



# A Physicists Approach to Anyons (Lagrangian)

Consider two identical non-interacting anyons with positions  $\mathbf{r}_1 = (x_1, y_1)$  and  $\mathbf{r}_2 = (x_2, y_2)$  in a harmonic potential. Let  $\phi = \arctan\left(\frac{y_2 - y_1}{x_2 - x_1}\right)$  be the relative angle between the two anyons and  $\dot{\phi} = \frac{d\phi}{dt}$ .

*Potential:*  $V(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{2}m\omega^2 (\mathbf{r}_1^2 + \mathbf{r}_2^2)$

*Statistical interaction due to braiding:*  $\mathcal{L}_{\text{int}} = \hbar\alpha\dot{\phi}, \quad \alpha \in [0, 1]$

*Classical Kinetic Energy:*  $T = \frac{1}{2}m (\dot{\mathbf{r}}_1^2 + \dot{\mathbf{r}}_2^2)$

**Lagrangian:**

$$\mathcal{L}(r_1, r_2, \dot{\mathbf{r}}_1, \dot{\mathbf{r}}_2, \dot{\phi}) = T + \mathcal{L}_{\text{int}} - V(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{2}m (\dot{\mathbf{r}}_1^2 + \dot{\mathbf{r}}_2^2) + \hbar\alpha\dot{\phi} - \frac{1}{2}m\omega^2 (\mathbf{r}_1^2 + \mathbf{r}_2^2)$$

# A Physicists Approach to Anyons (Lagrangian)

Consider two identical non-interacting anyons with positions  $\mathbf{r}_1 = (x_1, y_1)$  and  $\mathbf{r}_2 = (x_2, y_2)$  in a harmonic potential. Let  $\phi = \arctan\left(\frac{y_2 - y_1}{x_2 - x_1}\right)$  be the relative angle between the two anyons and  $\dot{\phi} = \frac{d\phi}{dt}$ .

*Potential:*  $V(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{2}m\omega^2 (\mathbf{r}_1^2 + \mathbf{r}_2^2)$

*Statistical interaction due to braiding:*  $\mathcal{L}_{\text{int}} = \hbar\alpha\dot{\phi}, \quad \alpha \in [0, 1]$

*Classical Kinetic Energy:*  $T = \frac{1}{2}m (\dot{\mathbf{r}}_1^2 + \dot{\mathbf{r}}_2^2)$

**Lagrangian:**

$$\mathcal{L}(r_1, r_2, \dot{\mathbf{r}}_1, \dot{\mathbf{r}}_2, \dot{\phi}) = T + \mathcal{L}_{\text{int}} - V(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{2}m (\dot{\mathbf{r}}_1^2 + \dot{\mathbf{r}}_2^2) + \hbar\alpha\dot{\phi} - \frac{1}{2}m\omega^2 (\mathbf{r}_1^2 + \mathbf{r}_2^2)$$

**Generalize to  $N$  anyons:** Let  $\phi_{ij} = \arctan\left(\frac{y_j - y_i}{x_j - x_i}\right)$ ,

$$\mathcal{L} = \sum_{i=1}^N \frac{m}{2} \dot{\mathbf{r}}_i^2 + \hbar\alpha \sum_{i < j}^N \dot{\phi}_{ij} - \frac{m\omega^2}{2} \sum_{i=1}^N \mathbf{r}_i^2$$

# A Physicists Approach to Anyons (Hamiltonian)

# A Physicists Approach to Anyons (Hamiltonian)

Rewrite  $N$ -anyon  $\mathcal{L}$ :

$$\mathcal{L} = \frac{m}{2} \sum_{i=1}^N [\dot{\mathbf{r}}^2 - \omega^2 \mathbf{r}_i^2] + \alpha \sum_{i < j}^N \dot{\mathbf{r}}_{ij} \cdot \frac{(-y_{ij} \hat{X} + x_{ij} \hat{Y})}{r_{ij}^2}$$

# A Physicists Approach to Anyons (Hamiltonian)

Rewrite  $N$ -anyon  $\mathcal{L}$ :

$$\mathcal{L} = \frac{m}{2} \sum_{i=1}^N [\dot{\mathbf{r}}^2 - \omega^2 \mathbf{r}_i^2] + \alpha \sum_{i < j} \dot{\mathbf{r}}_{ij} \cdot \frac{(-y_{ij} \hat{x} + x_{ij} \hat{y})}{r_{ij}^2}$$

Gauge potential:

$$\mathbf{A}_i(\mathbf{r}_i) = \alpha \sum_{j \neq i} \frac{\hat{z} \times \mathbf{r}_{ij}}{r_{ij}^2} = \alpha \sum_{j \neq i} \frac{-y_{ij} \hat{x} + x_{ij} \hat{y}}{r_{ij}^2}$$

# A Physicists Approach to Anyons (Hamiltonian)

Rewrite  $N$ -anyon  $\mathcal{L}$ :

$$\mathcal{L} = \frac{m}{2} \sum_{i=1}^N [\dot{\mathbf{r}}^2 - \omega^2 \mathbf{r}_i^2] + \alpha \sum_{i < j} \dot{\mathbf{r}}_{ij} \cdot \frac{(-y_{ij} \hat{x} + x_{ij} \hat{y})}{r_{ij}^2}$$

Gauge potential:

$$\mathbf{A}_i(\mathbf{r}_i) = \alpha \sum_{j \neq i} \frac{\hat{z} \times \mathbf{r}_{ij}}{r_{ij}^2} = \alpha \sum_{j \neq i} \frac{-y_{ij} \hat{x} + x_{ij} \hat{y}}{r_{ij}^2}$$

$i$ -th anyon Hamiltonian:

$$\mathcal{H}_i = \frac{1}{2m} \underbrace{(\mathbf{p}_i - \mathbf{A}_i(\mathbf{r}_i))^2}_{\text{canonical momentum}} + \frac{m\omega^2}{2} r_i^2$$

# A Physicists Approach to Anyons (Hamiltonian)

Rewrite  $N$ -anyon  $\mathcal{L}$ :

$$\mathcal{L} = \frac{m}{2} \sum_{i=1}^N [\dot{\mathbf{r}}^2 - \omega^2 \mathbf{r}_i^2] + \alpha \sum_{i < j} \dot{\mathbf{r}}_{ij} \cdot \frac{(-y_{ij} \hat{x} + x_{ij} \hat{y})}{r_{ij}^2}$$

Gauge potential:

$$\mathbf{A}_i(\mathbf{r}_i) = \alpha \sum_{j \neq i} \frac{\hat{\mathbf{z}} \times \mathbf{r}_{ij}}{r_{ij}^2} = \alpha \sum_{j \neq i} \frac{-y_{ij} \hat{x} + x_{ij} \hat{y}}{r_{ij}^2}$$

$i$ -th anyon Hamiltonian:

$$\mathcal{H}_i = \frac{1}{2m} \underbrace{(\mathbf{p}_i - \mathbf{A}_i(\mathbf{r}_i))^2}_{\text{canonical momentum}} + \frac{m\omega^2}{2} r_i^2$$

$N$ -anyon Hamiltonian:

$$\mathcal{H} = \frac{1}{2m} \sum_{i=1}^N (\mathbf{p}_i - \mathbf{A}_i(\mathbf{r}_i))^2 + \frac{m\omega^2}{2} \sum_{i=1}^N r_i^2$$

# A Physicists Approach to Anyons (Hamiltonian)

Rewrite  $N$ -anyon  $\mathcal{L}$ :

$$\mathcal{L} = \frac{m}{2} \sum_{i=1}^N [\dot{\mathbf{r}}^2 - \omega^2 \mathbf{r}_i^2] + \alpha \sum_{i < j} \dot{\mathbf{r}}_{ij} \cdot \frac{(-y_{ij} \hat{x} + x_{ij} \hat{y})}{r_{ij}^2}$$

Gauge potential:

$$\mathbf{A}_i(\mathbf{r}_i) = \alpha \sum_{j \neq i} \frac{\hat{\mathbf{z}} \times \mathbf{r}_{ij}}{r_{ij}^2} = \alpha \sum_{j \neq i} \frac{-y_{ij} \hat{x} + x_{ij} \hat{y}}{r_{ij}^2}$$

$i$ -th anyon Hamiltonian:

$$\mathcal{H}_i = \frac{1}{2m} \underbrace{(\mathbf{p}_i - \mathbf{A}_i(\mathbf{r}_i))^2}_{\text{canonical momentum}} + \frac{m\omega^2}{2} r_i^2$$

$N$ -anyon Hamiltonian:

$$\mathcal{H} = \frac{1}{2m} \sum_{i=1}^N (\mathbf{p}_i - \mathbf{A}_i(\mathbf{r}_i))^2 + \frac{m\omega^2}{2} \sum_{i=1}^N r_i^2$$

Expand:

$$\mathcal{H} = \frac{1}{2m} \sum_{i=1}^N p_i^2 + \frac{m\omega^2}{2} \sum_{i=1}^N r_i^2 - \frac{\alpha}{2m} \sum_{\substack{i=1 \\ j \neq i}}^N \frac{\ell_{ij}}{r_{ij}^2} + \frac{\alpha^2}{2m} \sum_{\substack{i=1 \\ j, k \neq i}}^N \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2}$$



# Interpreting the $N$ -anyon Hamiltonian

$$\mathcal{H} = \frac{1}{2m} \sum_{i=1}^N p_i^2 + \frac{m\omega^2}{2} \sum_{i=1}^N r_i^2 - \frac{\alpha}{2m} \sum_{\substack{i=1 \\ j \neq i}}^N \frac{\ell_{ij}}{r_{ij}^2} + \frac{\alpha^2}{2m} \sum_{\substack{i=1 \\ j, k \neq i}}^N \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2}$$

# Interpreting the $N$ -anyon Hamiltonian

$$\mathcal{H} = \underbrace{\frac{1}{2m} \sum_{i=1}^N p_i^2}_{\text{Mechanical momentum}} + \frac{m\omega^2}{2} \sum_{i=1}^N r_i^2 - \frac{\alpha}{2m} \sum_{\substack{i=1 \\ j \neq i}}^N \frac{\ell_{ij}}{r_{ij}^2} + \frac{\alpha^2}{2m} \sum_{\substack{i=1 \\ j, k \neq i}}^N \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2}$$

# Interpreting the $N$ -anyon Hamiltonian

$$\mathcal{H} = \underbrace{\frac{1}{2m} \sum_{i=1}^N p_i^2}_{\text{Mechanical momentum}} + \underbrace{\frac{m\omega^2}{2} \sum_{i=1}^N r_i^2}_{\text{Harmonic potential}} - \frac{\alpha}{2m} \sum_{\substack{i=1 \\ j \neq i}}^N \frac{\ell_{ij}}{r_{ij}^2} + \frac{\alpha^2}{2m} \sum_{\substack{i=1 \\ j, k \neq i}}^N \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2}$$

# Interpreting the $N$ -anyon Hamiltonian

$$\mathcal{H} = \underbrace{\frac{1}{2m} \sum_{i=1}^N p_i^2}_{\text{Mechanical momentum}} + \underbrace{\frac{m\omega^2}{2} \sum_{i=1}^N r_i^2}_{\text{Harmonic potential}} - \underbrace{\frac{\alpha}{2m} \sum_{\substack{i=1 \\ j \neq i}}^N \frac{\ell_{ij}}{r_{ij}^2}}_{\substack{\text{Relative angular} \\ \text{momentum} \\ \ell_{ij} = \mathbf{r}_{ij} \times \mathbf{p}_{ij}}} + \frac{\alpha^2}{2m} \sum_{\substack{i=1 \\ j, k \neq i}}^N \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2}$$

# Interpreting the $N$ -anyon Hamiltonian

$$\mathcal{H} = \underbrace{\frac{1}{2m} \sum_{i=1}^N p_i^2}_{\text{Mechanical momentum}} + \underbrace{\frac{m\omega^2}{2} \sum_{i=1}^N r_i^2}_{\text{Harmonic potential}} - \underbrace{\frac{\alpha}{2m} \sum_{\substack{i=1 \\ j \neq i}}^N \frac{\ell_{ij}}{r_{ij}^2}}_{\substack{\text{Relative angular} \\ \text{momentum} \\ \ell_{ij} = \mathbf{r}_{ij} \times \mathbf{p}_{ij}}} + \underbrace{\frac{\alpha^2}{2m} \sum_{\substack{i=1 \\ j, k \neq i}}^N \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2}}_{\text{Long-range interaction}}$$

# Nontrivial Braiding Effects in the Hamiltonian

Nontrivial braiding effects emerge from the *long-range interaction* term when  $N \geq 3$ .

# Nontrivial Braiding Effects in the Hamiltonian

Nontrivial braiding effects emerge from the *long-range interaction* term when  $N \geq 3$ .

$$\mathbf{N} = 2: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j, k \neq i}}^2 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{2m} \left( \frac{\mathbf{r}_{12} \cdot \mathbf{r}_{12}}{r_{12}^2 r_{12}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{21}}{r_{21}^2 r_{21}^2} \right) = \frac{\alpha^2}{mr_{12}^2}$$

# Nontrivial Braiding Effects in the Hamiltonian

Nontrivial braiding effects emerge from the *long-range interaction* term when  $N \geq 3$ .

$$\mathbf{N} = \mathbf{2}: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j, k \neq i}}^2 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{2m} \left( \frac{\mathbf{r}_{12} \cdot \mathbf{r}_{12}}{r_{12}^2 r_{12}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{21}}{r_{21}^2 r_{21}^2} \right) = \frac{\alpha^2}{mr_{12}^2} \longleftarrow \text{Coulomb-like interaction}$$



# Nontrivial Braiding Effects in the Hamiltonian

Nontrivial braiding effects emerge from the *long-range interaction* term when  $N \geq 3$ .

$$\mathbf{N} = 2: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j, k \neq i}}^2 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{2m} \left( \frac{\mathbf{r}_{12} \cdot \mathbf{r}_{12}}{r_{12}^2 r_{12}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{21}}{r_{21}^2 r_{21}^2} \right) = \frac{\alpha^2}{mr_{12}^2} \longleftarrow \text{Coulomb-like interaction}$$

$$\mathbf{N} = 3: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j, k \neq i}}^3 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{m} \left( \frac{1}{r_{12}^2} + \frac{1}{r_{13}^2} + \frac{1}{r_{23}^2} + \frac{\mathbf{r}_{12} \cdot \mathbf{r}_{13}}{r_{12}^2 r_{13}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{23}}{r_{21}^2 r_{23}^2} + \frac{\mathbf{r}_{31} \cdot \mathbf{r}_{32}}{r_{31}^2 r_{32}^2} \right)$$

# Nontrivial Braiding Effects in the Hamiltonian

Nontrivial braiding effects emerge from the *long-range interaction* term when  $N \geq 3$ .

$$\mathbf{N} = 2: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j,k \neq i}}^2 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{2m} \left( \frac{\mathbf{r}_{12} \cdot \mathbf{r}_{12}}{r_{12}^2 r_{12}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{21}}{r_{21}^2 r_{21}^2} \right) = \frac{\alpha^2}{mr_{12}^2} \longleftarrow \text{Coulomb-like interaction}$$

$$\mathbf{N} = 3: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j,k \neq i}}^3 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{m} \left( \underbrace{\frac{1}{r_{12}^2} + \frac{1}{r_{13}^2} + \frac{1}{r_{23}^2}}_{\text{Coulomb-like interaction}} + \frac{\mathbf{r}_{12} \cdot \mathbf{r}_{13}}{r_{12}^2 r_{13}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{23}}{r_{21}^2 r_{23}^2} + \frac{\mathbf{r}_{31} \cdot \mathbf{r}_{32}}{r_{31}^2 r_{32}^2} \right)$$

# Nontrivial Braiding Effects in the Hamiltonian

Nontrivial braiding effects emerge from the *long-range interaction* term when  $N \geq 3$ .

$$\mathbf{N} = 2: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j, k \neq i}}^2 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{2m} \left( \frac{\mathbf{r}_{12} \cdot \mathbf{r}_{12}}{r_{12}^2 r_{12}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{21}}{r_{21}^2 r_{21}^2} \right) = \frac{\alpha^2}{mr_{12}^2} \leftarrow \text{Coulomb-like interaction}$$

$$\mathbf{N} = 3: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j, k \neq i}}^3 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{m} \left( \underbrace{\frac{1}{r_{12}^2} + \frac{1}{r_{13}^2} + \frac{1}{r_{23}^2}}_{\text{Coulomb-like interaction}} + \underbrace{\frac{\mathbf{r}_{12} \cdot \mathbf{r}_{13}}{r_{12}^2 r_{13}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{23}}{r_{21}^2 r_{23}^2} + \frac{\mathbf{r}_{31} \cdot \mathbf{r}_{32}}{r_{31}^2 r_{32}^2}}_{\text{Nontrivial braiding}} \right)$$

# Nontrivial Braiding Effects in the Hamiltonian

Nontrivial braiding effects emerge from the *long-range interaction* term when  $N \geq 3$ .

$$\mathbf{N} = 2: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j,k \neq i}}^2 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{2m} \left( \frac{\mathbf{r}_{12} \cdot \mathbf{r}_{12}}{r_{12}^2 r_{12}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{21}}{r_{21}^2 r_{21}^2} \right) = \frac{\alpha^2}{mr_{12}^2} \leftarrow \text{Coulomb-like interaction}$$

$$\mathbf{N} = 3: \quad \frac{\alpha^2}{2m} \sum_{\substack{j=1 \\ j,k \neq i}}^3 \frac{\mathbf{r}_{ij} \cdot \mathbf{r}_{ik}}{r_{ij}^2 r_{ik}^2} = \frac{\alpha^2}{m} \left( \underbrace{\frac{1}{r_{12}^2} + \frac{1}{r_{13}^2} + \frac{1}{r_{23}^2}}_{\text{Coulomb-like interaction}} + \underbrace{\frac{\mathbf{r}_{12} \cdot \mathbf{r}_{13}}{r_{12}^2 r_{13}^2} + \frac{\mathbf{r}_{21} \cdot \mathbf{r}_{23}}{r_{21}^2 r_{23}^2} + \frac{\mathbf{r}_{31} \cdot \mathbf{r}_{32}}{r_{31}^2 r_{32}^2}}_{\text{Nontrivial braiding}} \right)$$

## Question

Why is this useful?

# Physical Implications of Nontrivial Braiding Effects

# Physical Implications of Nontrivial Braiding Effects

- The fractional quantum Hall effect is a physical manifestation of anyonic braiding in 2D electron systems (fractional charge, fractional statistics).

# Physical Implications of Nontrivial Braiding Effects

- ▶ The fractional quantum Hall effect is a physical manifestation of anyonic braiding in 2D electron systems (fractional charge, fractional statistics).
- ▶ Depending on the specific representation of the braid group, one can define topological properties of different flavors of anyons. The corresponding combination (fusion) rules can be used to describe the behavior of anyonic systems.

# Physical Implications of Nontrivial Braiding Effects

- ▶ The fractional quantum Hall effect is a physical manifestation of anyonic braiding in 2D electron systems (fractional charge, fractional statistics).
- ▶ Depending on the specific representation of the braid group, one can define topological properties of different flavors of anyons. The corresponding combination (fusion) rules can be used to describe the behavior of anyonic systems.
- ▶ Certain nonabelian anyons (defined by specific fusion rules) are alleged to enable fault-tolerant quantum computers, and is an ongoing area of research.



# Summary

## Main Takeaways:

# Summary

## Main Takeaways:

1. Representation theory is a powerful tool that can be used to obtain fundamental results in quantum mechanics and beyond.

# Summary

## Main Takeaways:

1. Representation theory is a powerful tool that can be used to obtain fundamental results in quantum mechanics and beyond.
2. Unitary representations of the braid group can act on  $(2 + 1)$ -dimensional quantum systems, resulting in anyons.

# Summary

## Main Takeaways:

1. Representation theory is a powerful tool that can be used to obtain fundamental results in quantum mechanics and beyond.
2. Unitary representations of the braid group can act on  $(2 + 1)$ -dimensional quantum systems, resulting in anyons.
3. Anyons exhibit fractional statistics in contrast to the boson/fermion dichotomy.

# Summary

## Main Takeaways:

1. Representation theory is a powerful tool that can be used to obtain fundamental results in quantum mechanics and beyond.
2. Unitary representations of the braid group can act on  $(2 + 1)$ -dimensional quantum systems, resulting in anyons.
3. Anyons exhibit fractional statistics in contrast to the boson/fermion dichotomy.
4. The nontrivial braiding effects of anyons results in useful physical properties that can be exploited for various physical applications.

# Summary

## Main Takeaways:

1. Representation theory is a powerful tool that can be used to obtain fundamental results in quantum mechanics and beyond.
2. Unitary representations of the braid group can act on  $(2 + 1)$ -dimensional quantum systems, resulting in anyons.
3. Anyons exhibit fractional statistics in contrast to the boson/fermion dichotomy.
4. The nontrivial braiding effects of anyons results in useful physical properties that can be exploited for various physical applications.

**Thank you for your attention!**

## SO(3) Calculations (pt. 1)

The state  $|\phi\rangle$  can be decomposed into a linear combination of the eigenvectors of  $J$ :

$$|\phi\rangle = \left( \sum_m |m\rangle \langle m| \right) |\phi\rangle = \sum_m \langle m|\phi\rangle |m\rangle,$$

where

$$\langle m|\phi\rangle = \langle m|U(\phi)|\mathcal{O}\rangle = \langle U^\dagger(\phi)m|\mathcal{O}\rangle = \langle e^{im\phi}m|\mathcal{O}\rangle = e^{-im\phi} \langle m|\mathcal{O}\rangle$$

is the projection of  $|\phi\rangle$  onto the eigenvector  $|m\rangle$  of  $J$ .

Thus,

$$\begin{aligned} J|\phi\rangle &= \sum_m e^{-im\phi} J|m\rangle = \sum_m m e^{-im\phi} |m\rangle = \sum_m i \frac{\partial}{\partial \phi} (e^{-im\phi} |m\rangle) = i \frac{\partial}{\partial \phi} |\phi\rangle \\ \implies \langle \phi|J|\psi\rangle &= \langle J^\dagger \phi|\psi\rangle = -i \frac{\partial}{\partial \phi} \langle \phi|\psi\rangle = -i \frac{\partial}{\partial \phi} \psi(\phi). \end{aligned}$$

## SO(3) Calculations (pt. 2)

$$\phi = \arctan\left(\frac{y}{x}\right)$$

$$\implies \frac{\partial}{\partial \phi} = (\mathbf{r} \times \nabla) \cdot \mathbf{e}_z \implies J = -i \frac{\partial}{\partial \phi} = -i (\mathbf{r} \times \nabla) \cdot \mathbf{e}_z = \frac{1}{\hbar} \hat{L}_z \implies \hat{\mathbf{L}} = \hat{\mathbf{r}} \times \hat{\mathbf{p}}$$

$$\hat{\mathbf{p}} = -i\nabla \implies \hat{L}_z = x\hat{p}_y - y\hat{p}_x$$

$$\hat{H} = \frac{\hat{\mathbf{p}}^2}{2m} + \hat{V}(\mathbf{r}), \quad [V(\mathbf{r}), \hat{L}_z] = 0, \quad [\hat{\mathbf{p}}^2, \hat{L}_z] = 0 \implies [\hat{H}, \hat{L}_z] = 0,$$

where the last line easily generalizes to  $\hat{\mathbf{L}}$ .



# Lie Algebra

$$\mathcal{J}^2 |j\rangle = (J_- J_+ + J_z + \mathcal{J}_z^2) |j\rangle = (0 + j + j^2) |j\rangle = j(j+1) |j\rangle ,$$

$$\mathcal{J}^2 |j, m\rangle = j(j+1) |j, m\rangle ,$$

$$J_z |j, m\rangle = m |j, m\rangle ,$$

$$J_{\pm} |j, m\rangle = \sqrt{j(j+1) - m(m \pm 1)} |j, m \pm 1\rangle .$$