

Demographic/Health Factor	Overall: N = 138
Age	
6-9 years	45 (32.6%)
10-13 years	71 (51.4%)
14-17 years	22 (15.9%)
Sex	
Male	57 (41.3%)
Female	81 (58.7%)
Weight (kg)	
15-35	87 (63.0%)
35-55	49 (35.5%)
>55	2 (1.4%)
Underweight	
Not Underweight	62 (44.9%)
Underweight	76 (55.1%)
Height (m)	
<1.15	7 (5.1%)
1.15-1.35	60 (43.5%)
1.35-1.55	59 (42.8%)
>1.55	12 (8.7%)
Growth Stunted	
Not Stunted	107 (77.5%)
Stunted	31 (22.5%)
Mode of Delivery	
Vaginal	126 (91.3%)
C-Section	12 (8.7%)
Ever Had Vaccination	
Yes	45 (32.6%)
No	93 (67.4%)
BCG Scar	
Yes	81 (58.7%)
No	57 (41.3%)
Deworming Pill	
Yes	90 (65.2%)
No	45 (32.6%)
NA/Don't Know	3 (2.2%)
Last Deworming Pill	
1-3 months	15 (16.9%)
4-6 months	5 (5.6%)
7-9 months	62 (69.7%)
>9 months	1 (1.1%)
Antibiotics Given	
Yes	17 (12.3%)
No	115 (83.3%)
NA/Don't Know	6 (4.3%)
Currently Prescribed Drug	
Yes	6 (4.3%)
No	128 (92.8%)
NA/Don't Know	4 (2.9%)
Anti-Malaria Drug in Last 3 Months	
Yes	2 (1.4%)
No	132 (95.6%)
NA/Don't Know	4 (2.9%)
Child Had Fever in Last 2 Weeks	
Yes	26 (18.8%)
No	110 (79.7%)
NA	2 (1.4%)

Demographic/Health Factor	Overall: N = 138
Currently Prescribed Drug	
Yes	6 (4.3%)
No	128 (92.8%)
NA/Don't Know	4 (2.9%)
Anti-Malaria Drug in Last 3 Months	
Yes	2 (1.4%)
No	132 (95.6%)
NA/Don't Know	4 (2.9%)
Child Had Fever in Last 2 Weeks	
Yes	26 (18.8%)
No	110 (79.7%)
NA	2 (1.4%)
Child Had Diarrhea in Last 2 Weeks	
Yes	19 (13.8%)
No	118 (85.5%)
Child Had Cough in Last 2 Weeks	
Yes	24 (17.4%)
No	113 (81.9%)
NA	1 (0.7%)
Child's Fingernails Trimmed	
Yes	79 (57.2%)
No	59 (42.8%)
Child Had Wheezing or Whistling in Chest	
Yes	0 (0.0%)
No	138 (100.0%)
Wheezing Attacks in Last Year	
0	97 (70.3%)
1-3	41 (29.7%)
Child Ever Had Asthma	
Yes	0 (0.0%)
No	138 (100.0%)
Child Had Hay Fever or Persistent Sneezing	
Yes	0 (0.0%)
No	138 (100.0%)
Child Ever Had Itchy Skin or Rash	
Yes	6 (4.3%)
No	132 (95.7%)
Body Part Affected by Rash	
Elbow Folds	14 (10.1%)
Behind the Knees	6 (4.3%)
Under the Buttocks	6 (4.3%)
Around the Neck	6 (4.3%)
Front of the Ankles	3 (2.2%)
Around the Eyes/Ears	0 (0.0%)
Child's Fingernails Dirty	
Yes	59 (42.8%)
No	79 (57.2%)
Frequency of Trimming Child's Fingernails	
Once Per Week	64 (46.4%)
Once Per Two Weeks	57 (41.3%)
Less Than Once/Month	11 (8.0%)
Don't Know	6 (4.3%)

Socioeconomic Factors	Overall: N = 138
Household Size	
2-3	6(4.3%)
4-6	91(65.9%)
7-9	37(26.8%)
>9	4(2.9%)
Older Siblings	
0	34(24.6%)
1-2	68(49.3%)
3-4	30(21.7%)
>4	6(4.3%)
Siblings Younger Than 12 Years Old	
0	28(20.3%)
1-2	87(63%)
3-4	22(15.9%)
>4	1(0.7%)
Knowledge of Intestinal Worm Transmission	
Yes	61(44.2%)
No	64(46.4%)
Don't Know	13(9.4%)
Knowledge of Why Worms Are Bad	
Yes	68(49.3%)
No	58(42%)
Don't Know	12(8.7%)
Knowledge of How to Avoid Worms	
Yes	61(44.2%)
No	65(47.1%)
Don't Know	12(8.7%)
Heard of Disease	
Ascharis	83(60.1%)
Trichuris	1(0.7%)
Hookworms	2(1.4%)
HIV/AIDS	94(68.1%)
Intestinal Worms	91(65.9%)
Malaria	98(71%)
Tuberculosis	27(19.6%)
Schistosoma/Bilharzia	11(8%)
None	18(13%)
Source of Disease Information	
Family	76(55.1%)
Health Professional	2(1.4%)
Teacher	51(37%)
Media	50(36.2%)
Don't Know	17(12.3%)
Residence	
Rural	35(25.4%)
Urban	103(74.6%)

Socioeconomic Factors	Overall: N = 138
Family Occupation	
Police	4(2.9%)
Teacher	5(3.6%)
Carpenter	9(6.5%)
Daily Labor	9(6.5%)
Security	9(6.5%)
Farmer	15(10.9%)
Government Employee	18(13%)
Merchant	25(18.1%)
Other	44(31.9%)
Maternal Education Status	
Illiterate	40(29%)
Primary School	55(39.9%)
High School	31(22.5%)
Higher Education	8(5.8%)
Don't Know	4(2.9%)
Technology in Household	
Electricity	129(93.5%)
Radio	70(50.7%)
TV	108(78.3%)
Family Member with Phone	131(94.9%)
Purpose of Phone Use	
Call/Radio/Internet	74(53.6%)
Call Only	58(42%)
Don't Know	6(4.3%)

Environmental Factors	Overall: N = 138
Latrine in School	
Yes	138(100%)
Latrine Has Doors	
Yes	78(56.5%)
No	60(43.5%)
Latrine Has Flies Around It	
Yes	137(99.3%)
No	1(0.7%)
Latrine Has Visible Stool Around It	
Yes	136(98.6%)
No	1(0.07%)
Don't Know	1(0.7%)
Use of School's Latrine	
Never	23(16.7%)
Sometimes	108(78.3%)
Always	5(3.6%)
NA/Don't Know	2(1.4%)
Family Owns a Latrine	
Yes	136(98.6%)
No	2(1.4%)
Latrine Location	
Outside	133(96.4%)
Inside	1(0.7%)
NA/Don't Know	4(2.9%)
Distance from House to Latrine (m)	
0-5	21(15.2%)
5-10	54(39.1%)
10-20	56(40.6%)
>20	1(0.7%)
NA/Don't Know	6(4.3%)
Distance from Kitchen to Latrine (m)	
0-5	30(21.7%)
5-10	59(42.8%)
10-20	40(29%)
>20	1(0.7%)
NA/Don't Know	8(5.8%)
Latrine Connected To	
Well	69(50%)
Ditch	7(5.1%)
River	5(3.6%)
Sewage	1(0.7%)
NA/Don't Know	56(40.6%)
House Floor Material	
Dust	51(37%)
Cement	47(34.1%)
Plastic Covered	6(4.3%)
Don't Know	43(24.6%)
Kitchen Location	
Separated	126(91.3%)
Within House	8(5.8%)
Don't Know	4(2.9%)
Kitchen Floor Material if Separated	
Dust	113(81.9%)
Cement	18(13%)
Don't Know	7(5.1%)
Kitchen Structures	
Roof	124(89.9%)
Walls	118(85.5%)
None	5(3.6%)

Environmental Factors	Overall: N = 138
Cooking Fuel Use	
Coal	114(82.6%)
Wood	98(71%)
Electric	58(42%)
Kerosene	3(2.2%)
Gas	1(0.7%)
Animals in House/Compound	
Pets	50(36.2%)
Chickens	42(30.4%)
Sheep/Goats	12(8.7%)
Cattle	10(7.2%)
None	59(42.8%)
Portable Water	
Yes	87(63%)
No	48(34.8%)
Don't Know	3(2.2%)
Source of Water if Not Portable	
River	21(30%)
Tap Water	18(25.7%)
Neighbor	17(24.3%)
Tank	6(8.6%)
Well	6(8.6%)
Truck	1(1.4%)
Water Treatment	
Yes	21(15.2%)
No	115(83.3%)
Don't Know	2(1.4%)
Method of Water Treatment	
Filter	17(77.3%)
Boil	2(9.1%)
Chemical	2(9.1%)
Don't Know	1(4.5%)
Frequency of Bathing in River	
Never	93(67.4%)
Sometimes	41(29.7%)
Always	3(2.2%)
NA/Don't Know	1(0.7%)
Frequency of Washing Clothes in River	
Never	80(58%)
Sometimes	49(35.5%)
Always	9(6.5%)
Frequency of Defecating in Open Field	
Never	110(79.7%)
Sometimes	27(19.6%)
Always	0(0%)
NA/Don't Know	1(0.7%)
Use of Toilet Paper After Defecating	
Never	13(9.4%)
Sometimes	92(66.7%)
Always	30(21.7%)
NA/Don't Know	3(2.2%)

Environmental Factors	Overall: N = 138
Washing Hands After Using Toilet	
Never	2(1.4%)
Sometimes	67(48.6%)
Always	69(50%)
Method of Washing Hands After Toilet	
Water	29(21%)
Soap and Water	108(78.3%)
NA/Don't Know	1(0.7%)
Frequency of Using Soap After Toilet	
Never	1(0.7%)
Sometimes	104(75.4%)
Always	23(16.7%)
NA/Don't Know	10(7.2%)
Washing Hands Before Eating	
Never	1(0.7%)
Sometimes	21(15.2%)
Always	115(83.3%)
NA/Don't Know	1(0.7%)
Method of Washing Hands Before Eating	
Water	19(13.8%)
Soap and Water	117(84.8%)
NA/Don't Know	2(1.4%)
Frequency of Using Soap Before Eating	
Never	0(0%)
Sometimes	87(63%)
Always	43(31.2%)
NA/Don't Know	8(5.8%)
Eating Soil	
Never	124(89.9%)
Sometimes	13(9.4%)
Always	0(0%)
NA/Don't Know	1(0.7%)
Washing Fruit	
Never	48(34.8%)
Sometimes	81(58.7%)
Always	8(5.8%)
NA/Don't Know	1(0.7%)
Favorite Fruit	
Mango	53(38.4%)
Banana	15(10.9%)
Avocado	12(8.7%)
Apple	7(5.1%)
Pineapple	6(4.3%)
Other	45(32.6%)
Eating Raw or Under-cooked Vegetables	
Never	1(0.7%)
Sometimes	98(71%)
Always	37(26.8%)
NA/Don't Know	2(1.4%)
Frequency of Washing Vegetables	
Never	60(43.5%)
Sometimes	44(31.9%)
Always	7(5.1%)
NA/Don't Know	27(19.6%)

Environmental Factors	Overall: N = 138
Walking Barefoot	
Never	7(5.1%)
Sometimes	61(44.2%)
Always	69(50%)
NA/Don't Know	1(0.7%)
Preferred Footwear at Home	
Barefoot	59(42.8%)
Sandals	74(53.6%)
Shoes	3(2.2%)
NA/Don't Know	2(1.4%)