Poisson Regression Article

1 Background

Figure 1: Siméon Denis Poisson. Source: Wikipedia



1.1 Understanding the Technique

1.1.1 The Poisson Distribution

Poisson regression extends the linear least-squares regression model to poisson distributed data. Often used to represent physical events occurring, the poisson distribution measures the probability of X events occuring over a given interval of time, and the average rate of those events occuring is λ. For example, the probability of $X = 6$ hurricanes happening over a year, when the historic average is $\lambda = \frac{4\,\text{hurricanes}}{\text{year}}$.

The poisson distribution can be viewed as a special case of the binomial distribution, which models the number of successes in a fixed number of trials (Oxford College of Emory University, Math Center, n.d.):

$$P(X = x) = f_x(x) = {}_nC_x p^x q^{n-x}$$

Where n is the number of trials, and x is the number of successes.

For the binomial distribution, the mean/expected value is np, which we'll call λ. So,

$$p = \frac{\lambda}{n}$$

If q is the probability of a failure, $q = 1 - p$, or

$$q = 1 - \frac{\lambda}{n}$$

Plugging back into our original equation:

$$\begin{aligned} P(X = x) = f_x(x) &= {}_nC_x(\frac{\lambda}{n})^x(1 - \frac{\lambda}{n})^{n-x} \\ &= \frac{n(n-1)(n-2)\cdots(n-x+1)}{x!} \cdot \frac{\lambda^x}{n^x}(1 - \frac{\lambda}{n})^{n-x} \\ &= \frac{n}{n} \cdot \frac{n-1}{n} \cdots \frac{n-x+1}{n} \cdot \frac{\lambda^x}{x!}(1 - \frac{\lambda}{n})^{n-x} \\ &= \frac{n}{n} \cdot \frac{n-1}{n} \cdots \frac{n-x+1}{n} \cdot \frac{\lambda^x}{x!}(1 - \frac{\lambda}{n})^{n}(1 - \frac{\lambda}{n})^{-x} \end{aligned}$$

Note that the second to last expression is an expression for $e^{-\lambda}$.

If we take the limit as our number of trials n approaches infinity while keeping x and λ fixed, all the terms except the second and third to last trend approach 1:

$$\lim_{n\to\infty} P(X = x) = f_x(x) = \frac{e^{-\lambda}\lambda^x}{x!}$$

This is the exact probability of x events occuring over λ time.

We can plot the probability mass function (PMF) to see the probability of x events occuring for every X, noting how different λ values change that distribution:

remark: notice how as λ increases the distribution is more bell curved shaped.

The poisson distribution is particularly well behaved because of it's distributions properties. It's first two moments are given (Soch and The Book of Statistical Proofs contributors 2020):


Expected Value / Mean:

$$\begin{aligned} E(x) &= \sum_{x\in\mathcal{X}} x \cdot f_X(x) \\ &= \sum_{x=0}^{\infty} x \cdot \frac{\lambda^x e^{-\lambda}}{x!} \\ &= \sum_{x=1}^{\infty} x \cdot \frac{\lambda^x e^{-\lambda}}{x!} \\ &= e^{-\lambda} \cdot \sum_{x=1}^{\infty} \frac{x}{x!}\lambda^x \\ &= \lambda e^{-\lambda} \cdot \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} \end{aligned}$$

Let $z = x - 1 \implies x = z + 1$:

$$E(X) = \lambda e^{-\lambda} \cdot \sum_{z=0}^{\infty} \frac{\lambda^z}{z!}$$

Figure 2: Poisson Distribution Examples



Remember that the power series expansion of $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$.

Finally,

$$\begin{aligned} E(X) =& \lambda e^{-\lambda} \cdot e^{\lambda} \\ =& \lambda \end{aligned}$$

Variance:

$$Var(X) = E(X^2) - E(X)^2$$

If we look at the expectation of $X(X-1)$:

$$\begin{aligned} E[X(X-1)] &= \sum_{x \in \mathcal{X}} x(x-1) \cdot f_X(x) \\ &= \sum_{x=0}^{\infty} x(x-1) \cdot \frac{\lambda^x e^{-\lambda}}{x!} \\ &= \sum_{x=2}^{\infty} x(x-1) \cdot \frac{\lambda^x e^{-\lambda}}{x!} \\ &= e^{-\lambda} \cdot \sum_{x=2}^{\infty} x(x-1) \cdot \frac{\lambda^x}{x \cdot (x-1) \cdot (x-2)!} \\ &= \lambda^2 \cdot e^{-\lambda} \cdot \sum_{x=2}^{\infty} \frac{\lambda^{x-2}}{(x-2)!}. \end{aligned}$$

let $z = x - 2 \implies x = z + 2$, hence:

$$E[X(X-1)] = \lambda^2 \cdot e^{-\lambda} \cdot \sum_{z=0}^{\infty} \frac{\lambda^z}{z!}$$

recall the power series expansion of $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$:

$$E[X(X-1)] = \lambda^2 \cdot e^{-\lambda} \cdot e^{\lambda} = \lambda^2$$

Since,

$$E[X(X-1)] = E(X^2 - X) = E(X^2) - E(X)$$

now:

$$E(X^2) - E(X) = \lambda^2 \implies E(X^2) = \lambda^2 + \lambda$$

Finally, by plugging in:

$$\begin{aligned} Var(X) =& E(X^2) - E(X)^2 \\ =& \lambda^2 + \lambda - \lambda^2 \\ =& \lambda \end{aligned}$$

We can verify this numerically. Here I simulate 10,000 poisson distributed datasets with $\lambda = 1$ and sample size $n = 25$:

```
set.seed(13)
sims = 10000 #Simulations
lambda = 1  #Frequency Parameter
n = 25      #Sample Size

means <- rep(NA, sims) #Place to store means
vars <- rep(NA, sims)  #Place to store variances

for(i in 1:sims){
  poisson_dat <- rpois(n, lambda) #Simulate random data
  means[i] <- mean(poisson_dat)   #Store the mean of the data
  vars[i] <- var(poisson_dat)     #Store the variance of the data
}
```

The means and variances should both be centered around 1:

Even with some outliers here and there, it checks out.

1.1.2 Poisson Regression

Where the typical linear regression goes as follows:


$$Y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \cdots + \beta_{p-1} x_{(p-1)i} + \epsilon_i$$

Since Y_i is assumed to be continuous, both sides of the equation can range from $(-\infty, \infty)$. To model a variable that is based on counts, and therefore strictly > 0, we have to transform the data to sit in the $(-\infty, \infty)$ range, and this can be achieved with a log transformation:

$$log(\lambda_i) = \beta_0 + \beta_1 x_{1i} + \cdots + \beta_{p-1} x_{(p-1)i}$$

where λ_i is the parameter for the Poisson distribution from which the ith response is drawn.

Figure 3: Poisson Means and Variances



If all of the data is not measured over the same unit of time, an offset variable is added to the equation:

$$\log(\lambda_i) = (\log(T_i) + \beta_0) + \beta_1 x_{1i} + \cdots + \beta_{p-1} x_{(p-1)i}$$

Lastly, a critical assumption of poisson regression is that the data behaves like it's poisson distributed, with equal mean and variance (Figure 3). Otherwise, another type of regression (negative binomial, quasi-poisson, etc.) should be used.

1.2 The Study



Jie et al. look to further investigate the Janzen-Connell effect, which looks to explain the mechanism behind biodiversity, or why a single tree or plant species doesn't just dominate a forest. In order for a plant to survive, it needs to not be eaten or destroyed by natural enemies (insects, animals, fungi, etc.). A common explanation behind biodiversity says that natural enemies specialize in eating/killing a specific plant over time, and those natural enemies accumulate in areas where there's high densities of those specific plants. Because of this, a new seedling might actually have an easier time in an area without a lot of members of its own species because those natural enemies have not accumulated so heavily. Essentially, there's always a push-and-pull between the plants and enemies in an area, and they balance each other out. (Jia et al. 2020).

For one of their analyses, the researchers here look to test factors that can affect a seedlings ability to survive around a lot of members of its own species (a.k.a recruitment):

- How does the density of a plant's own species affect its seedlings' recruitment?
- Which natural enemies drive seedling recruitment?


To control which natural enemies are present, a fence was used to keep out herbivores, and insecticides/fungicides were used to discourage insects/fungus.

Figure 4: Study Animals (Jia et al.)



To set up the experiment the researchers took 3 plots from a forest in China and divided them up into subplots:

Figure 5: Study Design Layout (Jia et al.)



Within each plot, half was fenced, and within each half, 10 squares were control, 10 had insecticide, and 10 had fungicide.

1.3 Goal of Statistical Analysis

For their regression equation, they regress seedling recruitment with fixed factors:

- Pesticide treatment
- Fencing exclosure
- Density of adult plants (same species)
- Density of adult plants (different species)
- Plot
- Census Date

Including a three-way interaction and pairwise two-way interactions between the first three.

And random factors:

- Subplot (random intercept)
- Species (random intercept)
- Effect of census on subplots (random effect)

Also, same species density has been log transformed and Z-scored

In *R*, our formula call looks like this:

```
formula <- recruits ~ pesticide*exclosure*con_crowding + het_crowding + plot + census  + (1| subplot) + (1|
```

2 Reproduction of the Study

2.1 Data Cleaning

I did some renaming of variables to fit my personal liking and some decoding for interpretation ease.

```
dat <- read_csv('seeds.csv')


dat$plot <- as.character(dat$plot)
dat$exclosure <- ifelse(dat$exclosure == 1, "Fenced", "Open")
dat$exclosure <- factor(dat$exclosure, levels = c("Open", "Fenced"))

dat <- dat |>
  mutate(pesticide = case_when(pesticide == "W" ~ "Control",
                               pesticide == "I" ~ "Insecticide",
                               pesticide == "F" ~ "Fungicide"))
```

2.2 Data Summary

Let's take a quick summary of our recruitment levels across treatment and exclosure:

Figure 6: Treatment and Exclosure Summary



Not a whole lot we can tell from this so far.

Here's how our recruitment levels differ by same species density:

Not really any conclusive evidence for how same species density affects seedling recruitment based on this alone.

For a frankenstein plot combining all of these:

Fun, but not super helpful.

As an aside, we can see how our data maps onto a poisson distribution:

Figure 7: Density Summary




Figure 8: Combined Summary





Figure 9: Poisson Distribution Mapped Onto Data



Our data appears more right-skewed than an assumed poisson distribution with mean recruits $\lambda = 3.6$. Let's see if this affects the assumptions of our model.

2.3 Model Assumptions

2.3.1 Gaussian Residuals

Since our model is a generalized linear mixed effect model, we need to check the assumption that our random effects are (multivariate) gaussian.

First we'll check the random intercepts for species:

Even with a small sample ($n = 16$ species), our random intercepts look roughly gaussian and follow the qq-plot pretty well.

Next, the random intercepts for subplots:

Even though the shapiro-wilks test indicates a deviation from normality, the histogram shows a roughly gaussian shape of our random intercepts for subplot centered at zero. The qq-plot looks a little wonky towards the right tail, but I don't think it's a huge problem.

Lastly, we check the multivariate normality of the effect of census on subplots:

The multivariate normality of the effects of census on subplots follows closely to a χ^2 distribution by fitting the qq-plot closely.

2.3.2 Dispersion

Dispertion ratios aren't quite as straightforward for mixed models, so we'll be using this approximation function provided by Dr. Ben Bolker, a mathematical statistician at McMaster University (Bolker et al. 2025).

Our dispersion ratio here is very good. The bottom left of the cluster of points highlights the high amount of 1's observed in the recruit measurement. Still, this is promising for our model.

Figure 10: Random intercepts of Species




Figure 11: Random intercepts of Subplots




Figure 12: Mahalanobis Distances of Census Effects on Subplots





Figure 13: Randomized Quantile Residuals of Poisson Model



2.3.3 Multicollinearity

Table 1: Model VIF

Term	VIF	Lower CI	Upper CI
Pesticide Treatment	4.353	3.919	4.851
Exclosure	3.287	2.974	3.649
Same Species Density	2.942	2.668	3.259
Different Species Density	1.241	1.167	1.349
Plot	1.137	1.078	1.240
Census Date	1.010	1.000	6.656
Pesticide Treatment: Exclosure	9.388	8.382	10.531
Pesticide Treatment: Same Species Density	7.227	6.467	8.093
Exclosure : Same Species Crowding	4.431	3.988	4.939
Pesticide Treatment : Exclosure : Same Species Crowding	10.772	9.609	12.092

The collinearity for the model looks good, with only some slight colinearity for pesticide treatment, exclosure, and same species density. VIF values for interaction terms are usually high, so there's not really anything to worry about here.

2.3.4 Sufficient Sample Size for Predictors

Ideally we should have at least 10 events counted per predictor:

$$\sum_{i=1}^{n=1019} \lambda_i = 3666 \geq 10p = 10(10) = 100 \quad \checkmark$$

2.4 Model Fit

We can obtain an overall model fit:

Table 2: Model Fit

AIC	AICc	BIC	R2_conditional	R2_marginal	ICC	RMSE	Sigma	Score_log	Score_spherical
4574.138	4575.065	4677.596	0.468	0.108	0.404	2.331	1	-2.002	0.027

Our fixed effects explain 10.8% of the variance in seedling recruitment. The total model including the random effects explains 46.8% of the variance in seedling recruitment. With our ICC, our random effects explain 40.4% of the variance in seedling recruitment.

The researchers mention testing the model fit when they drop some of the interaction terms. Since a three-way interaction can be very tricky to interpret, I wanted to test the fit between our normal model and a model with the three-way interaction term dropped in favor of just the pairwise two-way interactions.

Table 3: Model Comparison

	# Parameters	AIC	BIC	log Likelihood	Deviance	Chi-Sq	df	p
Dropped Interaction Model	19	4572.236	4665.841	-2267.118	4534.236	NA	NA	NA
Original Model	21	4574.138	4677.596	-2266.069	4532.138	2.097	2	0.35

Really not much difference between the two models, so begrudgingly, we'll move forward with the three-way interaction.

2.5 Model Inference

Now to the heart of the analysis:

- Do the fenced in plots have more seedlings?
- Does the pesticide treatment affect the number of seedlings?
- Is there an interactive effect between pesticide, exclosure, and same species crowding?

A general summary our model gives us:

Table 4: Model Summary

	Esti-mate	SE	Z	p	significance	Percent Change
Intercept	0.837	0.161	5.209	<0.001	***	
Pesticide = Fungicide	0.070	0.084	0.830	0.407		7.3
Pesticide = Insecticide	-0.146	0.089	-1.646	0.1		-13.6
Fenced	-0.167	0.089	-1.879	0.06	.	-15.4
Same Species Density	0.023	0.053	0.429	0.668		2.3
Different Species Density	-0.028	0.030	-0.938	0.348		-2.8
Plot 2	0.184	0.063	2.924	0.003	**	20.2
Plot 3	-0.159	0.067	-2.376	0.018	*	-14.7
Census Date = 2017 Spring	-0.400	0.053	-7.501	<0.001	***	-33
Pesticide = Fungicide : Fenced	0.178	0.120	1.481	0.139		19.5
Pesticide = Insecticide : Fenced	0.273	0.125	2.182	0.029	*	31.4
Pesticide = Fungicide : Same Species Density	0.178	0.064	2.804	0.005	**	19.5
Pesticide = Insecticide : Same Species Density	0.039	0.067	0.586	0.558		4
Fenced : Same Species Density	-0.082	0.066	-1.244	0.214		-7.9
Pesticide = Fungicide : Fenced : Same Species Density	0.104	0.093	1.111	0.266		10.9
Pesticide = Insecticide : Fenced : Same Species Crowding	0.131	0.095	1.376	0.169		14

where the percent change = $(e^{\text{estimate}} - 1) \times 100$.

Our model says, for the non-interaction fixed effects (deep breath):

- For the intercept, we'd expect to find $e^{0.837} = 3.598$ seedlings on a control subplot, that's open, has a same species & different species density of 0, on plot 1, and in the spring of 2016.
- For every standard deviation increase in different species density, we'd expect to see 2.8% fewer seedlings
- We'd expect to see 20.2% more seedlings on plot 2 than plot 1.
- We'd expect to 14.7% fewer seedlings on plot 3 than plot 1.
- We'd expect to see 33% fewer seedlings in the spring of 2017 compared to the spring of 2016

For the statistical discernability of our terms:

- There are marginally fewer seedlings in the fenced plots than the non-fenced plots ($Z = -1.879, p = 0.06$)
- There are discernibly more seedlings on plot 2 than plot 1 ($Z = 2.924, p = 0.003$)
- There are discernibly fewer seedlings on plot 3 than plot 1 ($Z = -2.376, p = 0.018$)
- There are discernibly fewer seedslings in the spring of 2017 than the spring of 2016 ($Z = -7.501, p < 0.001$)
- The two-way interaction between insecticide and fenced plots is statistically discernible ($Z = 2.182, p = 0.029$)
- The two-way interaction between fungicide and same species density is statistically discernible ($Z = 2.804, p = 0.005$)

Testing the effect sizes of our fixed effects (Jaeger 2025):

Table 5: Model Effect Sizes

Effect	R-Squared	Upper CI	Lower CI
Model	0.077	0.06	0.122
Pesticide	0.003	<0.001	0.015
Exclosure	0.003	<0.001	0.014
Census	0.04	0.02	0.067
Plot	0.017	0.006	0.038
Same Species Density	<0.001	<0.001	0.005
Different Species Density	<0.001	<0.001	0.005
Pesticide : Exclosure	0.003	<0.001	0.016
Pesticide : Same Species Density	0.004	<0.001	0.017
Exclosure : Same Species Density	<0.001	<0.001	0.007
Pesticide : Exclosure : Same Species Density	<0.001	<0.001	0.010

The largest effect of any term, besides the model, is the census term, which explains 4% of the variance in seed recruitments.

Now let's try to untangle all those interactions.

2.6 Model Interactions

2.6.1 Same Species Density Trend

First, we'll see if the density of a plant's own species affects its seedlings' recruitment through estimated marginal trends:

Table 6: Same Species Density Estimated Marginal Trend

Same Species Density Trend	SE	Z	p	Percent Change	Lower CI	Upper CI
0.093	0.033	2.85	0.004	9.769	2.954	17.035

Let's see the predicted number of seedlings across the same species density:

Interestingly, the overall effect of same species density is to positively affect the number of seedling recruits expected. Our model predicts that for every standard deviation increase in same species density, the number of seedling recruits increases by 9.8% (95% CI: [3.0, 17.0]). This trend is statistically disernible ($Z = 2.85, p = 0.004$).

2.6.2 Effect of Fencing in Plots

Next, we'll see if fencing off the plots from herbivores increases the number of seedlings:

Averaging across other variables, we see a contrast between the fenced-in and non-fenced-in plots:

Table 7: Contrast of Exclosure Types

Contrast	Estimate	SE	Z	p	Percent Change	Upper CI	Lower CI
Open - Fenced	0.016	0.052	0.313	0.754	1.64	12.526	-8.193

Plotting the predicted number of seedlings by exclosure type:

Our model predicts that an open plot will have 1.6% more seedlings than a fenced in plot (95% CI: [-8.2, 8.2]). As we can see from the table and plot, the global effect of exclosure type is not statistically discernible ($Z = 0.313, p = 0.754$).

Figure 14: Same Species Estimated Marginal Trend





Figure 15: Exclosure Type Estimated Marginal Means



2.6.3 Effect of Pesticides


We've tried to see the effects of one kind of natural threat to seedlings, herbivores, now we'll see if pesticides, thought to discourage insects and fungi, affects the number of seedlings:

Table 8: Contrast of Pesticide Types

Contrast	Estimate	SE	Z	p	Adjusted-p	Percent Change	Lower CI	Upper CI
Control - Fungicide	-0.159	0.060	-2.634	0.023	0.034	-14.730	-26.001	-1.742
Control - Insecticide	0.009	0.063	0.144	0.989	0.989	0.906	-12.858	16.845
Fungicide - Insecticide	0.168	0.061	2.766	0.016	0.034	18.338	2.605	36.483

Plotting the predicted number of seedlings by pesticide type:

Figure 16: Pesticide Type Estimated Marginal Means



The global effect of fungicide is statistically more favorable to seedlings than insecticide ($Z = -2.634, p = 0.023$, adjusted-$p = 0.034$) and the control ($Z = 2.766, p = 0.016$, adjusted-$p = 0.034$). Our model predicts that the fungicide plots will have 14.7% more seedlings (95% CI: [1.7, 26.0]) than the control plots, and 18.3% more seedlings (95% CI: [2.6, 36.5]) than the insecticide plots.

2.6.4 Pesticide Type and Exclosure Type Interaction


Based on a significant interaction term between fungicide and the fenced plots, we'll probe into that interaction here:

Table 9: Contrast of Insecticide Types by Exclosure Types

Contrast	Exclosure Type	Esti-mate	SE	Z	p	Adjusted-p	Percent Change	Lower CI
Control - Fungicide	Open	-0.070	0.084	-0.830	0.684	0.684	-6.771	-23.518
Control - Insecticide	Open	0.146	0.089	1.646	0.226	0.401	15.689	-5.986
Fungicide - Insecticide	Open	0.216	0.087	2.483	0.035	0.104	24.091	1.219
Control - Fungicide	Fenced	-0.249	0.086	-2.882	0.011	0.066	-22.009	-36.284
Control - Insecticide	Fenced	-0.128	0.089	-1.442	0.319	0.401	-11.987	-28.484
Fungicide - Insecticide	Fenced	0.121	0.086	1.413	0.334	0.401	12.850	-7.653

Plotting the predicted number of seedlings by exclosure and pesticide type:

Figure 17: Pesticide Type by Exclosure Type Estimated Marginal Means



Even though the interaction term in the overall model is significant, none of the contrasts remain statistically discernible after adjustment. It's interesting to see how the relationship between each pesticide type changes between fenced and non-fenced in plots.

2.6.5 Pesticide Type and Same Species Density Interaction


For our last interaction, we'll see if the effect of the same species density changes depending on the pesticide type used:

Table 10: Estimated Marginal Trend of Same Species Density by Pesticide Type

Pesticide Type	Same Species Density Trend	SE	Z	p	adjusted-p	Percent Change	Lower CI	Upper CI
Control	-0.018	0.042	-0.437	0.662	0.662	-1.825	-9.607	6.628
Fungicide	0.212	0.042	5.002	<0.001	<0.001	23.603	13.755	34.303
Insecticide	0.086	0.043	1.991	0.046	0.07	8.996	0.135	18.642

Visualizing those trends:

Figure 18: Estimated Marginal Trends of Same Species Density by Pesticide Type



Testing the differences in those trends:

Table 11: Contrasts of Estimated Marginal Trend of Same Species Density by Pesticide Type

Contrast	Estimate	SE	Z	p	adjusted-p	Percent Change	Lower CI	Upper CI
Control - Fungicide	-0.230	0.047	-4.937	<0.001	<0.001	-20.572	-28.800	-11.394
Control - Insecticide	-0.105	0.047	-2.203	0.071	0.071	-9.928	-19.410	0.670
Fungicide - Insecticide	0.126	0.048	2.639	0.023	0.034	13.401	1.417	26.801

We can see:

- The trend for same species density is discernibly greater for the fungicide treatment versus the control ($Z = 4.937, p < 0.001$, adjusted-$p < 0.001$). Our model predicts that for every standard deviation increase in same species density, the percent increase in seedlings is 20.572 greater (95% CI: [11.394, 28.800]) for the fungicide group than the control group.
- The trend for same species density is marginally greater for the insecticide treatment versus the control ($Z = 2.203, p = 0.071$, adjusted-$p = 0.071$). Our model predicts that for every standard deviation increase in

same species density, the percent increase in seedlings is 9.928 greater (95% CI: [-0.670, 19.410]) for the fungicide group than the control group.

- Lastly, the trend for same species density is discernibly greater for the fungicide treatment versus the insecticide treatment ($Z = 2.639, p = 0.023$, adjusted-$p = 0.034$). Our model predicts that for every standard deviation increase in same species density, the percent increase in seedlings is 13.401 greater (95% CI: [1.417, 26.801]) for the fungicide group than the insecticide group.

2.7 Conclusions

If we harken back to the study background, the Janzen-Connell effect tells us that we should expect new seedlings to struggle propagating in an area that is already densely populated with its own species, as there is a higher chance that natural enemies have begun to accumulate there. What we found through our data doesn't necessarily support this in every case however. Looking at the global trend of same species density, seedlings tends to be better off when there is a higher density of their own species Figure 14, though this is only true for groups with insecticide and fungicide Table 10. The trend for the control group isn't statistically discernible from zero, so we can't say for sure which direction this trend goes Table 10. However, if we look at the hypothesis generously, we might say that the groups being treated with pesticides, especially fungicide Table 11 would have fewer natural enemies to fight off, and in that case a higher same species density would likely just indicate that that species happens to do well there, and it makes sense for more seedlings to appear. Lastly when discussing the same species trend, the three-way interaction between exclosure type, pesticide, and same species density did not indicate that any of those trends differed by whether they were enclosed or not Table 4.

When investigating which natural enemies have the strongest effect on seedling recruitment, the global trend indicates that the fungicide had the most noticeable effect on recruitment, being noticeably greater than the control and insecticide groups Table 8. Across exclosure types, the effect of pesticides and being in the control groups shifted heavily Table 9. While little change was noticed in the fungicide group, being fenced in helped the insecticide group but hurt the control group Figure 17. The fungicide group was only better off compared to the insecticide group when the exclosure was open, but that effect reversed compared to the control when the exclosure was fenced Table 9. For the global effect of exclosure type, there was no statistically discernible difference between the open and fenced groups Table 7.

3 Disagreements With the Article

3.1 Results

3.1.1 Pesticide and Same Species Density Interaction

For the interaction between fungicide and same species density, I agree partially with the article when they say:

> "Protection from fungi increased … recruitment … of seedlings when conspecific adult densities were high (Fungicide × conspecific adult density interactions, recruitment: Poisson GLMM, $z = 4.87, P < 0.001$…)"

The interaction in my model summary wasn't quite as strong as there's though I still found it statistically discernible Table 4. Their wording makes me think they looked at the effect of the pesticides and varying levels of same species density, which I reversed, looking at the effect of same species density on the pesticides. Still, I'm assuming this value comes from their model summary, which should be the same as mine. Also, I've only heard of staggering levels of a continuous variable in an estimated marginal means/trends setting, not for a standard GLMM summary, but they don't mention either technique in the manuscript or supplemental materials.


For the interaction between insecticide and same species density, I don't get the same result as the article when they say:

> "Suppression of insects also increased seedling recruitment at high conspecific adult densities (Poisson GLMM: Insecticide × conspecific adult density interaction, $z = 2.19, P = 0.029$)…"

In my model summary, the interaction between same species density and the insecticide treatment was not statistically discernible from zero, and it wasn't particularly close Table 4.

If we compare their figure looking at the interaction between same species density and pesticide to mine Figure 18:

Figure 19: Jie et al.



So while the authors use the GLMM predictions and each dot is the mean and SE of the observed values at that density level, I opted for the estimated marginal trends, as I think that does a nicer job of untangling those interactions while making the trends visually understandable. Their plot also has a much narrower range of x values compared to mine, so you're seeing a much more truncated effect. I notice that their trends for the insecticide and fungicide groups have a logarithmic shape, whereas mine look more linear and quadratic for both groups respectively.

I'll try to recreate what I believe their analysis is doing:

Table 12: Contrasts of Estimated Marginal Means of Pesticide Type by Same Species Density (Post-hoc)

Contrast	Same Species Density (Z)	Esti-mate	SE	Z	p	adjusted-p	Percent Change	Lower CI	Upper CI
Control - Fungicide	-1	0.071	0.083	0.851	0.671	0.863	7.356	-11.705	30.531
Control - Insecticide	-1	0.114	0.084	1.346	0.37	0.573	12.029	-8.077	36.532
Fungicide - Insecticide	-1	0.043	0.085	0.500	0.871	0.98	4.353	-14.539	27.422
Control - Fungicide	0	-0.159	0.060	-2.634	0.023	0.052	-14.730	-26.001	-1.742
Control - Insecticide	0	0.009	0.063	0.144	0.989	0.989	0.906	-12.858	16.845
Fungicide - Insecticide	0	0.168	0.061	2.766	0.016	0.047	18.338	2.605	36.483
Control - Fungicide	1	-0.390	0.069	-5.674	<0.001	<0.001	-32.272	-42.341	-20.445
Control - Insecticide	1	-0.096	0.072	-1.323	0.382	0.573	-9.112	-23.261	7.647
Fungicide - Insecticide	1	0.294	0.068	4.295	<0.001	<0.001	34.196	14.297	57.559

This clears a few things up. I should mention however, I don't see in the article or the supplementary materials where they specify how they're doing this calculation, so I don't know for sure they're using the same cutoffs for low, medium, and high levels of same species density. We do see now that there are discernibly more seedlings in the fungicide

group compared to the control at 1 standard deviation above average for same species density ($Z = 5.674, p < 0.001$, adjusted-$p < 0.001$). We'd expect to see 32.272% more seedlings in the fungicide group compared to the control group (95% CI: [20.445, 42.341]) Table 12.

To their second claim, there is a clear disagreement with the article. At one standard deviation above average for same species density, there is no discernible difference between the control group and the insecticide group ($Z = 1.323, p = 0.382$, adjusted-$p = 0.573$). We'd expect to see the insecticide group have 9.112% more seedlings than the control group (95% CI: [-7.647, 23.261]) Table 12.

3.1.2 Effect of Fencing in Plots

For the effect of fencing in the plots, I come to a similar conclusion but obtained different numbers:

> "The effect of large vertebrates was not estimated to be large in any of the models (Recruitment: Poisson GLMM, $z = -0.23, P = 818$…)"

I'm assuming that $P = 818$ is a typo. Either way, in my model summary, the coefficient estimate for fencing in the enclosures was marginally discernible, with very different statistics Table 4. If we assume they're talking about the estimated marginal means effect that isolates that variable, my methods also found a non-discernible effect of the enclosures, but with different statistics once again Table 7.

3.2 Discussion

The first sentence of the discussion reads:

> "The deleterious effects of natural enemies on seedling recruitment and survival, assessed at the community scale, increase with the local density of conspecific adults."

I'd say this is only true for fungicide. Insecticide is not discernibly better than the control for seedlings at any level of same species density Table 7. They seem to change their wording slightly later in the paragraph, so I'm not sure why they it isn't consistent with the first sentence:

> "Thus, our results suggest that plant-associated fungi suppress the recruitment of tree species dominating the nearby canopy, which will increase turnover of tree species and could increase diversity."

Once again, they repeat:

> "While the evidence that natural enemies reduce seedling performance at high conspecific density in our system is in line with previous pesticide application experiments…"

This is only true of the fungicide group. I also tested the effect of the exclosure at different levels of same species density:

Table 13: Contrasts of Estimated Marginal Means of Exclosure Type by Same Species Density (Post-hoc)

Contrast	Same Species Density (Z)	Esti-mate	SE	Z	p	adjusted-p	Percent Change	Lower CI	Upper CI
Open - Fenced	-1	0.012	0.071	0.168	0.867	0.867	1.203	-12.012	16.403
Open - Fenced	0	0.016	0.052	0.313	0.754	0.867	1.640	-8.193	12.526
Open - Fenced	1	0.021	0.059	0.350	0.726	0.867	2.078	-9.012	14.520

As we can see, the effect of exclosure type, ostensibly meant to correspond with natural herbivores, is not discernible from zero at any level of same species density Table 13.

They repeat the same claim I think is too broad later on as well:

> "Instead, our results suggest that density of adult conspecifics is a better predictor of natural enemy impacts on recruitment and survival of seedlings."

3.2.1 Effect of Same Species Density on Unmanipulated Plots

Later in the discussion, they make the claim:

> "These natural enemy impacts appear to be important, even though the net effect of conspecific density is only weakly negative in plots when natural enemies are unmanipulated."

I calculated the effects of same species density at various groups of pesticide and exclosure types:

Table 14: Estimated Marginal Trends of Same Species Density by Pesticide and Exclosure (Post-hoc)

Pesticide	Exclosure	Same Species Density Trend	SE	Z	p	adjusted-p	Percent Change	Lower CI	Upper CI
Control	Open	0.023	0.053	0.429	0.668	0.668	2.305	-7.806	13.524
Fungicide	Open	0.201	0.054	3.751	<0.001	<0.001	22.297	10.088	35.859
Insecticide	Open	0.062	0.058	1.072	0.284	0.34	6.398	-5.008	19.174
Control	Fenced	-0.060	0.054	-1.102	0.271	0.34	-5.788	-15.268	4.753
Fungicide	Fenced	0.223	0.054	4.108	<0.001	<0.001	24.923	12.338	38.918
Insecticide	Fenced	0.110	0.053	2.082	0.037	0.075	11.658	0.649	23.871

Contrary to their claim, the effect of same species density on the unmanipulated plots, a.k.a., the open exclosure and control pesticide group, is not discernibly different than zero ($Z = 0.429, p = 0.668$, adjusted-$p = 0.668$). Important to their claim however, the esimated trend is actually slightly *positive*, as we'd expect for an unmanipulated plot, for every standard deviation increase in same species density would result in 2.305% *more* seedlings (95% CI: [-7.806, 13.524]) Table 14.

When the authors claim,

> "Our results extend the experimental evidence that seedling performance is suppressed by adult conspecifics to a community scale."

This is simply not true for seedling recruitment. None of the groups across pesticide and exclosure types had a statistically discernible negative trend of same species density on seedling recruitmentTable 14.

Later on the authors again claim:

> "…on the basis of the evidence presented here, insects and plant-associated fungi appear to play a greater role in mediating conspecific density dependence than large herbivores."


I don't see how they can claim this when the effect of insecticide on seedling recruitment is not discernible at any level of same species density Table 12.

3.2.2 Diagnosing the Differences

To try and understand how these differences arose, I checked to see if their were any differences in our code, and I think I understand some of them. As for the first difference in the interaction between fungicide and same species density, the authors of the paper changed the glm() formula once they found out that the three-way interaction between pesticide, exclosure, and same species density was not significant. After changing the model to only a two-way interaction between pesticide and same species density, that interaction between fungicide and same species density was a lot stronger.

Here's their first model with very similar estimates to mine Table 4:

Figure 20: Jie et al.



```
### Is the three-way interaction among pesticide, A.con_lg_sd and exclosure significant?
mod.recr.int3 <- glmer(recruits ~ exclosure*pesticide*A.con_lg_sd + A.het_lg_sd + szcensus + szsite  +
                         (1+census||quad.unique) + (1|sp.),
                       data=pest.recr.dat, family=poisson,
                       control=glmerControl(optimizer="optimx",
                                            optCtrl=list(method=c('bobyqa', 'Nelder-Mead'))))
```

87:1 (Untitled)


Console | Terminal | Markers | Background Jobs

R 4.5.1 · C:/Users/12697/Downloads/

```
                                Estimate Std. Error z value Pr(>|z|)
(Intercept)                      0.64562    0.15079   4.282 1.86e-05 ***
exclosure1                      -0.16687    0.08907  -1.873  0.06101 .
pesticideF                       0.06993    0.08470   0.826  0.40901
pesticideI                      -0.14581    0.08837  -1.650  0.09892 .
A.con_lg_sd                      0.02281    0.05348   0.426  0.66975
A.het_lg_sd                     -0.02829    0.02943  -0.961  0.33657
szcensus1                        0.19993    0.02629   7.604 2.87e-14 ***
szsite1                         -0.00834    0.03679  -0.227  0.82067
szsite2                          0.17597    0.03524   4.993 5.95e-07 ***
exclosure1:pesticideF            0.17849    0.12087   1.477  0.13974
exclosure1:pesticideI            0.27336    0.12533   2.181  0.02917 *
exclosure1:A.con_lg_sd          -0.08244    0.06694  -1.231  0.21815
pesticideF:A.con_lg_sd           0.17847    0.06427   2.777  0.00549 **
pesticideI:A.con_lg_sd           0.03922    0.06768   0.579  0.56229
exclosure1:pesticideF:A.con_lg_sd  0.10361    0.09420   1.100  0.27136
exclosure1:pesticideI:A.con_lg_sd  0.13068    0.09595   1.362  0.17321
```

Here's the second that now makes the interaction appear more significant. Note they have another type, where their $Z = 4.817$ and in the paper they say $Z = 4.87$:

Figure 21: Jie et al.



```
### Then, do Exclosures affect seedling recruitment?
mod.recr.excl <- glmer(recruits ~ pesticide*A.con_lg_sd + A.het_lg_sd + exclosure + szcensus + szsite  +
                         (1+census||quad.unique) + (1|sp.),
                       data=pest.recr.dat, family=poisson,
                       control=glmerControl(optimizer="optimx",
                                            optCtrl=list(method=c('bobyqa', 'Nelder-Mead'))))
```

102:1 (Untitled)

Console | Terminal | Markers | Background Jobs

R 4.5.1 · C:/Users/12697/Downloads/

```
Fixed effects:
                     Estimate Std. Error z value Pr(>|z|)
(Intercept)          0.569800   0.146566   3.888 0.000101 ***
pesticideF           0.159966   0.061407   2.605 0.009187 **
pesticideI          -0.009277   0.063602  -0.146 0.884030
A.con_lg_sd         -0.016149   0.042437  -0.381 0.703544
A.het_lg_sd         -0.032471   0.029786  -1.090 0.275648
exclosure1          -0.011788   0.051306  -0.230 0.818277
szcensus1            0.199907   0.026347   7.587 3.26e-14 ***
szsite1             -0.008852   0.037009  -0.239 0.810968
szsite2              0.177752   0.035534   5.002 5.67e-07 ***
pesticideF:A.con_lg_sd  0.226736   0.047067   4.817 1.45e-06 ***
pesticideI:A.con_lg_sd  0.103572   0.047879   2.163 0.030526 *
---
```

Later on, because the effect of exclosure is not statistically significant, the authors recode the model to exclude that term:

Figure 22: Jie et al.

```
############################
### Not include the exclosure?
## Meanwhile, the effect of exclosure treatment is not to be large, so is not considered further in the analyses.

## How about not include the exclosure treatment in the model?
mod.recr <- glmer(recruits ~ pesticide*A.con_lg_sd + A.het_lg_sd + szcensus + szsite  +
                    (1+census||quad.unique) + (1|sp.),
                  data=pest.recr.dat, family=poisson,
                  control=glmerControl(optimizer="optimx",
                                       optCtrl=list(method=c('bobyqa', 'Nelder-Mead'))))
```

These kind of post-hoc statistical adjustments, I believe, are very antithetical to what statisticians are telling us about responsible research ethics. There's a reason that high standard journals, like Nature, are starting to require pre-approved methods before running any analysis on an experiment. Picking and choosing variables like this by testing different models is certain to bias the experiment. At least with an algorithmic selection approach like Lasso, the decisions are not being made subjectively by humans. I would at least expect the researchers to provide p-value adjustments to their coefficients given they've created several models. Not as important, but also frustrating is the fact that their code doesn't run cleanly from start to finish, and different scripts require objects defined in the other (or not at all), indicating that nobody thought to check that they run on a new session in *R*.

4 Acknowledgement

I would like to thank the researchers who published this paper: Shihong Jia, Xugao Wang, Zuoqiang Yuan, Fei Lin, Ji Ye, Guigang Lin, Zhanqing Hao & Robert Bagchi. Transparency in data and code availability is hugely important for research ethics, and it makes my project possible.

5 AI Use

All of the guidance given by generative AI for this article can be found in this conversation.

6 Statistical Packages Used

- *tidyverse* for… everything it's *tidyverse* (Wickham et al. 2019)
- *ggplot2* for plotting (Wickham 2016)
- *ggthemes* for theme_clean() (Arnold 2024)
- *patchwork* for combined plots (Pedersen 2025)
- *lmerTest* for inference of GLMER model (Kuznetsova et al. 2017)
- *r2glmm* for effect sizes (Jaeger 2025)
- *performance* for model fit statistics (Lüdecke et al. 2021)

References

Arnold, Jeffrey B. 2024. *Ggthemes: Extra Themes, Scales and Geoms for 'Ggplot2'*. https://CRAN.R-project.org/package=ggthemes.

Bolker, Ben et al. 2025. *GLMM FAQ*. Https://bbolker.github.io/mixedmodels-misc/glmmFAQ.html.

Jaeger, Byron. 2025. *R2glmm: Computes r Squared for Mixed (Multilevel) Models*. https://CRAN.R-project.org/package=r2glmm.

Jia, Shuai, Xugao Wang, Zuoqiang Yuan, et al. 2020. "Tree Species Traits Affect Which Natural Enemies Drive the Janzen-Connell Effect in a Temperate Forest." *Nature Communications* 11 (1): 286. https://doi.org/10.1038/s41467-019-14140-y.

Kuznetsova, Alexandra, Per B. Brockhoff, and Rune H. B. Christensen. 2017. "lmerTest Package: Tests in Linear Mixed Effects Models." *Journal of Statistical Software* 82 (13): 1–26. https://doi.org/10.18637/jss.v082.i13.

Lüdecke, Daniel, Mattan S. Ben-Shachar, Indrajeet Patil, Philip Waggoner, and Dominique Makowski. 2021. "performance: An R Package for Assessment, Comparison and Testing of Statistical Models." *Journal of Open Source Software* 6 (60): 3139. https://doi.org/10.21105/joss.03139.

Oxford College of Emory University, Math Center. n.d. *The Connection Between the Poisson and Binomial Distributions*. Https://mathcenter.oxford.emory.edu/site/math117/connectingPoissonAndBinomial/.

Pedersen, Thomas Lin. 2025. *Patchwork: The Composer of Plots*. https://CRAN.R-project.org/package=patchwork.

Soch, Joram, and The Book of Statistical Proofs contributors. 2020. *The Book of Statistical Proofs*. Https://statproofbook.github.io/. https://doi.org/10.5281/zenodo.4305949.

Wickham, Hadley. 2016. *Ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. https://ggplot2.tidyverse.org.

Wickham, Hadley, Mara Averick, Jennifer Bryan, et al. 2019. "Welcome to the tidyverse." *Journal of Open Source Software* 4 (43): 1686. https://doi.org/10.21105/joss.01686.