

Assignment 3: Derivation

Maxwell Bo 43926871

May 17, 2017

1. (a) n is a **value** parameter. m is a **result** parameter.
- (b) Our post-condition, $mr\text{run}(A, n_0, m)$, expands to $lr\text{run}(A, n_0, m) \wedge (m < A.\text{len} \Rightarrow A_{n_0} \neq A_m)$. This satisfies the form $Q \triangleq Q_1 \wedge Q_2$. Q_1 was chosen as the invariant. Thus,

$$inv \triangleq lr\text{run}(A, n_0, m)$$

(c) Let

$$\begin{aligned} pre &\triangleq lr\text{run}(A, n, n+1) \\ post &\triangleq mr\text{run}(A, n_0, m) \end{aligned}$$

s.t.

$$\begin{aligned} &n, m : [pre, post] \\ \sqsubseteq &\{ \text{Composition: middle predicate is } inv \} \\ &n, m : [pre, inv]; \quad n, m : [inv, post] \\ \sqsubseteq &\{ \text{Assignment: } pre \Rightarrow inv[m \setminus n + 1] \} \\ &m := n + 1; \quad n, m : [inv, post] \end{aligned}$$

\therefore

$$\begin{aligned} inv[m \setminus n + 1] &\equiv lr\text{run}(A, n_0, m)[m \setminus n + 1] \\ &\equiv lr\text{run}(A, n_0, n + 1) \end{aligned}$$

\therefore

$$lr\text{run}(A, n, n + 1) \Rightarrow lr\text{run}(A, n_0, n + 1)$$

Let

$$guard \triangleq (m < A.\text{len} \wedge A_{n_0} = A_m)$$

s.t.

$$\begin{aligned} \sqsubseteq &\{ \text{Strengthen post: } inv \wedge \neg guard \Rightarrow post \} \\ &m := n + 1; \quad n, m : [inv, inv \wedge \neg guard] \end{aligned}$$

\therefore

$$\begin{aligned} &inv \wedge \neg guard \Rightarrow post \\ \equiv &\{ \text{Expansion of definitions} \} \\ &lr\text{run}(A, n_0, m) \wedge \neg(m < A.\text{len} \wedge A_{n_0} = A_m) \Rightarrow mr\text{run}(A, n_0, m) \\ \equiv &\{ \text{Expansion of functions} \} \end{aligned}$$

$$\begin{aligned}
& lrun(A, n_0, m) \wedge \neg(m < A.len \wedge A_{n_0} = A_m) \Rightarrow lrun(A, n_0, m) \wedge (m < A.len \Rightarrow A_{n_0} \neq A_m) \\
\equiv & \{ \text{De Morgan's law - negation of conjunction} \} \\
& lrun(A, n_0, m) \wedge (\neg(m < A.len) \vee \neg(A_{n_0} = A_m)) \Rightarrow lrun(A, n_0, m) \wedge (m < A.len \Rightarrow A_{n_0} \neq A_m) \\
\equiv & \{ P \Rightarrow Q \equiv \neg P \vee Q \} \\
& lrun(A, n_0, m) \wedge (\neg(m < A.len) \vee \neg(A_{n_0} = A_m)) \Rightarrow lrun(A, n_0, m) \wedge (\neg(m < A.len) \vee (A_{n_0} \neq A_m)) \\
\equiv & \{ \} \\
& \text{true}
\end{aligned}$$

$$\begin{aligned}
& \sqsubseteq \{ \text{Repetition} \} \\
& \quad m := n + 1; \\
& \quad \mathbf{do} (m < A.len \wedge A_{n_0} = A_m) \rightarrow \\
& \quad \quad n, m : [inv \wedge guard, inv \wedge (0 \leq V < V_0)] \\
& \quad \mathbf{od}
\end{aligned}$$

where

$$V \triangleq A.len - m$$

$$\begin{aligned}
& \sqsubseteq \{ \text{Assignment: } inv \wedge \neg guard \Rightarrow (inv \wedge (0 \leq V < V_0))[m \setminus m + 1] \} \\
& \quad m := n + 1; \\
& \quad \mathbf{do} (m < A.len \wedge A_{n_0} = A_m) \rightarrow \\
& \quad \quad m := m + 1 \\
& \quad \mathbf{od}
\end{aligned}$$

\therefore

$$\begin{aligned}
(inv \wedge (0 \leq V < V_0))[m \setminus m + 1] & \equiv (lrun(A, n_0, m) \wedge (0 \leq (A.len - m) < (A.len - m_0)))[m, m_0 \setminus m + 1, m] \\
& \equiv lrun(A, n_0, m + 1) \wedge (0 \leq (A.len - (m + 1)) < (A.len - m))
\end{aligned}$$

\therefore

2.

$$\begin{aligned}
pre & \triangleq A.len > 0 \\
post & \triangleq mrun(A, \ell, h) \wedge (\forall p, q \cdot mrun(A, p, q) \Rightarrow (h - \ell) \leq (q - p))
\end{aligned}$$

$$\ell, h : [pre, post]$$

$$\begin{aligned}
& \sqsubseteq \{ \text{Composition: middle predicate is } inv \} \\
& \quad \ell, h : [pre, inv]; \quad \ell, h : [inv, post]
\end{aligned}$$

where

$$inv \triangleq mrun(A, \ell, h)$$

$$\begin{aligned}
& \ell := 0; \\
& \mathbf{do} (m < A.len \wedge A_{n_0} = A_m) \rightarrow \\
& \quad m := m + 1 \\
& \mathbf{od}
\end{aligned}$$