

# Robust Standard Errors in Small Samples

Michal Kolesár

September 20, 2019

The package `ShiftShareSE` implements confidence intervals proposed by Adão et al. [2019] for inference in shift-share least squares and instrumental variables regressions, in which the regressor of interest (or the instrument) has a shift-share structure, as in Bartik [1991]. A shift-share variable has the structure  $X_i = \sum_{s=1}^S w_{is} X_s$ , where  $i$  indexes regions,  $s$  indexes sectors,  $X_s$  are sectoral shifters (or shocks), and  $w_{is}$  are shares, such as initial share of region  $i$ 's employment in sector  $s$ .

This vignette illustrates the use of the package using a dataset from Autor et al. [2013] (ADH hereafter). The dataset is included in the package as the list `ADH`. The first element of the list, `ADH$reg` is a data-frame with regional variables, the second element, `ADH$sic` is a vector of SIC codes for the sectors, and `ADH$W` is a matrix of shares. See `?ADH` for a description of the dataset.

## Examples

We now replicate column (1) of Table V in Adão et al. [2019]. First we load the package, define the vector of controls, and define a vector of 3-digit SIC codes:

```
library("ShiftShareSE")
ctrls <- "t2 + l_shind_manuf_cbp + l_sh_popedu_c + l_sh_popfborn +
         l_sh_empl_f + l_sh_routine33 + l_task_outsource + division"
sic <- floor(ADH$sic/10)
```

We cluster the standard errors at the 3-digit SIC code (using the option `sector_cvar`), and, following ADH, weight the data using the weights `ADH$reg$weights`. See `?reg_ss` and `?ivreg_ss` for full description of the options.

The first-stage regression:

```
reg_ss(as.formula(paste("shock ~ ", ctrl)), W = ADH$W,
       X = IV, data = ADH$reg, weights = weights, region_cvar = statefip,
       sector_cvar = sic, method = "all")
#> Estimate: 0.6310409
#>
#> Inference:
#>
#> Std. Error      p-value Lower CI Upper CI
#> Homoscedastic 0.02732516 0.000000e+00 0.5774846 0.6845973
#> EHW           0.08700719 4.083400e-13 0.4605100 0.8015719
#> Reg. cluster  0.09142372 5.113909e-12 0.4518537 0.8102281
#> AKM           0.05296055 0.000000e+00 0.5272402 0.7348417
#> AKMO          0.07671358 1.282891e-03 0.5375710 0.8382827
```

Note that for "AKMO", "Std. Error" corresponds to the normalized standard error, i.e. the length of the confidence interval divided by  $2z_{1-\alpha/2}$ .

The reduced-form and IV regressions:

```

reg_ss(as.formula(paste("d_sh_empl ~", ctrls)), W = ADH$W,
      X = IV, data = ADH$reg, region_cvar = statefip, weights = weights,
      sector_cvar = sic, method = "all")
#> Estimate: -0.4885687
#>
#> Inference:
#>
#> Std. Error      p-value    Lower CI    Upper CI
#> Homoscedastic 0.06332778 1.221245e-14 -0.6126889 -0.3644485
#> EHW          0.11244360 1.392685e-05 -0.7089541 -0.2681833
#> Reg. cluster  0.07578147 1.140306e-10 -0.6370977 -0.3400398
#> AKM           0.16419445 2.924641e-03 -0.8103839 -0.1667535
#> AKMO          0.25437489 4.218033e-04 -1.2368853 -0.2397541
ivreg_ss(as.formula(paste("d_sh_empl ~", ctrls, "| shock")),
        W = ADH$W, X = IV, data = ADH$reg, region_cvar = statefip,
        weights = weights, sector_cvar = sic, method = "all")
#> Estimate: -0.7742267
#>
#> Inference:
#>
#> Std. Error      p-value    Lower CI    Upper CI
#> Homoscedastic 0.1069532 4.523049e-13 -0.9838511 -0.5646022
#> EHW           0.1647892 2.623532e-06 -1.0972075 -0.4512459
#> Reg. cluster  0.1758096 1.063809e-05 -1.1188071 -0.4296462
#> AKM           0.2403730 1.277718e-03 -1.2453492 -0.3031041
#> AKMO          0.3318966 4.218033e-04 -1.6903240 -0.3893132

```

## Note on collinear sectors

Let  $W$  denote the share matrix with the  $(i, s)$  element given by  $w_{is}$ . Suppose that columns of  $W$  are collinear, so it that it has rank  $S_0 < S$ . Without loss of generality, suppose that the first  $S_0$  columns of the matrix are full rank, so that the collinearity is caused by the last  $S - S_0$  sectors. In this case, it is not possible to recover,  $\tilde{X}_s$ , the sectoral shifters with the controls partialled out, and the `reg_ss` and `ivreg_ss` functions will return an error message "Share matrix is collinear". The researcher can either (i) drop the collinear sectors, defining  $X_i = \sum_{s=1}^{S_0} w_{is} X_s$ , (ii) aggregate the sectors, or (iii) if the only controls are those with shift-share structure, and we have data on  $Z_s$ , we can estimate  $\tilde{X}_s$  by running a sector-level regression of  $X_s$  onto  $Z_s$ , and taking the residual. This third option is not currently implemented in this package. Note that options (i) and (ii) change the definition of the estimand. Since they involve changing the shock vector  $X_i$ , this has to be done before using the `reg_ss` and `ivreg_ss` functions.

## References

- Rodrigo Adão, Michal Kolesár, and Eduardo Morales. Inference in shift-share designs: Theory and inference. *Quarterly Journal of Economics*, forthcoming, August 2019. doi: 10.1093/qje/qjz025.
- David H. Autor, David Dorn, and Gordon H. Hanson. The China syndrome: Local labor market effects of import competition in the United States. *American Economic Review*, 103(6):2121–2168, October 2013. doi: 10.1257/aer.103.6.2121.

Timothy J. Bartik. *Who Benefits from State and Local Economic Development Policies?* W.E. Upjohn  
Institute for Employment Research, Kalamazoo, MI, 1991.