

	Slope a_{avg} (m s ⁻²)	Fit error δa (m s ⁻²)	Angle θ (rad)	Error $\delta\theta$ (rad)	$g = a /\sin\theta$ (m s ⁻²)	σ_g (m s ⁻²)	Average g (m s ⁻²)	σ_{SEM} of g (m s ⁻²)
Recording 1	-0.613 ± 0.019	0.019	0.0611	0.00175	10.05	0.43	10.05 $\pm$ 0.33	0.07
Recording 2	-0.621 ± 0.018	0.018	0.0611	0.00175	10.18	0.42		
Recording 3	-0.606 ± 0.014	0.014	0.0611	0.00175	9.93	0.37		

Table 1. Results for each recording. $\theta = 3.5^\circ = 0.0611$ rad with $\delta\theta = 0.1^\circ = 0.00175$ rad (digital-level resolution); $\sigma_g = g\sqrt{[(\delta a/a)^2 + (\delta\theta/\tan\theta)^2]}$. Average g is from the mean slope; its ± 0.33 is propagated from the three fit errors with the shared angle error added once (the value reported). $\sigma_{\text{SEM}} = \text{SD}(g_1, g_2, g_3)/\sqrt{3} = 0.07$ shows only the scatter between recordings and leaves out the angle error.