

# AI1110: Probability and Random Variable

## Assignment-1

Mayank Parasramka\*  
AI22BTECH11018

### Question: 11.16.4.4

'B' = Event of choosing any two tickets with no prize.

$$n(S_2) = {}^{10000}C_2 \quad (5)$$

$$n(B) = {}^{9990}C_2 \quad (6)$$

$$\Pr(B) = \frac{n(S_2)}{n(B)} \quad (7)$$

$$\therefore \Pr(B) = \frac{{}^{10000}C_2}{{}^{9990}C_2} \quad (8)$$

#### Problem Statement:

In a certain lottery 10,000 tickets are sold and ten equal prizes are awarded. What is the probability of not getting a prize if you buy

- (a) one ticket
- (b) two tickets
- (c) 10 tickets.

#### Solution:

Given,

Tickets sold = 10,000

Prizes awarded = 10

Tickets with no awards = 10,000 - 10 = 9990

- (a) If one ticket is bought,  
'S<sub>1</sub>' = Sample space = picking any one of the 10000 tickets.  
'A' = Event of not choosing one of the 10 prize tickets.

$$n(S_1) = {}^{10000}C_1 = 10000 \quad (1)$$

$$n(A) = {}^{9990}C_1 = 9990 \quad (2)$$

$$\Pr(A) = \frac{n(S_1)}{n(A)} = \frac{9990}{10000} \quad (3)$$

$$\therefore \Pr(A) = \frac{999}{1000} \quad (4)$$

- (b) If two tickets are bought,  
'S<sub>2</sub>' = Sample space = picking any two of the 10000 tickets.

- (c) If 10 tickets are bought,  
'S<sub>3</sub>' = Sample space = picking any 10 of the 10000 tickets.

'C' = Event of choosing any 10 tickets with no prize.

$$n(S_3) = {}^{10000}C_{10} \quad (9)$$

$$n(C) = {}^{9990}C_{10} \quad (10)$$

$$\Pr(C) = \frac{n(S_3)}{n(C)} \quad (11)$$

$$\therefore \Pr(C) = \frac{{}^{10000}C_{10}}{{}^{9990}C_{10}} \quad (12)$$

\*The student is with the Department of Artificial Intelligence,  
Indian Institute of Technology, Hyderabad, 502285, India.  
e-mail: ai22btech11018@iith.ac.in.