


Fall Detection using wearable sensor data




Motivation

- Falls are the second leading cause of accidental or unintentional injury deaths worldwide (behind only of road traffic injuries)
- Almost 50% of older adults who fall experience a minor injury
- 25% will experience a more serious injury such as a fracture
- Less than 50% who fell tell their doctor they have had a fall
- Each year, 2.8 million are treated in emergency departments for fall injuries

Research

- A comparison of accuracy of fall detection algorithms (**threshold-based vs. machine learning**) using waist-mounted tri-axial accelerometer signals from a comprehensive set of falls and non-fall trials (SFU, 2015)
- Waist sensor
- Tri-axial (X, Y and Z) Acceleration data
- Collected data, but not used:
 - Six other sensors (head, sternum, left and right thighs, left and right ankles)
 - Tri-axial measurements: Gyro and Magnetic Fields


Research




$$\text{Sensitivity (Recall)} = \frac{\text{True Positive}}{\text{True Positive} + \text{False Positive}}$$

$$\text{Specificity} = \frac{\text{True Negative}}{\text{True Negative} + \text{False Positive}}$$

Project Plan



Exploratory Data Analysis



Falls

Near Falls

Activities of Daily Life

Fall Detection using wearable sensor data

Thank You!

Gustavo Felhberg
Jorge Marciano
Muhammad Muhaimin

