

Prevalence of the photic sneeze reflex: A systematic review and meta-analysis

Josef Trinkl^{1,2} [\[0009-0008-9064-4972\]](#)

Stephan Munkwitz¹ [\[0000-0002-9559-5566\]](#)

Lucien Bickerstaff^{1,3} [\[0000-0002-2683-8130\]](#)

Taisuke Eto¹ [\[0000-0002-0347-7408\]](#)

Manuel Spitschan^{1,3,4,5} [\[0000-0002-8572-9268\]](#) *

¹Translational Sensory & Circadian Neuroscience, Max Planck Institute for Biological Cybernetics, Tübingen, Germany.

²University Hospital and Faculty of Medicine, University of Tübingen, Tübingen, Germany.

³TUM School of Medicine and Health, Chronobiology & Health, Technical University of Munich, Munich, Germany.

⁴TUM Institute for Advanced Study (TUM-IAS), Technical University of Munich, Garching, Germany

⁵TUMCREATE Ltd., Singapore, Singapore

* Correspondence: Manuel Spitschan (manuel.spitschan@tum.de)

Abstract

The photic sneeze reflex (PSR), in which exposure to bright light triggers sneezing, is widely recognised but inconsistently defined and measured. We conducted the first systematic review and meta-analysis of PSR prevalence to synthesise the epidemiological evidence, assess methodological quality, and identify priorities for future research. We included 18 articles comprising 31 study groups and extracted prevalence estimates, study characteristics, ascertainment methods, and epidemiological information. Fifteen eligible study groups classified as healthy were included in the primary meta-analysis. The pooled prevalence was 22% (95% CI, 15%-29%), with extreme between-study heterogeneity ($I^2 = 99.3\%$). Reported prevalence estimates and associations with participant characteristics varied widely, and nearly all studies were judged to be at high risk of bias. The pooled estimate should therefore be interpreted as a descriptive summary of the available evidence rather than a precise estimate of population prevalence. Future studies require a standardised operational definition, representative sampling, transparent reporting, and reproducible methods for assessing light-triggered sneezing.

Keywords: photic sneeze reflex, light sneezing, light sensitivity, prevalence, visual stimuli, epidemiology, systematic review

36 Introduction

37 The photic sneeze reflex (PSR) is a recognisable human response in which sudden
38 exposure to bright light, typically when moving from a darker environment into a well-
39 illuminated one, elicits sneezing. The phenomenon has been described as the photic sneeze
40 reflex, photic sneeze syndrome, sun sneezing, ACHOO (autosomal dominant compulsive
41 helio-ophthalmic outburst) syndrome, and light-induced sneezing. Despite its everyday
42 recognisability, the construct remains poorly standardised. Trigger conditions, latency,
43 frequency, and number of sneezes vary between individuals and populations, although
44 response patterns may be relatively stable within an individual (Semmes et al., 1995). The
45 absence of a shared operational definition complicates comparisons across studies.

46 Reported PSR prevalence estimates range from very low values to more than half of the
47 sampled population. This variation may reflect biological differences, but it also arises from
48 substantial methodological heterogeneity. Studies differ in their populations, recruitment
49 settings, sampling procedures, demographic coverage, and treatment of clinical and
50 ostensibly healthy groups. Ascertainment methods range from single self-report questions
51 and questionnaires to clinical observation and experimental light provocation, with
52 inconsistent wording and exposure conditions. Consequently, the prevalence of PSR is not
53 only biologically uncertain but also methodologically difficult to estimate and interpret.

54 PSR is relevant to sensory integration and reflex physiology because it may reflect
55 interactions between visual input and neural pathways involved in sneezing. Proposed
56 mechanisms include cross-talk between visual and trigeminal pathways, general
57 parasympathetic hypersensitivity, and parasympathetic generalisation through co-activation
58 of closely situated nerve branches (Everett, 1964). These accounts remain plausible but
59 unconfirmed, and no conclusive mechanistic model has been established.

60 Familial aggregation (Beckman and Nordenson, 1983; Everett, 1964; Forrester, 1985; Morris,
61 1987) and genome-wide association studies (Eriksson et al., 2010; Sasayama et al., 2018;
62 Wang et al., 2019) suggest a heritable contribution to PSR. Understanding the conditions
63 under which PSR occurs may also have practical relevance when unexpected sneezing
64 could create risk, including during ocular procedures (Morley et al., 2010; Wessels et al.,
65 1999). However, proposed mechanisms, inheritance patterns, and associations with
66 participant characteristics remain unresolved.

67 No systematic review has previously synthesised the prevalence evidence for PSR or
68 appraised the methodological quality of this literature. We therefore aimed to identify and
69 summarise studies reporting PSR prevalence, extract and classify epidemiological
70 information, assess risk of bias, estimate pooled prevalence in eligible healthy study groups,
71 and define priorities for more standardised future research.

72 Methods

73 Study design and reporting framework

74 This systematic review and meta-analysis of PSR prevalence estimates was conducted and
75 reported in accordance with the PRISMA 2020 statement (Page et al., 2021). No review
76 protocol was registered.

77 Eligibility criteria

78 Studies were eligible if they reported a PSR prevalence estimate or sufficient data to
79 calculate one. We applied no restrictions by study design, population, or language. Eligible
80 evidence included population surveys, clinical and laboratory studies, family studies, and
81 reports in which PSR prevalence was recorded incidentally. Study groups preselected for

photic sneezing were retained for the descriptive synthesis but excluded from pooled prevalence estimation.

Information sources and search strategy

The principal information source was the bibliography compiled previously (Trinkl et al., 2025), which contained 167 records relating to PSR. The bibliography was assembled using MEDLINE (PubMed), Google Books, and Google searches with the terms "photic sneeze", "sun sneeze*", "bright light" AND "sneeze", and variants in English, German, and French. Forward and backward citation searches in Google Scholar were used to supplement the bibliography, which was updated on 23 May 2021 and 31 January 2024 after the initial search on 9 November 2020. An additional unstructured Web of Science search on 10 December 2024 identified no further eligible articles. The bibliography is available at https://github.com/tscnlab/TrinklEtAl_ExpBrainRes_2025.

To update the existing reference library, we conducted a de novo MEDLINE search through PubMed using Medical Subject Headings (MeSH) and free-text terms. The complete Boolean search was:

("Photic sneeze reflex"[MeSH] OR "photic sneeze"[tiab] OR "sun sneeze"[tiab] OR "ACHOO syndrome"[tiab] OR "light-induced sneeze"[tiab] OR "sneezing and light"[tiab] OR "sneeze reflex"[tiab] OR "sneezing response"[tiab])

AND

("Prevalence"[MeSH] OR "epidemiology"[MeSH] OR prevalence[tiab] OR incidence[tiab] OR frequency[tiab] OR proportion[tiab] OR rate[tiab] OR distribution[tiab] OR epidemiologic study[tiab] OR population-based study[tiab] OR cross-sectional study[tiab] OR observational study[tiab] OR cohort study[tiab] OR case-control study[tiab] OR public health surveillance[tiab] OR health survey[tiab] OR national survey[tiab] OR risk factors[tiab] OR demographic factors[tiab] OR genetic predisposition[tiab]))

The final PubMed search was run on 28 June 2026 and returned 18 records.

Study selection

In total, 180 records were screened by title, abstract, and, where appropriate, full text. Eighteen studies reporting PSR prevalence or data from which prevalence could be calculated were included. MS performed the initial screening and selection, and JT subsequently assessed the included studies and extracted the data. Duplicate records and reports without usable human PSR prevalence data were excluded. The study-selection process is summarised in the PRISMA flow diagram (**Figure 1**).

Data extraction

For each eligible study, we extracted bibliographic information, study design, sample size, recruitment method, country and setting, participant demographics, occupation, and reported health status. We also recorded inclusion and exclusion criteria, the operational definition of PSR, the exact ascertainment question where available, the method used to identify photic sneezers, the light source and exposure conditions, potential confounders, and the numbers of participants with and without PSR. When a report contained multiple populations or subgroups, these were extracted separately to support consistent classification and analysis.

Classification of study populations and subgroups

Study groups were classified according to their reported health status as healthy or patient/clinical samples. Groups preselected because participants or families had PSR were described narratively but excluded from pooled prevalence estimates. The primary

quantitative synthesis included eligible healthy study groups; patient and clinical groups were analysed separately as an exploratory comparison.

Risk-of-bias assessment

Risk of bias was assessed using the Joanna Briggs Institute Critical Appraisal Checklist for Prevalence Studies (Joanna Briggs Institute, 2020), with the appraisal focused specifically on the validity of each reported PSR prevalence estimate. Two authors independently assessed each of the 31 study groups (JT with either SM or TE). Initial agreement was reached for 175 of 279 item-level judgements (62.7%). Differences arose mainly from the interpretation of items Q5 (targeting coverage bias) and Q7 (standardisation of measurement). The assessors agreed on operational rules for these items and resolved all remaining disagreements by consensus with reference to the JBI guidance (Munn et al., 2015). For Q1, the sampling frame was judged in relation to the PSR prevalence question rather than the original purpose of the study. For Q3, a required sample size of 289 was calculated using an expected prevalence of 0.25, a 95% confidence level, and precision of 0.05. Overall risk-of-bias judgements were based on the number and likely influence of item-level concerns on the prevalence estimate.

Data synthesis

Study characteristics, ascertainment methods, prevalence estimates, reported associations, and methodological limitations were synthesised narratively. Individual studies were summarised to retain relevant contextual information that could not be represented in the quantitative analysis.

Statistical analysis

Random-effects meta-analyses of proportions were conducted in R (version 4.5.1) using RStudio (version 2025.09.1+401) and the metaprop function of the meta package (version 8.2.1). (Balduzzi et al., 2019). For each study group, prevalence was calculated as the number of participants classified as having PSR divided by the total sample size. The Freeman-Tukey double-arcsine transformation was used to stabilise variances for proportions near the boundaries, and pooled estimates were back-transformed to the original proportion scale. Between-study heterogeneity was quantified using I^2 and τ^2 . The primary analysis included eligible healthy study groups; patient and clinical groups were pooled separately for descriptive comparison.

Data and code availability

The PSR bibliography is available at https://github.com/tscnlab/TrinklEtAl_ExpBrainRes_2025 (Trinkl et al., 2025). The study-level extraction sheet and item-level JBI risk-of-bias assessments are provided in the supplementary material. Analysis code and figure-generation materials should be deposited with the final submission.

Results

Eighteen studies reported PSR prevalence or sufficient data to calculate it. Seven studies contributed more than one distinct population, yielding 31 study groups in total. Twenty groups were classified as healthy and 11 as patient or clinical samples. (Table 1) Five healthy groups were preselected for PSR and were therefore excluded from quantitative synthesis, leaving 15 eligible healthy groups for the primary meta-analysis. Patient and clinical groups were analysed separately as an exploratory comparison.

Nearly all included studies were judged to be at high risk of bias (Table 2). The most frequent concerns involved the representativeness of the sampling frame and recruitment method (Q1

and Q2), sample coverage and response-rate management (Q5 and Q9), and the validity and consistency of PSR ascertainment (Q6 and Q7). Most studies used convenience or otherwise narrowly defined samples, provided incomplete demographic information, or did not report the exact wording used to identify PSR. Definitions also differed in whether they referred specifically to sunlight, included artificial light, or left the light source unspecified. Although several studies reported high response rates and analysed the identified samples appropriately, these strengths did not resolve the broader limitations in sampling and outcome measurement.

Individual study summaries

Sneezing in response to light (Everett, 1964)

Everett investigated associations between PSR and heredity, sex, allergy, and nasal disorders in three samples: psychiatric house officers, 75 hospitalised medical and surgical patients, and students from three colleges. The student survey yielded 339 analysable responses. The study reported associations with sex and ethnicity but not with allergic rhinitis or other allergies, and proposed parasympathetic excitation as a possible common mechanism. The paper also discussed familial patterns, related reflexes, and possible practical implications.

The photic sneeze reflex in the human newborn: a preliminary report (Anderson and Rosenblith, 1968)

During a study of visual following in 275 newborns, the investigators presented a shiny but silent bicycle bell for 1-3 minutes and used a penlight when the bell elicited no response. Seventy-six infants were reported to have sneezed, including 36 who sneezed only during presentation of the bell. Among infants with documented sneeze counts, most sneezed once. Subgroup analyses considered sex and ethnicity, and a separate case described light-triggered sneezing in a newborn with trisomy 13.

Clinical and statistical observations on so-called vasomotor rhinitis (Takahashi, 1942)

Takahashi examined 79 patients with vasomotor rhinitis at the outpatient clinic of Kyoto Imperial University. Light-related sneezing was assessed in 37 patients, of whom 14 (37.8% $\pm$ 7.9%) sneezed after moving from indoors into direct sunlight. Some patients also responded to strong artificial light. On the basis of these observations, the author used the term "Lichtniesen" for light-induced sneezing.

Solar sneeze reflex (Lang and Howland, 1987)

Lang and Howland reported 15 patients with allergic or vasomotor rhinitis who sneezed on entering sunlight, corresponding to approximately 11% of the clinic population. After treatment of rhinitis, four patients no longer reported sunlight-triggered sneezing and three reported an attenuated response, although changes were not clearly related to rhinitis type. A separate survey of healthcare professionals at the same clinic identified PSR in 50 of 144 respondents (35%).

The photic sneeze reflex and ocular anaesthesia (Wessels et al., 1999)

Wessels et al. prospectively assessed PSR and involuntary sneezing during local anaesthesia in 557 patients undergoing ocular surgery. Before drug administration, participants were asked whether they had ever sneezed when entering sunlight or bright light; 78 (14%) responded affirmatively. Twenty-nine patients (5.2%) sneezed after anaesthesia. Sneezing was more frequent after periocular than retrobulbar injection, but the association between a history of PSR and procedure-related sneezing was not statistically significant. The study also found no significant difference in procedure-related sneezing between male and female patients.

Factors prompting sneezing during local anaesthetic injections to the eyelids (Morley et al., 2010)

Morley et al. analysed 314 procedures in 294 patients receiving intravenous sedation and local anaesthetic injections for oculoplastic surgery. A history of photic sneezing was reported by 23% of participants, and sneezing during or within 5 minutes of injection occurred in 51 procedures involving 47 patients. Procedure-related sneezing was associated with male sex, upper-eyelid infiltration, bilateral infiltration, deeper sedation, and midazolam use, whereas opioid derivatives were associated with lower risk. A history of photic sneezing was also associated with procedure-related sneezing (RR 2.6, 95% CI 1.6-4.0).

Selective amplification of ipRGC signals accounts for interictal photophobia in migraine (McAdams et al., 2020)

McAdams et al. studied visual discomfort and pupil responses in 20 control participants and 40 participants with migraine, divided according to the presence or absence of aura. As part of the assessment, participants were asked whether they tended to sneeze when moving from a dark room into bright light. Between 15% and 20% of each group responded affirmatively, with no significant between-group difference.

Individual differences in the sneezing reflex: an inherited physiological trait in humans? (Beckman and Nordenson, 1983)

Among 460 blood donors in Umeå, Sweden, 109 (24%) reported sneezing after sudden exposure to strong light. Women were more numerous among photic sneezers, but the association with sex was not statistically significant. The authors also reported a preliminary family study: in seven families with one photic-sneezing parent, 9 of 14 children reportedly had the trait, whereas no PSR was reported in 11 families in which neither parent had the trait.

Autosomal dominant transmission of the photic sneeze reflex (Peroutka and Peroutka, 1984)

A poll of 25 neurologists at Johns Hopkins Hospital identified PSR in 36%, although only 8% reported prior awareness of the phenomenon. The authors also described a family in which PSR was present in one author, his father, his brother, and subsequently his infant daughter, but not in his mother or wife. The family history was presented as evidence consistent with dominant transmission.

Sneezing on exposure to bright light as an inherited response (Forrester, 1985)

Forrester reported that 29 of 114 consecutive patients attending a general practice (25%) responded affirmatively when asked about sneezing in bright light. Six pedigrees were obtained. Across 15 couples in which one partner reported PSR, 17 of 37 children were also reported to have the trait.

ACHOO syndrome: prevalence and inheritance (Morris, 1987)

Morris presented two case reports with pedigrees interpreted as supporting autosomal dominant inheritance with variable expressivity. Some relatives also reportedly sneezed in response to eyebrow plucking or hair pulling. In an additional questionnaire administered at a meeting of neurologists, 4 of 20 respondents (20%) reported sneezing on exposure to bright light.

Questionnaire survey of sneezing induced by light stimulation in Tohoku (Kodama et al., 1992)

Kodama et al. surveyed 749 people aged from their teens to their sixties in workplaces, schools, and public settings in the Tohoku region of Japan. PSR was reported by 187 respondents (25%; 99 male and 88 female). Reported prevalence was highest among people

in their twenties and somewhat lower in older age groups. Most photic sneezers reported a single sneeze at the moment of glare, although a small minority reported occasional multiple sneezes. Responses indicated substantial individual variation in triggering conditions, including sunlight, artificial light, and the intensity of the transition. Forty-three respondents reported a relative with the same response.

Sneezing in response to bright light: is it a cause of accidents? (Deshmukh, 1995)

Deshmukh combined two family case reports with surveys of college students, high-school students, and operating-room staff. Across the three surveys, 79 of 508 respondents (15.6%) reported sneezing on sudden exposure to bright light. The family reports were interpreted as consistent with dominant inheritance, and the article discussed circumstances in which an unexpected sneeze could present a safety risk.

The photic sneeze response: a descriptive report of a clinic population (Semes et al., 1995)

Semes et al. distributed questionnaires to 500 consecutive patients attending a university optometric clinic; 367 questionnaires were returned and 122 respondents (33.2%) identified as photic sneezers. PSR was more frequently reported by White than Black respondents, and an association with self-reported deviated nasal septum was also observed, although relevant subgroup information was incomplete. Most photic sneezers reported that sunlight triggered sneezing rarely or occasionally, and 107 of 118 respondents with available data reported three or fewer sneezes per episode.

Prevalence of the photo-induced sneezing reflex in a German clinical sample (Kulas et al., 2017)

Kulas et al. administered a standardised questionnaire to 962 patients attending an otorhinolaryngology outpatient clinic. The questionnaire assessed occurrence and frequency of photic sneezing, additional triggers, avoidance strategies, demographic characteristics, familial predisposition, medical history, and substance use. PSR was reported by 542 participants (56%), although 301 described only occasional sneezing; excluding these responses produced an estimate closer to those reported in other studies. Familial predisposition and nicotine consumption were associated with PSR, whereas other examined associations were not statistically significant.

Web-based, participant-driven studies of common genetic traits (Eriksson et al., 2010)

Eriksson et al. analysed 535,076 single-nucleotide polymorphisms in customers of a direct-to-consumer genetic testing company across 22 traits. For photic sneezing, 1,750 of 5,390 participants of Northern European ancestry (32.5%) answered yes to the question, "Do you have a tendency to sneeze when exposed to bright sunlight?" The analysis identified one genome-wide significant intergenic locus on 2q22.3 and a suggestive association near NR2F2 on chromosome 15.

Genome-wide association study of photic sneeze syndrome in a Japanese population (Sasayama et al., 2018)

Sasayama et al. analysed survey and genotype data from 11,409 participants in Japan. Participants who selected "light sneeze reflex" from a list of medical conditions were classified as photic sneezers, producing a prevalence estimate of 3.2%. The analysis supported previously reported loci and identified two additional suggestive associations in the Japanese sample. The ascertainment method, which treated PSR as one item within a list of conditions, differed substantially from the more direct questions used in several other studies.

Genome-wide association study of the photic sneeze reflex in a Chinese population (Wang et al., 2019)

Wang et al. analysed questionnaire and genotype data from 3,417 participants in China and reported a PSR prevalence of 25.6%. Prevalence was higher in male than female

participants (30.1% vs 21.1%). Two independent genetic variants were associated with PSR. One locus on 2q22.3 had been reported previously, whereas a locus on 3p12.1 was newly identified in this analysis. Both associated variants were intergenic, and the 3p12.1 locus was located near CADM2.

Synthesis of results

The 18 included studies had sample sizes ranging from 6 to 11,409 participants and comprised analytical ($n = 6$) and descriptive ($n = 12$) observational designs. Study populations included community and online samples, students, healthcare professionals, blood donors, newborns, family members, and patients recruited in clinical settings. Because several reports included more than one population, 31 distinct study groups were extracted: 20 classified as healthy and 11 as patient or clinical samples. Five healthy groups were excluded from quantitative synthesis because they were preselected for PSR, leaving 15 groups in the primary meta-analysis.

Across the 15 eligible healthy study groups, prevalence estimates ranged from 3% to 36%. The random-effects pooled prevalence was 22% (95% CI, 15%-29%), with extreme heterogeneity ($I^2 = 99.3\%$; $\tau^2 = 0.5102$). Across patient and clinical groups, prevalence estimates ranged from 11% to 56%, with an exploratory pooled estimate of 24% (95% CI, 15%-34%) and similarly high heterogeneity ($I^2 = 97.1\%$; $\tau^2 = 0.5432$). These pooled values therefore summarise highly heterogeneous evidence and should not be interpreted as precise population estimates.

Almost all studies ascertained PSR by self-report, usually through a questionnaire or a direct question about sneezing after exposure to sunlight or bright light. Response formats ranged from binary self-identification to graded frequency or intensity scales and open-ended answers. Only four studies reported the exact wording of the ascertainment question (Eriksson et al., 2010; McAdams et al., 2020; Wang et al., 2019; Wessels et al., 1999) and only one study directly observed sneezing during light exposure (Anderson and Rosenblith, 1968).

All but one study were judged to be at high risk of bias under the JBI appraisal. The principal concerns were non-representative or poorly described sampling methods, inconsistent definitions of PSR and its triggering light source, limited validity of outcome ascertainment, and incomplete or inappropriate statistical analysis.

Reported associations with sex, age, ancestry or ethnicity, and health status were inconsistent and were not examined using comparable methods across studies. Several family and genetic studies suggested a heritable contribution to PSR, but the available evidence did not permit robust estimation of inheritance patterns or subgroup differences.

Discussion

This review provides the first systematic synthesis of PSR prevalence studies, the first pooled prevalence estimate, an explicit appraisal of operational definitions and methodological quality, and a proposed framework for more standardised future assessment. The pooled estimate of 22% across 15 eligible healthy study groups suggests that PSR is not rare, but its interpretation is constrained by extreme heterogeneity and the high risk of bias in nearly all studies. Differences in recruitment, population characteristics, ascertainment questions, light sources, and response definitions mean that studies may not have measured the same underlying construct. The exploratory analysis of patient and clinical samples showed a similar pooled estimate, but the constituent populations were diverse and do not define a single clinically meaningful target population. Reported associations with sex, age,

ancestry or ethnicity, health status, and familial occurrence were inconsistent and should not be treated as established.

The review has several strengths. It draws together literature spanning multiple decades, languages, settings, and study designs; separates distinct study groups; and applies a structured prevalence-specific risk-of-bias appraisal. Its main limitation is the quality of the underlying evidence. Many studies were not designed primarily to estimate PSR prevalence, most relied on convenience samples and self-report, and almost all were at high risk of bias. The evidence base also overrepresents populations of European ancestry and provides insufficient information for reliable comparison across geographic or demographic groups. Accordingly, the pooled prevalence is best understood as a descriptive summary of the available literature rather than a general or global population estimate.

Future studies should use a shared operational definition and distinguish self-reported sunlight-triggered sneezing from responses observed during controlled light provocation. A minimum assessment should include a clearly worded screening question, graded measures of frequency and reproducibility, latency and sneeze count, and separate questions about sunlight and artificial light. Experimental studies should report the source, spectrum, intensity, geometry, duration, and temporal profile of light exposure, as well as pre-exposure conditions. Epidemiological studies require more representative sampling and fuller reporting of recruitment setting, ancestry or ethnicity, age, sex and gender, health status, relevant clinical conditions, medications, and other sneeze triggers. These improvements would allow future research to determine whether observed subgroup differences reflect biological variation or methodological artefact. To facilitate harmonisation across studies, we additionally propose a standardised PSR screening questionnaire incorporating these core elements. The questionnaire is intended as a practical minimum screening tool for self-reported photic sneeze responses and can be included as a supplementary assessment alongside study-specific measures.

Proposed minimum recommendations for future PSR prevalence studies

Operational definition

- The photic sneeze reflex is a predisposition to sneeze following exposure to a bright light stimulus or a sudden increase in ambient light intensity.

Recommended screening question

- Do you sneeze when you are suddenly exposed to bright light, for example when stepping from a darker place into sunlight?
Yes / No / Unsure / Light-induced nasal sensation without sneezing

Frequency, latency, multiplicity, and reproducibility

- How often do you sneeze when exposed to bright light?
Always / Often / Sometimes / Rarely / Never
- What is the typical latency between light exposure and the first sneeze?
- How many sneezes typically occur during a light-triggered episode?
- What is the typical interval between successive sneezes?
- Has this response occurred consistently across your lifetime?
Yes / No / Unsure
- When the same light transition is repeated, how often does it trigger sneezing?
Always / Often / Sometimes / Rarely / Never

Sunlight and artificial light

- Does natural sunlight trigger sneezing?
Yes / No / Unsure
- Does artificial light trigger sneezing?
Yes / No / Unsure

Distinguish self-reported and experimentally provoked PSR

- State whether PSR was assessed by structured questionnaire or interview or observed during controlled light stimulation. Report the instrument, mode of administration, outcome definition, and criteria used to classify a participant as having PSR.

Report light exposure conditions and participant characteristics

- Report the light source; direct or diffuse illumination; continuous, pulsed, or flash exposure; spectral characteristics; illuminance at eye level; pre-exposure conditions; duration and rate of the dark-to-light transition; time of day; relevant environmental conditions; and the use of glasses, sunglasses, or contact lenses.
- Report participant age, sex and gender, ancestry or ethnicity, family history of PSR, allergic or non-allergic rhinitis, migraine, photophobia, relevant ophthalmological or neurological conditions, smoking and exposure to nasal irritants, medications that may affect sneezing, and other substances or stimuli that trigger sneezing.

Conclusion

This systematic review identified a substantial but methodologically weak literature on PSR prevalence. The pooled estimate of 22% across eligible healthy study groups indicates that PSR is commonly reported, but extreme heterogeneity and pervasive risk of bias prevent precise inference about prevalence in any general or global population. The principal limitations are inconsistent definitions, non-representative sampling, and reliance on poorly documented self-report measures. A standardised operational definition, reproducible ascertainment methods, representative sampling, and transparent reporting are required before the epidemiology and determinants of PSR can be characterised reliably.

Supplementary information and declarations

Supplementary information

Data extraction form and item-level risk-of-bias assessments

Authors' contributions

Conceptualisation: MS. Data Curation: MS, JT, TE. Formal Analysis: JT, SM, TE. Funding Acquisition: MS. Investigation: JT, MS. Methodology: JT, MS. Project Administration: MS. Resources: —. Software: —. Supervision: MS. Validation: —. Visualisation: —. Writing – Original Draft Preparation: JT. Writing – Review & Editing: JT, SM, LB, TE, MS.

Funding

Open Access funding enabled and organized by Projekt DEAL. This work was supported by the Max Planck Society (Max Planck Research Group Translational Sensory and Circadian Neuroscience to M.S.), the Wellcome Trust (204686/Z/16/Z to M.S.) and the Alexander von Humboldt Foundation (Ref 3.5 - 1244173 - JPN - HFST-P to T.E.). For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission.

Data availability

The study-level data, risk-of-bias assessments, analysis code, and figure-generation materials will be made available under a CC BY licence in the project repository [permanent repository URL to be added].

Declarations

Conflict of interest

The authors have no conflicts of interest to declare.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Declaration of AI-assisted technologies

During preparation of this manuscript, the authors used ChatGPT to support language editing and the organisation of information. The authors subsequently reviewed and revised all content and take full responsibility for the final manuscript.

469 Photoc Sneeze Survey Questionnaire

470 Please complete the survey below. Thank you!

471 Thank you for agreeing to participate in this survey.

472 Please note: Some questions may seem repetitive. We include these to check consistency. Please answer

473 all questions to the best of your knowledge and ability.

1. Do you have a tendency to sneeze when exposed to bright sunlight?

- ☐ Yes
- ☐ No, what are you talking about

2. Do you sneeze in response to bright light?

- ☐ Yes
- ☐ No

3. Does bright light give you a tickle sensation?

- ☐ Yes
- ☐ No

4. Do you only sneeze when exposed to outdoor bright light (daylight, including through windows) or indoor bright light (electric light)?

- ☐ Only outdoors
- ☐ Only indoors
- ☐ Both
- ☐ Don't know

5. When you feel like you are about to sneeze, do you try to elicit it by looking at a bright light?

- ☐ Always
- ☐ Very Often
- ☐ Sometimes
- ☐ Rarely
- ☐ Never

6. How old were you when you first discovered that you sneeze in response to bright light?
Approximate age in years.
If you are not sure, leave this field empty.

7. How old are you now?
Exact age in years.

8. You've been in a dark area for two hours or more, and suddenly enter bright light, for example bright sunlight after being indoors. Do you sneeze?

- ☐ Always
- ☐ Usually
- ☐ Sometimes
- ☐ Never

9. What happens when the light you enter is only moderate? Do you sneeze?

- ☐ Always
- ☐ Usually
- ☐ Sometimes
- ☐ Never

10. And what about entering dim light after being in a dark area?

- ☐ Always
- ☐ Usually
- ☐ Sometimes
- ☐ Never

11. How quickly do you

- ☐ Within less than 5 seconds

sneeze when you enter bright light?	☐ After 5-10 seconds ☐ After more than 10 seconds ☐ After more than 1 minute ☐ After more than 5 minutes
12. When you are exposed to the same bright-light conditions on different occasions, how consistently does bright light make you sneeze?	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never ☐ Don't know
13. How many sneezes typically occur following a bright-light exposure that triggers sneezing?	☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 or more ☐ Varies ☐ Don't know
14. Please select "Sometimes" for this question. Thank you for paying attention!	☐ Always ☐ Usually ☐ Sometimes ☐ Never
15. Please select which of the following general statements you agree with (multiple answers are possible).	☐ I sneeze in response to causes unrelated to light ☐ I use light to elicit a sneeze caused by other causes ☐ I sneeze in response to light without prior feeling of needing to sneeze

474

475 Sneezing in daylight

476

477 Please select to what extent exposure to different levels of DAYLIGHT elicits a sneeze.

Stimulus or situation	Response
16. Direct sunlight	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
17. Full daylight (not direct sun)	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
18. Overcast day	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
19. Sunset/sunrise	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never

20. Twilight	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
--------------	---

478

21. Please select "Disagree" for this question. Thank you for paying attention!

☐ Strongly Agree
☐ Agree
☐ Undecided
☐ Disagree
☐ Strongly Disagree

479

480 Sneezing in electric light

481

482 Please select to what extent exposure to different levels of ELECTRIC LIGHT elicits a sneeze.

Stimulus or situation	Response
22. Bright room illumination	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
23. Moderate room illumination	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
24. Dim room illumination	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
25. Monitors, smartphone displays, TVs	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
26. Direct view of light sources (e.g. lamps)	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
27. Car headlights (at night)	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
28. Other (please describe below)	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never

483

484 Aspects of light making you sneeze

485

30. Which of the following aspects of light direction would in your experience trigger a sneeze?

- ☐ The light must come from above
- ☐ The light must come from below
- ☐ The light can come from anywhere
- ☐ Don't know

31. Does it matter whether the light is relatively uniform across your entire vision?

- ☐ The light must be uniform
- ☐ The light does not need to be uniform
- ☐ Don't know

Time of day

In this section, we would like to ask you whether there are any specific times during the day that you are more likely to sneeze due to BRIGHT LIGHT.

33. Are there any specific times of day that you are more likely to sneeze due to BRIGHT LIGHT?

- ☐ Yes
- ☐ No
- ☐ Don't know

34. During which time window are you more likely to sneeze due to BRIGHT LIGHT?

- ☐ 0:00 - 2:00 am
- ☐ 2:00 - 4:00 am
- ☐ 4:00 - 6:00 am
- ☐ 6:00 - 8:00 am
- ☐ 8:00 - 10:00 am
- ☐ 10:00 - 12:00 am
- ☐ 12:00 - 2:00 pm
- ☐ 2:00 - 4:00 pm
- ☐ 4:00 - 6:00 pm
- ☐ 6:00 - 8:00 pm
- ☐ 8:00 - 10:00 pm
- ☐ 10:00 - midnight

Season

In this section, we would like to ask you whether there are any specific seasons during which you are more likely to sneeze in response to bright light.

35. Do you experience a change in your susceptibility to sneeze due to BRIGHT LIGHT across the season?

- ☐ No change
- ☐ Slight change
- ☐ Moderate change
- ☐ Marked change
- ☐ Extremely marked change
- ☐ Don't know

36. During which season are you most likely sneeze due to BRIGHT LIGHT?

- ☐ Winter (December 21 - March 20)
- ☐ Spring (March 21 - June 20)
- ☐ Summer (June 21 - September 20)
- ☐ Autumn (September 21 - December 20)

37. Please select "Agree" for this question. Thank you for paying attention!

- ☐ Strongly agree
- ☐ Agree
- ☐ Undecided
- ☐ Disagree
- ☐ Strongly disagree

Sneezing due to other stimuli

We would like to know about other stimuli or objects that make you sneeze. Please select to what extent other stimuli elicit a sneeze.

Stimulus or situation	Response
38. Inhaling cold air	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
39. Inhaling dust	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
40. Eating chocolate	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
41. Eating mint or mint-flavoured products	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
42. Smelling mint or mint-flavoured products	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
43. Smelling pepper	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
44. Eating spicy foods	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
45. Drinking wine	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
46. Smelling perfume and fragrances	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
47. Smelling cleaning products	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never
48. Experiencing sexual arousal	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never ☐ Don't know

49. Experiencing strong emotions	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never ☐ Don't know
50. Plucking eyebrows	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never ☐ Don't know
51. Plucking nose hair	☐ Always ☐ Very Often ☐ Sometimes ☐ Rarely ☐ Never ☐ Don't know

501

502 **Some information about you**

503

504 Finally, we would like to know a few things about you.

52. What is your sex? ☐ Male ☐ Female

If you are one or more of non-binary, transgender, have variations of sex characteristics, sometimes also known as intersex, the answer you give can be different from what is on your birth certificate.

If you are not sure how to answer, use the sex registered on your official documents, such as passport or driving

licence, or
whichever
answer
best
describes
your sex.

53. Do you _____
have any _____
other _____
comments _____
? _____

54. I certify ☐ Yes
that I ☐ No
answered
the above
questions
accurately.


505

506 **Researcher note**

507 Ancestry and ethnicity, relevant health status, and medication use are not included as questionnaire items here.
508 Depending on the study aims, population, and analysis plan, researchers should consider collecting these data
509 separately when and if applicable.

510


511 **Figures and Tables**



- 1 **Figure 1.** PRISMA 2020 flow diagram showing the study-selection process (adapted from Page et al., 2021).

2

3




4

5 **Figure 2.** Forest plot of PSR prevalence in the 15 eligible healthy study groups.

6

7



8

9 **Figure 3.** Forest plot of PSR prevalence in study groups classified as patient or clinical
10 samples.

11

Study group	Country	Population	Health status category	Sample size	Age	Sex	PSR definition	Diagnostic method	PSR question	Light stimulus	Prevalence proportion
Sasayama et al. (2018)	Japan	Japanese population	Healthy	11409	Mean age: Sneezers: 46.3 years Non-sneezers: 50.0 years	sneezers: 176 m, 184 f non-sneezers: 5849 m, 5200 f	tendency to sneeze when the eye is exposed to bright light	questionnaire	individuals who checked "light sneeze reflex" in a list of medical conditions/diseases in online survey were treated as PSR cases.	bright light	0.03
Everett (1964) house officers	USA	House officers	Patient	18	Adults	6 m, 12 unclear	tendency to sneeze when suddenly exposed to bright light	self-report	not specified	bright light	0.33
Everett (1964) Patients		Medical and surgical patients	Patient	75	Adults	49 m, 26 f		self-report			0.11
Everett (1964) Students		Medical students, college juniors & seniors	Healthy	339	Young adults	193 m, 146 f		questionnaire			0.18
Lang & Howland (1987) Patients	USA	Rhinitis patients	Patient	139	24 to 61 years	5 m, 10 f	Sneezing upon entering sunlight; Tendency to sneeze one to 20 times in sunlight	self-report	not specified	sunlight	0.11
Lang & Howland (1987) healthcare professionals		Healthcare professionals	Healthy	144	not specified	unclear					0.35
Deshmukh (1995) college students	USA	College students	Healthy	31	Mean age: 20 years	unclear	reflex action when exposed to sudden bright light	self-report	not specified	bright light	0.19
Deshmukh (1995) operating staff		Operating room staff	Healthy	36	Mean age: 38 years						0.33
Deshmukh (1995) highschool students		High school students	Healthy	441	Mean age: 16 years						0.14
Wessels et al. (1999)	USA	Patients undergoing eye surgery	Patient	557	16 to 97 years Mean age: 69.9 years	223 m, 334 f	sneezing is provoked by exposure to bright light	self-report	Have you ever noticed yourself sneezing when going into the sun or into bright light?	bright light	0.14
McAdams et al. (2020) healthy participants	USA	Control group	Healthy	20	Mean age: 31 years	7 m, 13 f	sneezing in response to sunlight	self-report	subjects were asked if they "tend to sneeze when [they] step out of a dark room into bright sunlight"	bright sunlight	0.20
McAdams et al. (2020) migraineurs with aura		Migraineurs with aura	Patient	20	Mean age: 31 years	1 m, 19 f					0.15
McAdams et al. (2020) migraineurs without aura		Migraineurs without aura	Patient	20	Mean age: 30 years	3 m, 17 f					0.15
Morris (1987) neurologists	USA	Neurologist Grand Round	Healthy	20	not specified	15 m, 5 f	sneezing on exposure to bright light	questionnaire	not specified	bright light	0.20
Morley et al. (2010)	UK	Patients undergoing eye surgery	Patient	248	19 to 105 years	unclear	sneezing on exposure to bright sunlight	self-report	not specified	bright sunlight	0.23
Beckman & Nordenson (1983) blood donors	Sweden	Blood donors	Healthy	460	not specified	unclear	sudden exposure to strong light may induce sneezing	self-report	not specified	bright light	0.24
Beckman & Nordenson (1983) preliminary study		Families from a preliminary study	Healthy	73				not specified			0.22
Kodama et al. (1992)	Japan	Japanese population	Healthy	749	10 to 69 years	391 m, 358 f	light-sneeze reflex	questionnaire or interview	not specified	bright light	0.25
Kulas et al. (2016)	Germany	Patients in otorhinolaryngology outpatient center	Patient	962	not specified	440 m, 534 f	sneezing in response to bright light, sunlight and artificial light	questionnaire	not specified	bright light	0.56
Forrester (1985)	UK	Patients at a general surgery practice	Patient	114	not specified	unclear	sneezing in response to bright light	unclear	not specified	bright light	0.25
Wang et al. (2019)	PR China	Chinese population	Healthy	3417	16 to 70 years	1700 m, 1717 f	uncontrollable reflexive sneezing in response to sudden exposure to bright light	self-report	"in most cases, when you enter from a dark or dim environment suddenly into a bright environment, would you: A. sneeze, B. be a little responsive but do not sneeze, and C. have no response"	bright light/environment	0.26
Anderson & Rosenblith (1968)	USA	Newborns	Healthy	275	Newborns	137 m, 138 f	sneezing in response to bright light	clinical observation	not specified	reflection of a shiny bicycle bell or flashlight shining into the eyes	0.28
Eriksson et al. (2010)	Online	Northern European descent	Healthy	5390	not specified	unclear	tendency to sneeze when moving from relative darkness into bright light — most often sunlight	questionnaire	Do you have a tendency to sneeze when exposed to bright sunlight?	bright light; emphasis on sunlight	0.32
Semes et al. (1995)	USA	Patients receiving	Patient	367	12 to 89 years	unclear	sneezing in response to	questionnaire	not specified	unclear. Some answers	0.33

		eye examination			Mean age sneezers: 45.8 years non-sneezers: 44.0 years		bright light			refer to sunlight, but it was also asked for sneezing in response to other bright light stimuli	
Peroutka & Peroutka (1984) neurologists poll	USA	Neurologists	Healthy	25	not specified	unclear	sneezing in response to bright light	poll between neurologists	not specified	bright light	0.36
Takahashi (1942)	Japan	Rhinitis patients at outpatient clinic	Patient	79	1 to 60 (2 cases over 60)	62 m, 17 f	sneezing in response to bright light (Lichtniesen)	clinical observation & direct interview	not specified	bright light	0.38

12

13 **Table 1.** Characteristics of the included studies and study groups. m, male; f, female.

14

15

	Risk of bias									
	D1	D2	D3	D4	D5	D6	D7	D8	D9	Overall
Sasayama et al. (2018)	X	X	+	X	+	X	+	X	+	X
Everett (1964), house officers	X	X	X	+	X	X	X	X	X	X
Everett (1964), patients	X	X	X	+	+	X	X	X	+	X
Everett (1964), students	X	X	+	+	+	X	+	X	+	X
Lang & Howland (1987), patients	X	X	X	X	X	X	X	X	-	X
Lang & Howland (1987), healthcare professionals	X	X	X	X	X	X	X	X	-	X
Deshmukh (1995), college students	X	X	X	X	+	X	X	X	+	X
Deshmukh (1995), operating staff	X	X	X	X	+	X	X	X	+	X
Deshmukh (1995), highschool students	X	X	+	X	+	X	X	X	+	X
Deshmukh (1995), case report 1	X	X	X	X		X	X			X
Deshmukh (1995), case report 2	X	X	X	X		X	X			X
Wessels et al. (1999)	X	X	+	+	+	X	+	+	+	X
McAdams et al. (2020), healthy participants	X	X	X	X	+	X	X	X	+	X
McAdams et al. (2020), migraineurs with aura	X	X	X	X	+	X	X	X	+	X
McAdams et al. (2020), migraineurs without aura	X	X	X	X	+	X	X	X	+	X
Morris (1987), case report 1	X	X	X	X		X	X			X
Morris (1987), case report 2	X	X	X	X		X	X			X
Morris (1987), neurologists	X	X	X	X	+	X	+	X	-	X
Morley et al. (2010)	X	X	X	+	X	X	X	X	X	X
Beckman & Nordenson (1983), blood donors	X	X	+	X	+	X	X	X	+	X
Beckman & Nordenson (1983), preliminary study	X	X	X	X	+	X	X	X	+	X
Kodama et al. (1992)	X	X	+	X	X	X	X	X	+	X
Kulas et al. (2016)	X	X	+	X	X	X	+	X	X	X
Forrester (1985)	X	X	X	X	+	X	X	X	+	X
Wang et al. (2019)	X	X	+	+	+	X	+	+	+	X
Anderson & Rosenblith (1968)	X	X	X	+	+	-	-	X	X	X
Eriksson et al. (2010)	+	+	+	+	+	+	+	X	+	+
Semes et al. (1995)	X	X	+	X	X	X	+	X	X	X
Peroutka & Peroutka (1984), neurologists poll	X	X	X	X	+	X	X	X	+	X
Peroutka & Peroutka (1984), case report	X	X	X	X		X	X			X
Takahashi (1942)	X	X	X	X	X	X	-	X	+	X

D1: JBI Q1
D2: JBI Q2
D3: JBI Q3
D4: JBI Q4
D5: JBI Q5
D6: JBI Q6
D7: JBI Q7
D8: JBI Q8
D9: JBI Q9

Judgement
 High
 Unclear
 Low
 Not applicable

16

17 **Table 2.** Item-level risk-of-bias judgements using the JBI Critical Appraisal Checklist for
18 Prevalence Studies, visualised in a colour-blind-friendly format adapted from robvis
19 (McGuinness and Higgins, 2021).

20 References

- 21 Anderson, R.B., Rosenblith, J.F., 1968. The photic sneeze reflex in the human newborn: A preliminary
22 report. *Dev. Psychobiol.* 1, 65–66. <https://doi.org/10.1002/dev.420010113>
- 23 Balduzzi, S., Rücker, G., Schwarzer, G., 2019. How to perform a meta-analysis with R: a practical
24 tutorial. *Evid. Based Ment. Health* 22, 153–160. [https://doi.org/10.1136/ebmental-2019-](https://doi.org/10.1136/ebmental-2019-300117)
25 300117
- 26 Beckman, L., Nordenson, I., 1983. Individual differences with respect to the sneezing reflex: an
27 inherited physiological trait in man? *Hum. Hered.* 33, 390–391.
28 <https://doi.org/10.1159/000153407>
- 29 Deshmukh, N., 1995. Sneezing Response to Bright Light: Is it a Cause of Accidents? *Guthrie J.* 64, 104–
30 105. <https://doi.org/10.3138/guthrie.64.3.104>
- 31 Eriksson, N., Macpherson, J.M., Tung, J.Y., Hon, L.S., Naughton, B., Saxonov, S., Avey, L., Wojcicki, A.,
32 Pe'er, I., Mountain, J., 2010. Web-Based, Participant-Driven Studies Yield Novel Genetic
33 Associations for Common Traits. *PLoS Genet.* 6, e1000993.
34 <https://doi.org/10.1371/journal.pgen.1000993>
- 35 Everett, H.C., 1964. Sneezing in response to light. *Neurology* 14, 483–483.
36 <https://doi.org/10.1212/WNL.14.5.483>
- 37 Forrester, J.M., 1985. Sneezing on exposure to bright light as an inherited response. *Hum. Hered.* 35,
38 113–114. <https://doi.org/10.1159/000153527>
- 39 Joanna Briggs Institute, 2020. Checklist for Prevalence Studies. Joanna Briggs Institute.
- 40 Kodama, M., Sato, A., Kuchiiwa, S., 1992.
41 東北地方における光刺激によって誘発されるくしゃみ反射に関するアンケート調査 -
42 Questionnaire Survey on Sneezing Reflex Induced by Light Stimulus in Tohoku. *Igaku*
43 *Seibutsugaku* 125, 215–219.
- 44 Kulas, P., Hecker, D., Schick, B., Bozzato, A., 2017. Investigations on the prevalence of the photo-
45 induced sneezing reflex in the German population, a representative cross-sectional study. *Eur.*
46 *Arch. Oto-Rhino-Laryngol. Off. J. Eur. Fed. Oto-Rhino-Laryngol. Soc. EUFOS Affil. Ger. Soc. Oto-*
47 *Rhino-Laryngol. - Head Neck Surg.* 274, 1721–1725. [https://doi.org/10.1007/s00405-016-](https://doi.org/10.1007/s00405-016-4256-2)
48 4256-2
- 49 Lang, D.M., Howland, W.C., 1987. Solar sneeze reflex. *JAMA* 257, 1330–1331.
50 <https://doi.org/10.1001/jama.1987.03390100068019>
- 51 McAdams, H., Kaiser, E.A., Igdalova, A., Haggerty, E.B., Cucchiara, B., Brainard, D.H., Aguirre, G.K.,
52 2020. Selective amplification of ipRGC signals accounts for interictal photophobia in migraine.
53 *Proc. Natl. Acad. Sci. U. S. A.* 117, 17320–17329. <https://doi.org/10.1073/pnas.2007402117>
- 54 Morley, A.M.S., Jazayeri, F., Ali, S., Malhotra, R., 2010. Factors prompting sneezing in intravenously
55 sedated patients receiving local anesthetic injections to the eyelids. *Ophthalmology* 117,
56 1032–1036. <https://doi.org/10.1016/j.opthta.2009.09.007>
- 57 Morris, H.H., 1987. ACHOO syndrome. Prevalence and inheritance. *Cleve. Clin. J. Med.* 54, 431–433.
58 <https://doi.org/10.3949/ccjm.54.5.431>
- 59 Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., Shamseer, L.,
60 Tetzlaff, J.M., Akl, E.A., Brennan, S.E., Chou, R., Glanville, J., Grimshaw, J.M., Hróbjartsson, A.,
61 Lalu, M.M., Li, T., Loder, E.W., Mayo-Wilson, E., McDonald, S., McGuinness, L.A., Stewart, L.A.,
62 Thomas, J., Tricco, A.C., Welch, V.A., Whiting, P., Moher, D., 2021. The PRISMA 2020
63 statement: an updated guideline for reporting systematic reviews. *BMJ* n71.
64 <https://doi.org/10.1136/bmj.n71>
- 65 Peroutka, S.J., Peroutka, L.A., 1984. Autosomal dominant transmission of the “photic sneeze reflex.”
66 *N. Engl. J. Med.* 310, 599–600.
- 67 Sasayama, D., Asano, S., Nogawa, S., Takahashi, S., Saito, K., Kunugi, H., 2018. A genome-wide
68 association study on photic sneeze syndrome in a Japanese population. *J. Hum. Genet.* 63,
69 765–768. <https://doi.org/10.1038/s10038-018-0441-z>
- 70 Semes, L.P., Amos, J.F., Waterbor, J.W., 1995. The photic sneeze response: a descriptive report of a
71 clinic population. *J. Am. Optom. Assoc.* 66, 372–377.

72 Takahashi, S., 1942. 所謂血管運動神經性鼻炎ノ臨床的統計的觀察 - Clinical and Statistical
73 Observations on So-called Vasomotor Rhinitis. Kyoto Imp. Univ. Fac. Med.
74 Trinkl, J., Bickerstaff, L., Munkwitz, S., Spitschan, M., 2025. Stimulus conditions eliciting sneezing in
75 response to bright light. Exp. Brain Res. 243, 56. [https://doi.org/10.1007/s00221-024-06988-](https://doi.org/10.1007/s00221-024-06988-4)
76 4
77 Wang, M., Sun, X., Shi, Y., Song, X., Mi, H., 2019. A genome-wide association study on photic sneeze
78 reflex in the Chinese population. Sci. Rep. 9, 4993. [https://doi.org/10.1038/s41598-019-](https://doi.org/10.1038/s41598-019-41551-0)
79 41551-0
80 Wessels, I.F., Wessels, D.A., Zimmerman, G.J., 1999. The photic sneeze reflex and ocular anesthesia.
81 Ophthalmic Surg. Lasers 30, 208–211.
82
83