

Statistical analysis plan for the 'Pharyngeal electrical stimulation for acute stroke dysphagia trial' (PhEAST) (ISRCTN98886991)

Lisa J Woodhouse, PhD;¹ Iris I Mhlanga, MSc;¹ Cristina Roadevin, PhD;² Jacqui K Benfield, PhD;¹ Lisa F Everton, PhD;^{1,3} Gwenllian Wilkinson; PhD;^{1,3} Sarah Greatrex;¹ Cameron JC Skinner, MSc;¹ Gemma Squires;¹ Amanda Buck;¹ Corinne Latulipe, BSc;¹ Kennedy M Cadman;¹ Nikola Sprigg, PhD FRCP;^{1,4} Kailash Krishnan, PhD FRCP;^{1,4} Jason P Appleton, PhD MRCP;^{1,4,5} Karl Matz, MD;⁶ Helle K Iversen, PhD MD;⁷ Satish Mistry, PhD;⁸ Marilyn James, PhD;² Timothy J England, PhD FRCP;^{1,5} Shaheen Hamdy; PhD FRCP;^{8,9} Alan A Montgomery, PhD;² Philip M Bath, DSc FMedSci;^{1,4} for PhEAST Investigators

1. Stroke Trials Unit, University of Nottingham, Nottingham UK
2. Nottingham Clinical Trials Unit, University of Nottingham, Nottingham UK
3. Speech & Language Therapy, Nottingham University Hospitals NHS Trust, Nottingham, UK
4. Stroke, Nottingham University Hospitals NHS Trust, Nottingham UK
5. Stroke, University Hospitals of Derby and Burton NHS Foundation Trust, Derby UK
6. Centre for Clinical Neurosciences and Prevention, University of Continuing Education Krems, Krems Austria
7. Department of Neurology, University hospital Rigshospitalet, Denmark
8. Phagenesis Ltd, Manchester UK
9. Centre for Gastrointestinal Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester UK

Short title: PhEAST trial statistical analysis plan

Correspondence: Professor Philip M Bath
Stroke Trials Unit, Mental Health & Clinical Neuroscience, University of Nottingham,
Nottingham NG7 2UH
Email: philip.bath@nottingham.ac.uk

PhEAST SAP

11/08/2026

Version 0.871

Keywords: dysphagia; pharyngeal electrical stimulation; statistical analysis plan; stroke; trial

ABSTRACT

Introduction

Post stroke dysphagia is common, associated with poor functional outcome and lacks treatment strategies beyond behaviour therapies delivered by speech & language therapists. Here, we present the statistical analysis plan for the ongoing pharyngeal electrical stimulation for acute stroke dysphagia trial (PhEAST). PES is a candidate treatment for dysphagia present in non-ventilated stroke patients.

Methods

PhEAST is an investigator-initiated international prospective randomised open-label blinded-endpoint phase-4 superiority trial involving 650 participants with tube-dependent post-stroke dysphagia. Consenting patients are randomised to PES versus no PES given on top of standard care with PES given daily for 6 days. The primary outcome is the dysphagia severity rating scale (DSRS), a measure of swallowing impairment, made at days 14 and 90 and analysed using repeated measures regression.

Conclusion

We present the statistical analysis plan for the main analyses based on data up to day 90 along with planned secondary analyses including presentation of baseline data, health economics, cognition and extended follow-up to 12 months.

INTRODUCTION

Post-stroke dysphagia (PSD, swallowing problems after stroke) is common affecting 60% or more of people with acute stroke and is associated with a poor outcome.¹ Whilst many patients regain safe swallowing, others continue to suffer from dysphagia for weeks, months or even years. Although food, fluids and drugs will initially need to be administered enterally via a nasogastric tube (NGT), a permanent feeding tube (percutaneous endoscopic gastrostomy [PEG], radiologically-inserted gastrostomy [RIG] or jejunostomy tube [J-tube]) will be needed to facilitate discharge from hospital and long-term feeding if dysphagia persists. Poor outcomes result from aspiration, e.g. causing pneumonia, and reduced nutritional intake, e.g. causing malnutrition. Either can lead to poor functional outcome and death. The management of PSD and staff time in hospital result in high healthcare costs so any effective intervention that reduces swallowing impairment would likely have significant health economic benefits.

The mainstay of management of PSD are screening (usually by nurses),² diagnosis and then treatment. Behavioural interventions (e.g. Shaker, chin tuck against resistance and base of tongue exercises), postural changes and compensatory strategies (e.g. thickening drinks, modifying food and clearing swallows) are the most widely used therapeutic approaches.³ In most countries, these are administered by speech and language therapists (SLTs, also called speech and language pathologists) although occupational therapists (OTs) have the same role in other countries, e.g. Denmark. SLTs may also administer physical interventions involving thermal or tactile stimulation.⁴ In addition to pharmacotherapy, a variety of devices have also been tested including acupuncture, neuromuscular electrical stimulation (NMES), repetitive transcranial magnetic stimulation (rTMS) and pharyngeal electrical stimulation (PES).^{3,4} None of these are used widely, largely because there is insufficient large high quality trial data to inform guidelines and practice. In a Cochrane meta-analysis, the potential for efficacy was seen for all the above interventions as judged by reductions (significant or non-significant tendencies) in swallowing impairment and dysphagia.⁴ However, most trials were small ($N < 100$) and the meta-analyses showed considerably heterogeneity ($I^2 > 50\%$). Further, there was evidence of publication bias for most interventions to the extent that any evidence of efficacy might disappear if missing trials had been published and included in the analyses.

Although devices generally have a *conformité européenne* (European conformity) mark and/or FDA approval, gatekeepers to their use, such as insurers, healthcare commissioners or the UK's National Institute for Health and Care Excellence (NICE) may not approve them for general use. For example, NICE noted for NMES that "*For dysphagia after a stroke, the evidence suggests potential benefit, but is limited in quality and quantity. Therefore, this procedure should only be used with special arrangements for clinical governance, consent, and audit or research*".⁵ The solution to these concerns is to perform large definitive phase-3/4 randomised controlled trials that assess efficacy or effectiveness, safety and health economics and so provide level 1a/b evidence that influences healthcare gatekeepers, guidelines and clinical practice.

Pharyngeal electrical stimulation (PES)

PES has a robust theoretical base and a well-defined treatment paradigm, including only requiring 3-6 days of treatment.⁶ Human swallowing has bilateral representation in the brain with a 'dominant' cortex (unrelated to handedness).⁶ Dysphagia often follows a stroke affecting the dominant swallowing cortex and may be exacerbated in recurrent stroke. Swallowing is dependent on afferent feedback via bulbar cranial nerves innervating the pharynx. Increased sensory input from the pharynx, as delivered as PES, has been shown to drive long-term beneficial changes in the cortical control of swallowing⁷ with reorganisation of the swallowing cortex.⁷⁻⁹

A phase-1 study¹⁰ suggested that PES should be delivered at 5 Hz for 10 minutes, a paradigm that maximises the effect on brain excitability.^{10,11} In a randomised dose-comparison phase-2a trial in patients with subacute non-ventilated stroke, PES reduced radiological aspiration as measured using the penetration aspiration scale score (PAS).¹² PES also reduced swallowing impairment (assessed as the dysphagia severity rating scale, DSRS) and length of stay in hospital in patients with PSD in a sham-controlled parallel group phase II trial.¹² In a NIHR RfPB-funded multicentre phase-2b randomised sham-controlled trial, the study was feasible in respect of recruitment, compliance and retention; PES non-significantly reduced clinical dysphagia and length of stay in hospital.¹³ An individual patient data meta-analysis of these three small phase II trials (n=73) found that PES significantly reduced aspiration (PAS) and swallowing impairment (DSRS) and was safe and well

tolerated.¹⁴

The multicentre international randomised single-blind phase-3 STEPS trial assessed PES versus sham in 162 patients with a recent non-ventilated ischaemic or haemorrhagic stroke and radiological penetration or aspiration (PAS>=3). The study was feasible (recruitment, compliance and retention) and PES was safe but did not reduce aspiration (PAS) or swallowing impairment (DSRS) relative to sham.¹⁵ Likely explanations for the neutral results included: i) undertreatment in the PES group (average stimulation 14.8mA); ii) partial stimulation in the sham group (through assessment of threshold and tolerability levels); and iii) recruitment of patients whose swallow impairment was too mild (and so who were likely to improve anyway).

PHADER was an international single-arm study in 245 patients with unventilated stroke, ventilated stroke, ventilated-related, traumatic brain injury or other causes of neurogenic dysphagia.¹⁶ PES was associated with improved DSRS and PAS over 90 days, both overall and in each diagnostic group including in both non-ventilated and ventilated stroke. Analyses from the recent PhEED trial and MAPS audit registry also suggest that PES is associated with improved swallowing over a period of at least 90 days.¹⁷*ref Importantly, there was no control group in these three studies

In spite of this accumulation of clinical trial data and having a CE mark and FDA designation, PES lacks an adequate evidence base on efficacy, safety and information on health economics for non-ventilated stroke patients. Further, NICE say that “... *more research is needed on pharyngeal electrical stimulation. This procedure should only be done as part of a formal research study and a research ethics committee needs to have approved its use. More research is needed on details of patient selection (including the cause of dysphagia and the timing of the intervention) and effects on length of hospital stay compared with usual care*”.¹⁸ The PhEAST trial is designed to add significantly to the evidence base and address NICE questions.

PES is effective for accelerating decannulation of intubated stroke patients post-ventilation^{19,20} and there is common agreement for using PES in this group in European, UK and US guidelines.²¹⁻²³ Importantly, PhEAST does not address this population but rather focuses on non-ventilated stroke patients with dysphagia.

METHODS

Design

The pharyngeal electrical stimulation for acute stroke dysphagia trial (PhEAST) is assessing the safety and effectiveness of PES in 650 patients with PSD from the UK, Austria and *Denmark. Full information is given in the latest protocol (version 12, 19 May 2026: <https://stroke.nottingham.ac.uk/pheast/docs/>) and a published version of this.²⁴ Adults with acute or subacute stroke (recruitment between 2 and 31 days) and who are tube-dependent for fluids and food (functional oral intake scale,²⁵ FOIS score 1-3) are randomised 1:1 to six days of PES versus no PES; practically, this amounts to PES vs NGT and all participants receive best medical-SLT/usual care. The primary outcome is swallow impairment assessed using the dysphagia severity rating scale (DSRS)^{12,26} at 14 and 90 days and blinded to treatment assignment to avoid observer bias. Analysis will compare the two treatment groups using adjusted repeated measures regression. Although PhEAST extends follow-up to 365 days as a result of secondary funding, this was not the original design and we will focus on reporting data up to 90 days in the primary results publication.

The present paper details the statistical analysis plan (SAP) and health economics analysis plan (HEAP) as well as intended secondary publications, as presented in the accompanying supplements. This information is presented during recruitment, blinded to treatment assignment and prior to locking of the trial database so that analyses are not data-driven or selectively reported.²⁷ Following on from our practice for the ENOS, TARDIS and RIGHT-2 trials,²⁸⁻³⁰ we include not just information on the primary publication (PES vs. no sham) but also detail additional planned secondary publications, including baseline characteristics, health economics, swallowing up to one year and cognition up to one year. The SAP structure follows that published in 2017 by Gamble *et al.*^{31,32}

Supporting information

Additional supporting information may be found in the online version of this article with the accompanying Supplements.

Supplement:

1. Statistical analysis plan.

2. Health economics analysis plan.
3. Description of main results publication with follow-up to 90 days (day 90 was the original final outcome assessment).
4. Description of baseline data publication including demographics and stroke and dysphagia presentation.
5. Description of health economics to 90 days publication. (A similar format will be used out to day 365 if performed.)
6. Description of planned swallowing/dysphagia publication with follow-up to one year.
7. Description of planned cognition/dementia publication with follow-up to one year.
8. Description of planned responder analysis publication.

Other planned analyses/publications include:

1. Qualitative assessment of PES delivery.
2. Assessment of trial logistics, delivery and challenges.
3. Calculating swallowing and eating scales from clinical information.
4. Prevalence and use of oral trials of fluids/food.

DISCUSSION

PhEAST is the largest randomised trial of PES to date for any indication and is designed to provide definitive evidence on effectiveness and safety in patients with tube-dependent post-stroke dysphagia. Indeed, it may be the largest randomised trial ever in post-stroke dysphagia. Recruitment will complete at the end of October 2026 with a target of 650 participants. We anticipate completing day 90 follow-ups in quarter 1 2027 and presenting the main results at the European Stroke Organisation Conference (Vienna) in May 2027. Follow-ups at days 180 and 365 will complete later in 2027 with intended presentation at the International Stroke Conference (Los Angeles) in February 2028.

ACKNOWLEDGEMENTS

We thank the patients who joined the trial and their relatives for supporting enrolment; NIHR research delivery network staff who screened, recruited and treated patients; and speech & language therapists who screened and followed-up patient outcomes at day 14. PMB is Stroke Association Professor of Stroke Medicine and is an Emeritus NIHR Senior Investigator.

STATEMENT OF ETHICS

Study approvals were obtained in the UK from the national research ethics committee (REC): East of England - Essex (approval 21/EE/0252, date 6 Dec 2021), Health Research Agency (7 Jan 2022), Health Research Agency Wales (7 Jan 2022), and Scotland (21/SS/0075, 9 Feb 2022). In Austria, approval was given by the Lower Austrian Ethics Commission (GS1-EK-3/201-2021). In Denmark, approval was given by the Medical Research Ethics Committee (MREC) (16-0302-7). All participants or a relative gave written informed consent/consentee assent.

CONFLICT OF INTEREST

- LJW, IM, CR, JKB, SG, CJCS, GS, AB, CL, KMC, MJ, AAM, PMB – were funded, at least in part, by the HTA PhEAST grant (NIHR HTA: NIHR132016).
- SM: is employed by Phagenesis Ltd as a Senior Clinical Scientist
- SH: is Chief Scientific Officer for Phagenesis Ltd
- PMB: has previously consulted for Phagenesis Ltd (consultancy ceased in 2022) and been involved in running earlier studies of PES. Is a member of the European Stroke Organisation post-stroke dysphagia guidelines group.
- The remaining authors report no relevant conflicts of interest.

FUNDING SOURCES

Funding from the NIHR Health Technology Assessment (NIHR132016); the funder had no role in the design, data collection, data analysis, and reporting of this study. Phagenesis Ltd provide the devices (base station and treatment catheters) and training for their use; the company commented on the SAP but did not decide on contents.

AUTHOR CONTRIBUTIONS

- PMB, SH, TJE, AAM, MJ, NS, LFE, HKI, KM are co-investigators contributing to conceptualisation, funding acquisition, methodology, supervision, formal analysis and writing.
- LJW, IIM, CR, JKB, GW, SG, CJCS, GS, AB, CL, KMC, were/are trial management committee members contributing to trial administration, investigation, data curation, software, validation and statistical analysis.
- CR, MJ are health economists and wrote the HEAP.
- JPA, KK are medical monitors.
- SM contributed resources and writing.
- PMB wrote the original draft of this manuscript; all other authors contributed to review and editing.

DATA AVAILABILITY

PhEAST SAP

11/08/2026

Version 0.871

Once the trial is completed and published, data from PhEAST will be added to trial-level and individual patient data (IPD) meta-analyses of PES in acute/subacute stroke.^{4,14} IPD will be made available to the 'virtual international stroke trials archive' (VISTA)³³ and subsequently over the web, as with the International Stroke Trial.³⁴

REFERENCES

1. Cohen DL, Roffe C, Beavan J, et al. Post-stroke dysphagia: A review and design considerations for future trials. *Int J Stroke* 2016; 11(4): 399-411.
2. Labeit B, Michou E, Hamdy S, et al. The assessment of dysphagia after stroke: state of the art and future directions. *Lancet Neurol* 2023; 22(9): 858-70.
3. Labeit B, Michou E, Trapl-Grundschober M, et al. Dysphagia after stroke: research advances in treatment interventions. *Lancet Neurol* 2024; 23(4): 418-28.
4. Bath PM, Lee HS, Everton LF. Swallowing therapy for dysphagia in acute and subacute stroke. *Cochrane Database Syst Rev* 2018; 10(10): CD000323.
5. NICE. Transcutaneous neuromuscular electrical stimulation for oropharyngeal dysphagia in adults: NICE, 2018.
6. Hamdy S, Aziz Q, Rothwell JC, et al. The cortical topography of human swallowing musculature in health and disease. *Nat Med* 1996; 2(11): 1217-24.
7. Hamdy S, Rothwell JC, Aziz Q, Singh KD, Thompson DG. Long-term reorganization of human motor cortex driven by short-term sensory stimulation. *Nat Neurosci* 1998; 1(1): 64-8.
8. Hamdy S, Aziz Q, Rothwell J, et al. Explaining oropharyngeal dysphagia after unilateral hemispheric stroke. *The Lancet* 1997; 350(9097): 686-92.
9. Suntrup S, Teismann I, Wollbrink A, et al. Pharyngeal electrical stimulation can modulate swallowing in cortical processing and behavior - magnetoencephalographic evidence. *Neuroimage* 2015; 104: 117-24.
10. Fraser C, Power M, Hamdy S, et al. Driving plasticity in human adult motor cortex is associated with improved motor function after brain injury. *Neuron* 2002; 34(5): 831-40.
11. Jefferson S, Mistry S, Michou E, Singh S, Rothwell J, Hamdy S. Reversal of a virtual lesion in human pharyngeal motor cortex by high frequency contralesional brain stimulation. *Gastroenterology* 2009; 137(3): 841-9.
12. Jayasekaran V, Singh S, Tyrrell P, et al. Adjunctive functional pharyngeal electrical stimulation reverses swallowing disability after brain lesions. *Gastroenterology* 2010; 138(5): 1737-46.
13. Vasant D, Michou E, Tyrrell P, et al. OC-063 Pharyngeal Electrical Stimulation (pes) In Dysphagia Post-acute Stroke: A Double blind, Randomised Trial. *Gut* 2014; 63(Supplement 1): A31.
14. Scutt P, Lee H, Hamdy S, Bath P. Pharyngeal Electrical Stimulation for Treatment of Poststroke Dysphagia: Individual Patient Data Meta-Analysis of Randomised Controlled Trials. *Stroke Research and Treatment* 2015; 2015(429053): 8.
15. Bath PM, Scutt P, Love J, et al. Pharyngeal Electrical Stimulation for Treatment of Dysphagia in Subacute Stroke: A Randomized Controlled Trial. *Stroke* 2016; 47(6): 1562-70.
16. Bath PM, Woodhouse LJ, Suntrup-Krueger S, et al. Pharyngeal electrical stimulation for neurogenic dysphagia following stroke, traumatic brain injury or other causes: Main results from the PHADER cohort study. *EClinicalMedicine* 2020; 28: 100608.
17. Harvey RL, Smith R, Bathula R, et al. Pharyngeal electrical stimulation to treat dysphagia in acute stroke: learnings from cases in the PhEED clinical trial. *J Rehabil Med* 2025; 57: jrm43538.

18. **NICE. Pharyngeal electrical stimulation for neurogenic dysphagia.** London: National Institute for Health and Care Excellence, 2024.
19. **Suntrup S, Marian T, Burchard Schroder J, et al. Electrical pharyngeal stimulation for dysphagia treatment in tracheotomized stroke patients: a randomized controlled trial. *Intensive Care Medicine* 2015; 41(9): 1629-37.**
20. **Dziewas R, Stellato R, van der Tweel I, et al. Pharyngeal electrical stimulation for early decannulation in tracheotomised patients with neurogenic dysphagia after stroke (PHAST-TRAC): a prospective, single-blinded, randomised trial. *Lancet Neurol* 2018; 17(10): 849-59.**
21. **Dziewas R, Michou E, Trapl-Grundschober M, et al. European Stroke Organisation and European Society for Swallowing Disorders guideline for the diagnosis and treatment of post-stroke dysphagia. *Eur Stroke J* 2021; 6(3): LXXXIX-CXV.**
22. **Intercollegiate Stroke Working Party. National Clinical Guideline for Stroke for the UK and Ireland.** London: Royal College Physicians, 2023.
23. **Prabhakaran S, Gonzalez NR, Zachrison KS, et al. 2026 Guideline for the Early Management of Patients With Acute Ischemic Stroke: A Guideline From the American Heart Association/American Stroke Association. *Stroke* 2026.**
24. **Benfield JK, Everton LF, Woodhouse LJ, et al. Pharyngeal electrical stimulation versus none for post stroke dysphagia: rationale, design and protocol for the parallel group superiority pharyngeal electrical stimulation for acute stroke dysphagia trial (PhEAST) (ISRCTN98886991). *Cerebrovasc Dis Extra* 2026: 1-13.**
25. **Crary MA, Mann GD, Groher ME. Initial psychometric assessment of a functional oral intake scale for dysphagia in stroke patients. *Arch Phys Med Rehabil* 2005; 86(8): 1516-20.**
26. **Everton LF, Benfield JK, Hedstrom A, et al. Psychometric assessment and validation of the dysphagia severity rating scale in stroke patients. *Sci Rep* 2020; 10(1): 7268.**
27. **Collins R, MacMahon S. Reliable assessment of the effects of treatment on mortality and major morbidity, I: clinical trials. *Lancet* 2001; 357(9253): 373-80.**
28. **Bath PM, Houlton A, Woodhouse L, et al. Statistical analysis plan for the 'Efficacy of Nitric Oxide in Stroke' (ENOS) trial. *Int J Stroke* 2014; 9(3): 372-4.**
29. **Bath PM, Robson K, Woodhouse LJ, et al. Statistical analysis plan for the 'Triple Antiplatelets for Reducing Dependency after Ischaemic Stroke' (TARDIS) trial. *Int J Stroke* 2015; 10(3): 449-51.**
30. **Scutt P, Appleton JP, Dixon M, et al. Statistical analysis plan for the 'Rapid Intervention with Glyceryl trinitrate in Hypertensive stroke Trial-2 (RIGHT-2)'. *Eur Stroke J* 2018; 3(2): 193-6.**
31. **Gamble C, Krishan A, Stocken D, et al. Guidelines for the Content of Statistical Analysis Plans in Clinical Trials. *JAMA* 2017; 318(23): 2337-43.**
32. **Hemming K, Kearney A, Gamble C, et al. Prospective reporting of statistical analysis plans for randomised controlled trials. *Trials* 2020; 21(1): 898.**
33. **Ali M, Bath PM, Curram J, et al. The Virtual International Stroke Trials Archive. *Stroke* 2007; 38(6): 1905-10.**
34. **Sandercock PA, Niewada M, Czlonkowska A, International Stroke Trial Collaborative G. The International Stroke Trial database. *Trials* 2011; 12(1): 101.**
35. **Enos Trial Investigators. Efficacy of nitric oxide, with or without continuing antihypertensive treatment, for management of high blood**

pressure in acute stroke (ENOS): a partial-factorial randomised controlled trial. *Lancet* 2015; 385(9968): 617-28.

36. Bath PM, Woodhouse LJ, Appleton JP, et al. Antiplatelet therapy with aspirin, clopidogrel, and dipyridamole versus clopidogrel alone or aspirin and dipyridamole in patients with acute cerebral ischaemia (TARDIS): a randomised, open-label, phase 3 superiority trial. *Lancet* 2018; 391(10123): 850-9.

37. Bath PM, Scutt P, Blackburn DJ, et al. Intensive versus Guideline Blood Pressure and Lipid Lowering in Patients with Previous Stroke: Main Results from the Pilot 'Prevention of Decline in Cognition after Stroke Trial' (PODCAST) Randomised Controlled Trial. *PLoS One* 2017; 12(1): e0164608.

38. Sprigg N, Flaherty K, Appleton JP, et al. Tranexamic acid for hyperacute primary IntraCerebral Haemorrhage (TICH-2): an international randomised, placebo-controlled, phase 3 superiority trial. *Lancet* 2018; 391(10135): 2107-15.

39. Right-Investigators. Prehospital transdermal glyceryl trinitrate in patients with ultra-acute presumed stroke (RIGHT-2): an ambulance-based, randomised, sham-controlled, blinded, phase 3 trial. *Lancet* 2019; 393(10175): 1009-20.

40. Weir CJ, Lees KR. Comparison of stratification and adaptive methods for treatment allocation in an acute stroke clinical trial. *Stat Med* 2003; 22: 705-26.

41. Hernández AV, Steyerberg EW, Butcher I, et al. Adjustment for strong predictors of outcome in traumatic brain injury trials: 25% reduction in sample size requirements in the IMPACT study. *J Neurotrauma* 2006; 23(9): 1295-303.

42. Optimising the Analysis of Stroke Trials C, Gray LJ, Bath PM, Collier T. Should stroke trials adjust functional outcome for baseline prognostic factors? *Stroke* 2009; 40(3): 888-94.

43. Gray LJ, Collier T, Bath PMW, on behalf of the OAST Collaboration. Calculation of sample size for stroke trials assessing functional outcome: comparison of binary and ordinal approaches. European Stroke Conference; 2008; Nice, France; 2008.

44. Lees KR, Bath PMW, Schellinger PD, et al. Contemporary outcome measures in acute stroke research: choice of primary outcome measure *Stroke* 2012; 43(4): 1163-70.

45. Schellinger PD, Bath PMW, Lees K, et al. Assessment of additional endpoints relevant to the benefit of patients after stroke - what, when, where, in whom. *Int J Stroke* 2012; 7(3): 227-30.

46. Bath PMW, Woodhouse L, Scutt P, et al. Management of high blood pressure in acute stroke: Efficacy of Nitric Oxide in Stroke (ENOS), a partial-factorial randomised controlled trial. *Lancet* 2015; 385(9968): 617-28.

47. Woodhouse LJ, Scutt P, Hamdy S, et al. Route of Feeding as a Proxy for Dysphagia After Stroke and the Effect of Transdermal Glyceryl Trinitrate: Data from the Efficacy of Nitric Oxide in Stroke Randomised Controlled Trial. *Transl Stroke Res* 2018; 9(2): 120-9.

48. Wilkinson G, Everton LF, Krishnan K, Benfield J, Hamdy S, Bath PM. Mode of nutrition as a reflection of swallowing ability in acute and sub-acute stroke: Validation of a bedside tool. *J Stroke Cerebrovasc Dis* 2025; 34(12): 108484.

49. Cichero JA, Lam P, Steele CM, et al. Development of International Terminology and Definitions for Texture-Modified Foods and Thickened Fluids Used in Dysphagia Management: The IDDSI Framework. *Dysphagia* 2017; 32(2): 293-314.

50. Steele CM, Namasivayam-MacDonald AM, Guida BT, et al. Creation and Initial Validation of the International Dysphagia Diet Standardisation Initiative Functional Diet Scale. *Arch Phys Med Rehabil* 2018; 99(5): 934-44.
51. Belafsky PC, Mouadeb DA, Rees CJ, et al. Validity and reliability of the Eating Assessment Tool (EAT-10). *Ann Otol Rhinol Laryngol* 2008; 117(12): 919-24.
52. Jenkinson C, Fitzpatrick R, Crocker H, Peters M. The Stroke Impact Scale: validation in a UK setting and development of a SIS short form and SIS index. *Stroke* 2013; 44(9): 2532-5.
53. Zung WW, Richards CB, Short MJ. Self-rating depression scale in an outpatient clinic. Further validation of the SDS. *Arch Gen Psychiatry* 1965; 13(6): 508-15.
54. Quinn TJ, Dawson J, Lees JS, Chang TP, Walters MR, Lees KR. Time spent at home poststroke: "Home-time" a meaningful and robust outcome measure for stroke trials. *Stroke* 2008; 39: 231-3.
55. EMEA. Points to Consider on Multiplicity Issues in Clinical Trials. *EMA* 2002.
56. Schulz KF, Grimes DA. Multiplicity in randomised trials II: subgroup and interim analyses. *Lancet* 2005; 365(9471): 1657-61.
57. Wilkinson G, McLaughlin C, Rehman H, Hamdy S, Bath PM. Treatment of post-stroke dysphagia with pharyngeal electrical stimulation: the multicentre assessment of pharyngeal electrical stimulation (MAPS) Clinical Registry. *Submitted* 2026.
58. Kenedi H, Campbell-Vance J, Reynolds J, et al. Implementation and Analysis of a Free Water Protocol in Acute Trauma and Stroke Patients. *Crit Care Nurse* 2019; 39(3): e9-e17.
59. Rosenbek J, Robbins J, Roecker E, Coyle J, Wood J. A penetration-aspiration scale. *Dysphagia* 1996; 11(2): 93-8.
60. Wardlaw JM, Doubal F, Brown R, et al. Rates, risks and routes to reduce vascular dementia (R4vad), a UK-wide multicentre prospective observational cohort study of cognition after stroke: Protocol. *Eur Stroke J* 2021; 6(1): 89-101.
61. Galovic M, Stauber AJ, Leisi N, et al. Development and Validation of a Prognostic Model of Swallowing Recovery and Enteral Tube Feeding After Ischemic Stroke. *JAMA Neurol* 2019.
62. Scutt P, Lee HS, Hamdy S, Bath PM. Pharyngeal Electrical Stimulation for Treatment of Poststroke Dysphagia: Individual Patient Data Meta-Analysis of Randomised Controlled Trials. *Stroke Res Treat* 2015; 2015: 429053.
63. Ali M, Bath PMB, Davis SM, et al. The virtual international stroke trials archive (VISTA). *Stroke* 2007; 38: 1905-10.
64. Curtis L, Burns A. Unit Costs of Health and Social Care 2023. Canterbury: Personal Social Services Research Unit, 2023.
65. White IR, Horton NJ, Carpenter J, Pocock SJ. Strategy for intention to treat analysis in randomised trials with missing outcome data. *BMJ (Clinical research ed)* 2011; 342: d40.
66. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011; 30(4): 377-99.
67. Lyden P, Brott T, Tilley B, et al. Improved reliability of the NIH Stroke Scale using video training. NINDS TPA Stroke Study Group. *Stroke* 1994; 25(11): 2220-6.
68. Wei L, Lachin J. Two-Sample Asymptotically Distribution-Free Tests for Incomplete Multivariate Observations. *Journal of the American Statistical Association* 1984; 79(387): 653-61.

69. Lachin JM. Applications of the Wei-Lachin multivariate one-sided test for multiple outcomes on possibly different scales. *PLoS One* 2014; 9(10): e108784.
70. Wardlaw JM, Woodhouse LJ, Mhlanga, II, et al. Isosorbide Mononitrate and Cilostazol Treatment in Patients With Symptomatic Cerebral Small Vessel Disease: The Lacunar Intervention Trial-2 (LACI-2) Randomized Clinical Trial. *JAMA Neurol* 2023; 80(7): 682-92.
71. Bamford J, Sandercock P, Dennis M, Burn J, Warlow C. Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet* 1991; 337(8756): 1521-6.
72. Dhillon PS, Butt W, Podlasek A, et al. Effect of proximal blood flow arrest during endovascular thrombectomy (ProFATE): Study protocol for a multicentre randomised controlled trial. *Eur Stroke J* 2023; 8(2): 581-90.
73. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *Canadian Medical Association Journal* 2005; 173(5): 489-95.
74. Cumming TB, Blomstrand C, Bernhardt J, Linden T. The NIH stroke scale can establish cognitive function after stroke. *Cerebrovascular diseases (Basel, Switzerland)* 2010; 30(1): 7-14.
75. Nasreddine ZS, Phillips NA, Bédirian V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society* 2005; 53(4): 695-9.
76. Tombaugh TN, Kozak J, Rees L. Normative data stratified by age and education for two measures of verbal fluency: FAS and animal naming. *Arch Clin Neuropsychol* 1999; 14(2): 167-77.
77. Desmond DW, Tatemichi TK, Hanzawa L. The telephone interview for cognitive status (TICS): Reliability and validity in a stroke sample. *Int J Geriatr Psychiatry* 1994; 9: 803-7.
78. Burns A, Harrison JR, Symonds C, Morris J. A novel hybrid scale for the assessment of cognitive and executive function: The Free-Cog. *Int J Geriatr Psychiatry* 2021; 36(4): 566-72.
79. Brooke P, Bullock R. Validation of a 6 item cognitive impairment test with a view to primary care usage. *Int J Geriatr Psychiatry* 1999; 14(11): 936-40.
80. Zung WWK. A self-rating depression scale. *Arch Gen Psychiatry* 1965; 12: 63-70.

SUPPLEMENT 1

STATISTICAL ANALYSIS PLAN – PHARYNGEAL ELECTRICAL STIMULATION FOR ACUTE STROKE DYSPHAGIA TRIAL (PhEAST)

Authors: Philip M Bath, Lisa J Woodhouse, Alan M Montgomery.

ABBREVIATIONS

ADE	Adverse device effect
AE	Adverse event
BI	Barthel index
BLR	Binary logistic regression
CF	Informed consent form
CI	Chief Investigator overall
CPHR	Cox proportional hazards regression
CRF	Case Report Form
DAP	Data Analysis Plan
D/D	Death or discharge
DMC	Data Monitoring Committee
DSRS	Dysphagia severity rating scale
EAT-10	Eating assessment tool
eCRF	Electronic case record form
EDAR	Eating and drinking with acknowledged risk
EOT	End of Trial
EQ-VAS	EuroQuality of Life-visual analogue scale
EQ-5D-5L	EuroQoL-five dimensions-5 levels
FEES	Fibreoptic endoscopic evaluation of swallowing
FOIS	Functional oral intake scale
FSS	Feeding status scale
FWP	Free water protocol
GCP	Good Clinical Practice
GCS	Glasgow coma scale
HTA	Health Technology Assessment
ICH	Intracerebral haemorrhage
IDDSI	International dysphagia diet standardisation initiative
IDDSI-FDS	IDDSI-functional diet scale
IQCODE	Informant questionnaire on cognitive decline in the elderly
IS	Ischaemic stroke
ITT	Intention-to-treat
J-tube	Jejunostomy tube
LPFV	Last participant first visit
LPLV	Last participant last visit
mA	Milliampere
MHRA	Medicines and Healthcare products Regulatory Agency
mITT	Modified intention-to-treat
MLR	Multiple linear regression
MMSE	Mini-mental state examination
MoCA	Montreal cognitive assessment
mRS	Modified Rankin scale
NGT	Nasogastric tube
NJT	Nasojejunal tube
NHS	National Health Service

PhEAST SAP

11/08/2026

Version 0.871

NIHR	National Institute for Health and care Research
NIHSS	National Institutes of Health stroke scale
NMES	Neuro-muscular electrical stimulation
OLR	Ordinal logistic regression
OT	Occupational therapist
PAS	Penetration aspiration scale
PEG	Percutaneous endoscopic gastrostomy
PEJ	Percutaneous endoscopic jejunostomy
PES	Pharyngeal electrical stimulation
PI	Principal Investigator at a local centre
PIS	Participant Information Sheet
PLR	Personal legal representative (health)
PP	Per protocol
PSD	Post stroke dysphagia
REC	Research Ethics Committee
R&D	Research and Development department
RIG	Radiologically-inserted gastrostomy
RMR	Repeated measures regression
rTMS	Repetitive transcranial magnetic stimulation
SADE	Serious adverse device effect
SAE	Serious adverse event
SAR	Serious adverse reaction
SLT	Speech & language therapist
SPC	Summary of Product Characteristics
SWAT	Study within a trial
TMG	Trial Management Group
TSC	Trial Steering Committee
USADE	Unexpected serious adverse device effect
VFS	Videofluoroscopy
ZDS	Zung depression scale

SECTION 1. ADMINISTRATIVE INFORMATION

1 Title and trial registration

1a Title: Pharyngeal electrical stimulation for acute stroke dysphagia trial

Acronym: PhEAST

1b Registration: ISRCTN98886991. IRAS project number: 304658. Research ethics: England 21/EE/0252 (date 06/12/2021), Scotland 21/SS/0075 (date 09/02/2022). WHO U1111-1273-9942. University of Nottingham sponsor protocol 21068

2 SAP version: 1.0 (15 July 2026)

3 Protocol version: 11.0 (15 January 2026)

4 SAP revisions

4a Revision history: None, this is the first version.

4b Justification for each revision: Not relevant.

4c Timing of revisions: Not relevant.

5 Roles and responsibilities

Authors: Philip M Bath, Lisa J Woodhouse, Alan A Montgomery.

Responsible statisticians: Lisa J Woodhouse (senior statistician, blinded), Iris Mhlanga (statistician, unblinded), Alan A Montgomery (consultant statistician, blinded)

PhEAST SAP


11/08/2026

Version 0.871

Chief Investigator: Philip M Bath (blinded)

Contributors and roles: Philip M Bath, Lisa J Woodhouse, Iris Mhlanga, Alan Montgomery, for the PhEAST Investigators

6 Signatures

Role	Name	Signature	Date
Author, Statistician	Lisa J Woodhouse		7 Aug 2026
Author, Senior Statistician	Alan A Montgomery		7 Aug 2026
Author, Chief Investigator	Philip M Bath		7 Aug 2026

SECTION 2. INTRODUCTION

7 Background and rationale: See main text.

8 Objectives: To assess whether pharyngeal electrical stimulation (PES) is safe and efficacious at improving post-stroke dysphagia.

8a Primary Objective:

To assess whether 6 days of PES versus no PES (on top of best medical/usual care) accelerates return to oral intake of food and drink as assessed using the dysphagia severity rating scale (DSRS) with blinding to treatment.

8b Secondary Objectives:

To assess whether PES with usual care versus usual care:

- Improves swallowing and feeding.
- Reduces pneumonia, antibiotic usage, dependence on PEG/RIG feeding, and hospital length of stay. †
- Reduces dependence and disability.
- Is effective in predefined subgroups including by age, sex, cause (stroke type, severity), timing of intervention – see full list in section 27f. †
- Participant subgroups predict response to PES.
- Improves quality-of-life.
- Is cost effective.
- Reduces cognitive impairment.

† Some of objectives are driven, in part, by NICE questions.¹⁸ NICE refer to PhEAST as part of their review and guidance.

SECTION 3. STUDY METHODS

9 Trial design

PhEAST is an international prospective randomised open-label blinded-endpoint (PROBE) parallel-group superiority phase-4 effectiveness trial of PES versus no PES in 800 hospitalised patients with acute/subacute PSD. All participants receive best medical/usual care. Data collection is performed via a secure internet site with real-time data validation.

10 Randomisation

Allocate to PES : no PES 1:1 using minimisation, with 95/5 random element to reduce predictability, on country (non-UK vs UK), age (<75/75+), sex, DSRS (<12/12), impairment (NIHSS <15/15+), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior) and time to randomisation (<15/15+ days). (The minimisation-randomisation system is that used in our prior trials: ENOS, TARDIS, PODCAST, TICH-2, RIGHT-2.³⁵⁻³⁹) Including these minimisation variables as covariates in multivariable regression models improves precision of estimated between-group effects and so increases statistical power.⁴⁰

Investigators (doctors, research nurses-coordinators/healthcare professionals, speech and language therapists (SLT) (or occupational therapists [OT] in Denmark may enrol participants. Randomised treatment assignment will occur when essential baseline data are entered into the trial computer system by investigators. As such, allocation is concealed from investigators up to the time that they have screened, consented and collected and completed entry of key baseline data into the trial database.

11 Sample size/power considerations

11a Sample size calculation: The primary outcome, dysphagia severity rating scale

(DSRS, *S Table 1.1*),^{12,26} will be compared between PES and no PES using multiple linear regression with adjustment for minimisation variables. The null hypothesis (H_0) is that PES does not alter DSRS at day 14 in participants with PSD. The alternative hypothesis (H_1) is that PES alters DSRS at days 14 and 90 in participants with PSD.

S Table 1.1: DSRS and subscales including old 'food' consistencies and new IDDSI levels.^{12,26}

Score	Fluids	Score	Diet	Score	Supervision
4	No oral fluids	4	Non oral feeding	4	No oral feeding
3	IDDSI level 4 – extremely thick	3	IDDSI level 4 - pureed diet or level 5 – minced & moist	3	Therapeutic feeding (SALT/trained staff)
2	IDDSI level 3 – moderately thick	2	IDDSI level 6 – soft & bite sized	2	Feeding by third party (untrained)
1	IDDSI level 1 – slightly thick or level 2 mildly thick	1	IDDSI level 7 - easy to chew	1	Eating with supervision
0	IDDSI level 0 - thin	0	IDDSI level 7- regular	0	Eating independently

Patients on 15 teaspoons of intake or less per day (of any consistency) are given a DSRS score of 11 (4 – fluid scale; 4 - diet scale; 3 – supervision scale, such limited trials to be considered almost NBM.)
If more than 15 teaspoons, are given per day (any consistency), oral trials are scored from the fluid and diet subscales (i.e. 3 onwards) and can be either trials of food or fluid or trials of food and fluids.
DSRS supervision score 3 is always chosen when a patient is on limited or consistent oral trials and still requires NG/ PEG tube, i.e. although having oral intake there remains some dependency on tube feeding.

Assuming 1:1 randomisation; alpha 5% (two-tailed); power 90%; DSRS difference $\Delta=1.2$ (this target difference is above the lower level of 0.3 of the estimated minimal clinically important difference²⁶); standard deviation 5.0 (*Table 3*); losses 3% (greater than seen in previous PES trials); crossovers 3%; sample rounded up; a sample size of $N=800$ is needed (PES $n=400$, control $n=400$). The sample size estimation used a standard t test sample size formula. The assumptions are based on pilot trials and STEPS.¹²⁻¹⁵ We and others have shown that adjustment for covariates improves statistical power⁴¹⁻⁴³ or reduces sample size; however, we have not taken account of this in the above sample size calculation since the relevance of these findings to analysis of DSRS remains unclear. Nevertheless, it is likely that covariate adjustment will improve statistical power so that the final power will probably be greater than assumed here.

This calculation does not take account of the subsequent change in analysis plan which will use DSRS assessments at both days 14 and 90 with analysis using repeated measures regression; see section 26a below for further information.

11b Sensitivity calculations: The sample size varies by the difference in DSRS between PES and no PES (*S Table 1.2*).

S Table 1.2: Sample size for a variety of differences in mean DSRS between PES and no PES.

Data are shown for the original plan to compare DSRS between Pes and no PES using multiple linear regression. The illustrated mean differences all exceed the lower level of 0.3 of the estimated minimal clinically important difference.²⁶

Mean difference	1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7
Total trial size	1200	1000	800	700	600	500	500	400

S Table 1.3: Power of trial by achieved final sample size.

Data are shown for the revised plan (devised in 2025, approved by TSC in March 2026 during recruitment and prior to data lock and unblinding). 90% power is achieved with 636 participants.

Size (N)	822	795	769	742	716	689	663	636	610	583	557	530
Power (%)	96	95	94	94	93	92	91	90	89	87	86	85

12 Framework: PES will be tested versus no PES for superiority.

13 Statistical interim analyses and stopping guidance

13a Interim analyses: One formal interim analysis was planned to be performed to guide the independent Data Monitoring Committee (DMC) at the funder stop-go time-point occurring at 12 months after first enrolment. However, this was not performed because the recruitment of 65 participants at that stage was judged insufficient to justify formal analysis. Since performing the interim analysis later would not influence trial progression, we did not run the analysis.

13b Adjustments of significance level: There is no planned adjustment to the significance level; it remains at $P \leq 0.05$.

13c Stopping rules: The stopping rules for the Data Monitoring Committee regarding effectiveness were based on the combination of presence of 'proof beyond a reasonable doubt' and the likelihood that the results would inform clinical practice. The possible DMC recommendations at any assessment were:

1. Stop enrolment if the study is negative: statistical evidence that DSRS or fatal SAE rates are significantly higher in the PES than sham group ($p < 0.01$);
2. Stop enrolment if the study is positive: the combination of statistical evidence that DSRS is significantly lower in the PES than sham group "beyond reasonable doubt" and the overall trial results will lead to a change in clinical practice, e.g. by taking account of delta DSRS and evidence that at least some secondary dysphagia-swallowing or other related outcomes are also being benefitted, e.g. one or more of FOIS, IDDSI-FDS, EAT-10, FSS, death, length of hospital stay, pneumonia, antibiotic use.
3. Continue enrolment if the study is neutral: or if conditions 1 and 2 are not present.
4. Modify study design – if it appears that:
 - a. Sample size calculation assumptions were incorrect, e.g., if standard deviation exceeds 6.0;
 - b. Apparent study design aspects will lead to incorrect study conclusions;
 - c. Specific clinical procedures jeopardise the safe execution of the study.

The DMC performed their final assessment of unblinded data in October 2025 and recommended continuation of the trial (as per #3 above).

14 Timing of final analyses

Prior to each database lock, the trial managers/trial co-ordinators will chase outstanding data queries and the lock will take place in accordance with the documented data lock procedure once notification has been given to the trial programmer by the chief investigator. Both interim and final locks will be documented and primarily consist of the creation of a read-only copy of the live database, with each copy available to the trial statistician via the online data extract process. The main publication will focus on data up to day 90 as per the original funded plan.

A soft lock will be performed once the final participant has been recruited (LPFV). The trial statistician will share the results of analyses based on this SAP of the interim data with the chief investigator, co-chief investigator and deputy chief investigator; together they will draft the first version of the main publication and will

not participate further in participant- or analysis-related decision-making until data cleaning has completed following last participant last visit (LPLV). All other members of the trial team will remain blinded to the results.

The hard lock will be performed once the final participant has had their day 90 visit data collected (LPLV) and data cleaning has completed. The Main publication will be updated with the final results out to day 90 (*Supplement 3*).

Participants will remain in the trial until day 365; analyses for day 180 and 365 data will be performed once all participants have reached day 365 and data cleaning has completed. A secondary publication will focus on these longer-term follow-ups. (*Supplement 6*).

15 Timing of outcome assessments: Outcomes will be assessed blinded-to-allocation at days 14 (protocolised range 13-17), 90 (76-104), 180 (159-201), 365 (337-393) unless otherwise specified (*S Table 1.4*). Day 14 assessments are performed in hospital. Day 90,^{36,38,39,44-46} 180 and 365 are performed by central telephone (postal) follow-up. If the participant is an inpatient at day 90, 180 or 365, follow-up may be completed by the hospital site staff. If data are collected in hospital prior to day 90, this will be counted as day 14 if there is no existing day 14 data. If data are collected outside the protocolised windows, data will be assigned as follows:

- Day 14 missing and data collected prior to day 76 will be counted as day 14 data.
- Day 90 missing and data collected prior to day 159 will be counted as day 90 data.
- Day 180 missing and data collected prior to day 337 will be counted as day 365 data.

S Table 1.4: Timing of assessments.

Timing	Screen	Baseline	Day 1-9	Day 14 (13-17)	Discharge or death	Day 90 (83-97) †	Day 180 †	Day 365 †
Location	Hospital	Hospital	Hospital	Hospital	Hospital	Centrally ‡	Centrally ‡	Centrally ‡
Eligibility	+							
Consent/proxy consent	+							
Primary: DSRS, ^{12,26} blinded to treatment ¹	+	+		+		+	+	+
FOIS, ²⁵ mRS ⁴⁴	+	+		+		+	+	+
FSS, ^{47,48} IDDSI FDS, ^{49,50} EAT-10 ⁵¹		+		+		+	+	+
NGT/PEG/RIG <i>in situ</i>		+		+		+	+	+
NIHSS, Cog-4, GCS, CFS		+		+				
Stroke impact scale, ⁵² global		+		+		+	+	+
Cognition, e.g. MoCA MMSE TICS IQCODE		+		+		+	+	+
Randomisation		+						
Quality of life: EQ-5D-5L, EQ-VAS		+		+	+	+	+	+
BI, ZDS ⁵³		+		+		+	+	+
PES: threshold, tolerability, stimulation, duration, no. of catheters			x 6					
SAEs			<>					
Device-related (S)ADEs			<	>				
Targeted outcomes: chest infection, pneumonia, antibiotics			<	=	>			
Fatal SAEs			<	=	=	>		
All-cause mortality			+	+	+	+	+	+
Weight				+				
Disposition					+	+	+	+
Resource use					+	+	+	+
Home time ⁵⁴						+		

BI: Barthel index; CFS; clinical frailty scale; DSRS: dysphagia severity rating scale;²⁶ EAT-10: eating questionnaire; EQ-5D: EuroQoL 5-dimension 5-level; EQ-VAS: EuroQoL visual analogue scale; FOIS: functional oral intake scale (modified as per section 36a);²⁵ FSS; feeding status scale;^{47,48} GCS: Glasgow coma scale; Hosp: hospital; IDDSI: international dysphagia diet standardisation initiative; IDDSI-FDS: IDDSI-functional diet scale (modified as per section 36c); IQCODE: informant questionnaire on cognitive decline in the elderly; MMSE: mini-mental state examination; MoCA: Montreal cognitive assessment; NIHSS: National Institutes of Health stroke scale; QoL: quality of life; ZDS: Zung depression scale, abbreviated.⁵³

† Telephone (postal) follow-up

‡ If participant is an inpatient at day 90, 180 or 365, follow up can be completed by site staff in person.

Hospital discharge to/disposition: home, residential home, nursing home, hospital, death

SECTION 4. STATISTICAL PRINCIPLES

16 Levels of statistical significance

The result of the analysis of the primary outcome will be shown with a p-value, with $P \leq 0.05$ judged as significant. Otherwise, p-values will not be shown unless essential.

17 Adjustments for multiplicity

There will be no adjustment made for multiplicity; rather secondary analyses will be used for hypothesis generating.^{55,56}

18 Confidence intervals

The results of analyses and comparisons will be reported with 95% confidence intervals (95% CI).

19 Adherence and protocol deviations

19a Definition of adherence: Adherence will be assessed by examining the participant's 6 daily treatment forms as recorded on the REDCap database and based on evidence of treatment administration: threshold, tolerability, calculated stimulation and actual stimulation levels (maximum for each is 50 mA); and exposure assessed as number of treatments received (maximum is 6) and length of each treatment (maximum is 10 minutes).

Adherence will be considered sufficient if actual stimulation level is >20 mA, actual stimulation is the same as calculated stimulation level, six treatments are delivered and each treatment is given for >9 minutes 50 seconds.

19b Presentation of adherence: Adherence will be presented as:

- Mean (standard deviation) for each of threshold, tolerability, calculated stimulation and actual stimulation levels.
- Median [interquartile range] for each of number of treatments and their length.

19c Protocol violations: These are a major deviation from the trial protocol, for example where a participant is enrolled in spite of not fulfilling all the inclusion and exclusion criteria, or where deviations from the protocol could negatively affect participant safety, trial delivery or interpretation. Listed protocol violations are:

1. Enrolment without consent.
2. Enrolment but ineligible.
3. Non-reporting of primary outcome measure.
4. Non-reporting of serious adverse event (SAE), serious adverse device event (SADE), serious unanticipated adverse device event (SUADE).
5. PES treatment beginning after day 1, i.e. not started on day of or after randomisation.
6. Primary outcome (DSRS) at day 14 not performed between days 13 and 17.
7. Primary outcome (DSRS) at day 14 assessor not blinded to treatment assignment.
8. Other dysphagia therapy device(s) used during the trial period of days 0-90.
9. Any other major violation of the trial protocol.

All protocol violations must be reported immediately to the Chief Investigator, via the online electronic case report form. The CI will notify the Sponsor if a violation has an impact on participant safety or integrity of the trial data. The Sponsor will advise on appropriate measures to address the occurrence, which may include reporting of a serious GCP breach, internal audit of the trial and seeking counsel of the trial committees. Violations, deviations and device deficiencies will be presented in listings in the main publication (*Supplement 3*).

19d Protocol deviations: These are minor deviations from the protocol that affect the conduct of the trial in a minor way. This includes any deviation from the trial protocol that is not listed as a Protocol Violation. Deviations will be collected, via the

online electronic case report form. The list of deviations is not exhaustive and so there will be a drop-down menu as well as a free text box in the eCRF.

All protocol deviations must be reported immediately to the Chief Investigator via the online electronic case report form (eCRF). The CI will notify the Sponsor if a deviation or violation has an impact on participant safety or integrity of the trial data. The Sponsor will advise on appropriate measures to address the occurrence, which may include reporting of a serious GCP breach, internal audit of the trial and seeking counsel of the trial committees. Deviations will be presented in listings in the main publication (*Supplement 3*).

19e Device deficiencies: Apparent device deficiencies (whether of base station or catheters) must be reported on the relevant eCRF; these will be forwarded to the manufacturer for further assessment. Deficiencies will be presented in listings in the main publication (*Supplement 3*).

20 Analysis populations

All available data will be used and levels of missing data will be reported.

1. **Safety:** All randomised participants – primary safety analysis.
2. **Modified intention to treat (mITT):** All randomised participants with recorded DSRS data at days 14 and/or 90 analysed using repeated measures regression and without imputation – primary efficacy analysis.
3. **Intention to treat (ITT):** All randomised participants with recorded or imputed DSRS data at days 14 and 90 analysed using repeated measures regression and imputations made using regression imputation – sensitivity efficacy analysis.
4. **Per protocol set (PP):** All participants in the mITT set who are deemed to have no protocol violations that could interfere with the objectives of the study – sensitivity efficacy analysis.

Participants will be included or excluded from analyses of these populations as follows:

Excluded from all analyses

- Enrolled without valid consent.
- Enrolled with consent but no randomisation.

Excluded from per protocol analyses

- Enrolled but ineligible:
 - Already in another trial where a co-enrolment agreement is not in place.
 - FOIS >3.
 - Randomised <2 or >31 days.
 - Randomised to PES but did not receive it.
 - Randomised to no PES but did receive it.
- Dysphagia therapy device(s) used between randomisation and day 90.
- Received PES treatment late, i.e. day >1 after randomisation.
- Primary outcome not performed between days 13 to 17 and/or days 76 to 104.
- Non-reporting of primary outcome measure at both days 14 and 90.
- Received active treatment with another device prior to the 90 day follow-up.

SECTION 5. TRIAL POPULATION

21 Screening data: Sites will attempt to keep screening data in logs; where kept, these will be aggregated and presented.

22 Eligibility

22a Inclusion

1. Hospitalised adults.

2. Age ≥ 18 years.
3. Recent (2-31 days) ischaemic or haemorrhagic, anterior or posterior circulation, stroke (as diagnosed clinico-radiologically) at a stroke centre.
4. Clinical dysphagia with dependency on tube feeding defined as a functional oral intake scale (FOIS²⁵) score of:
 - i. Nothing by mouth, feeding by NGT/PEG,
 - ii. Tube dependent feeding with minimal attempts of food or liquids, or
 - iii. Tube dependent feeding with consistent oral intake of food or liquids.
5. Baseline DSRS supervision score of either 3 (requiring therapeutic feeding by SALT team; on oral trials) or 4 (No oral feeding) (Supplement 1, Table 2).
6. Conscious or semi-conscious defined as a NIHSS item 1a score of 0, 1 or 2 (where the patient requires repeated stimulation to arouse).

Women of childbearing age may be included since the treatment time is short (6 daily 10 minute sessions) with no residual effects.

22b Exclusion

1. Non-stroke dysphagia, e.g. due to traumatic brain haemorrhage, subarachnoid haemorrhage, brain tumour, Parkinson's disease, multiple sclerosis, severe dementia, head or neck cancer.
2. Pre-stroke dysphagia for whatever reason and whether resolved or not.
3. Pre-stroke dependency, i.e. modified Rankin scale (mRS⁴⁴) is 4 or 5.
4. NIHSS item 1a score of 2 (where the patient only responds to pain) or 3.
5. Ongoing or anticipated ventilation/intubation/tracheostomy. Patients who have been ventilated (and been decannulated if they required a tracheotomy) are eligible if they remain dysphagic.
6. Ongoing or planned treatment of dysphagia with other:
 - a) Forms of electrical / magnetic stimulation, e.g. neuromuscular electrical stimulation (NMES), transcranial direct-current stimulation (TDCS), repetitive transcranial magnetic stimulation (rTMS), Ampcare; or
 - b) Devices, e.g. expiratory muscle strength training (EMST), IQoro, IOPI, biofeedback using EMG electrodes, or chin tuck against resistance using a ball/chin depressor.
 - c) These exclusions apply for the duration of trial follow-up.
7. Malignant middle cerebral artery syndrome (although this typically presents before 4 days). Patients are eligible once stable after hemicraniectomy.
8. Pacemaker, cochlear implant or implantable cardioverter-defibrillator.
9. Need for $>35\%$ of oxygen.
10. Participants not likely to be in the treating hospital for at least 14 days (including the 6-day treatment period), i.e. expected to be repatriated to, or rehabilitated at, a non-participating organisation.
11. Two or more NGT tubes pulled out within the last week unless nasal bridle in place.
12. Investigator feels patient will not tolerate PES catheter.
13. Pregnancy if known at time of enrolment. A pregnancy test is not required.
14. Participating in another randomised controlled treatment trial for post-stroke dysphagia.
15. Palliative care.
16. Presence of a pharyngeal pouch.
17. Investigator believes dysphagia will be short-term, e.g. signs of impending recovery in swallowing (enrolment is reserved for patients whom the investigator considers are unlikely to resolve spontaneously over the next few days).
18. Participant is risk-feeding at time of screening.

23 Recruitment

Recruitment will be summarised in a CONSORT flow diagram showing recruitment, randomisation and follow-ups at days 14 and 90.

24 Withdrawal/follow-ups

24a Withdrawals: Withdrawals are considered as:

- Withdrawal from treatment (applies only to PES group).
- Withdrawal from individual follow-ups.
- Withdrawal from all follow-ups.
- Missed follow-ups.
- Lost to follow-up.

24b Timing of withdrawals: These will be categorised by days 14, 90, 180 and 365.

24c Presentation of withdrawals: The timing of withdrawals and losses will be summarised in the CONSORT flow diagram. They will be shown by treatment group in a Kaplan-Meier graph and rates compared using Cox proportional hazards regression.

25 Baseline patient characteristics

25a Listing of baseline characteristics: These will comprise demographics and clinical measures and are listed in *Supplement 3*.

25b Summarisation of baseline characteristics: Data will be shown as number (%), median [interquartile range] or mean (standard deviation), as appropriate.

SECTION 6. ANALYSIS

26 Outcome definitions

26a Outcomes

Primary outcome: The primary outcome is the 14-level dysphagia severity rating scale (DSRS), a composite measure of swallowing impairment comprising three ordered categorical subscales each with score 0-4 so total score is 0-12 (*Supplement Table 1.1*). For PhEAST, the DSRS is extended to include death with total score = 13, as we have done previously.^{15 57} Detailed guidance for scoring patients on oral trials was developed for PhEAST using the DSRS. These are: allocating a score of 11 for patients on minimal oral trials (≤ 15 teaspoons of any consistency in total per day) and scoring from the food and fluid sub-scales for patients on consistent oral trials (> 15 teaspoons of any consistency in total per day) alongside a score of 3 on the supervision scale. These same definitions were used to operationally define the difference between a FOIS of 2 and 3. These scores reflect the limitations of current scales to accurately capture scores for patients on oral trials which is a common approach used in the management of PSD. Although the FOIS is one of the few scales to include reference to oral trials, it was necessary to operationally define the wording used for the purpose of the study. Originally the intention was to assess the primary outcome at day 14 reflecting the design of earlier PES studies. However, post-stroke recovery paradigms value assessment of outcome at day 90 by which time effective interventions should have had a substantial effect. Additionally, PES studies find that DSRS continues to improve over 90 days whilst some recovery occurs in untreated participants. This is similar to recover trajectories of motor and speech impairments following stroke. Rather than just switching the timing of the primary outcome from day 14 to day 90, we decided to keep the first and add the second and use a repeated measures analysis; this will have the added effect of increasingly statistical power. This decision to change was made during recruitment, prior to data lock and masked to randomisation and is explained in protocol version 11 and the protocol publication.²⁴

To avoid observer bias, the primary outcome measure (DSRS) is assessed by a blinded: i) speech & language therapist at day 14 while the participant is in hospital; and ii) follow-up coordinator at day 90 by telephone; both are masked to treatment assignment. Treatment catheters will be removed before the day 14 assessment and replaced by a NGT, as necessary, to maintain blinding in hospital.

All-cause death is included in the extended DSRS for multiple reasons: i) death is a common outcome after severe stroke with dysphagia, e.g. following pneumonia resulting from aspiration; ii) to avoid missing a “kill or cure” effect; iii) to avoid excluding participants from analyses (and so maintain statistical power); iv) to avoid “missingness” which may be informative; and v) to anchor analyses. Many other scales used in stroke or dysphagia include death, e.g. modified Rankin scale, Barthel index, quality of life (EQ-5D), feeding status scale.

Patients with post-stroke dysphagia may receive fluids and food in spite of speech & language therapy advice against this. Specifically, some patients may wish to receive water to drink even if they have thin liquid aspiration, so-called free water protocol.⁵⁸ Equally, patients who are near to end-of-life may wish to feed even though they have liquid and/or food aspiration, known as “eating and drinking with acknowledged risk” (EDAR). (Of note, EDAR present at baseline is ineligible for the trial.) The main analyses described here will use the “professional” scores derived from SLT advice. A sensitivity analysis will use “pragmatic” scores derived from what the participant is actually doing. For example, a participant may have a professional SLT-scored DSRS of 4 (fluids 1, diet 2, supervision 1) but a pragmatic DSRS of 0 (fluids 0, diet 0, supervision 0), the latter reflecting participant request. The issue of using these latter scores is that they may make the effect of control/no PES look better so that DSRS scores in the two treatment groups, PES and no PES, appear more similar.

Secondary endpoints: These are listed by timing in *S Table 1.4*. Day 90/180/365 assessments will be made centrally by telephone blinded to baseline scores, treatment and in-hospital data (or by post if telephone does not work). If the participant is an inpatient at day 90, 180 or 365, the follow-up may be completed by the hospital site staff.

Note 1: These outcomes have all been sensitive to therapeutic change in previous studies.

Penetration/aspiration: The use of fiberoptic endoscopic evaluation of swallowing (FEES) or videofluoroscopy (VFS) is not mandated in the protocol but if they are carried out the penetration aspiration scale score⁵⁹ and date will be recorded via the eCRF.

Health economics: See *Supplement 2*: health economics analysis protocol (HEAP).

Safety endpoints: PES has an excellent safety record in previous trials¹⁸ whilst participants with PSD, who usually have severe stroke, will have multiple adverse events and SAEs. Hence, we will limit recording to:

- SAEs over 0-9 days.
- Procedure/device-related (S)AEs/SADEs/USADEs over days 0-14.
- Discontinuations due to (S)AEs.
- Fatal SAEs over days 10-90.

Events will be adjudicated blinded to participant information and treatment assignment.

Device deficiencies: Device deficiencies will be recorded and forwarded to the company for further investigation. These will be listed as related to:

- The base station
- The treatment catheter

Cognition: Post-stroke cognitive impairment (PSCI) and post-stroke dementia (PSD) follow stroke damage to cognitive neural pathways and/or the presence of concomitant small vessel disease and are common with rates up to 35% at 5 years.

However, studies into PSCI/PSD are complicated by multiple issues, notably: a tendency to recruit patients with milder stroke (who are easier to enrol); assessment of cognition in the presence of existing cognitive impairment or concurrent dysphasia; difficulty in follow up, e.g. because the patient is in a care home; or the patient has died. As a result, adequate assessment of cognition and its temporal trajectory in patients with severe ischaemic stroke (IS) or intracerebral haemorrhage (ICH) is often not performed.

We expect most patients with post-stroke dysphagia in the PhEAST trial will have severe stroke, and so the PhEAST cognition study will assess cognition at days 0 (baseline), day 14 (timing of main study primary outcome), day 90 (timing of main study secondary outcomes) and days 180 and 365 after randomisation. Dysphagia measures at day 180 and day 365 will be collected as well as the cognition measures. The measures used for the cognition study will be:

- Cognition (participant): DSM-5-7 level and DSM-5-4-level ordinal cognition scales,⁶⁰ t-MoCA, t-TICS, t-MMSE, semantic verbal fluency (animal naming), phonemic verbal fluency (letter F), clinical diagnosis of dementia (from participants or carers/informants).
- Cognition (from informant): IQCODE (separate informant consent for this).
- Dependency and disability (necessary for diagnosis of dementia): modified Rankin scale (mRS ⁴⁴), and Barthel index (BI).
- Frailty: clinical frailty index (CFI).
- Mood/depression (which complicates PSCI/PSD diagnosis): Zung depression scale.
- Health economics: EQ-5D-5L, EQ-VAS.
- Stroke impact scale.
- Recurrent stroke after index event.

Overall, the cognition study will provide information on cognition and its trajectory over the first year after severe stroke, a neglected research area and of considerable importance to this population and their family and carers. Additionally, it will enhance the main trial itself through providing extended follow-up information on swallowing.

26b Units: Units will be shown in tables.

26c Calculations/transformations: Quality of life (EQ-5D-5L) using UK weightings.

26d NICE: NICE list several key efficacy and safety outcomes to be assessed in research:¹⁸

- “Degree of aspiration” – we assess that as chest infection or pneumonia.
- “Change in severity of dysphagia” – we assess that as swallow impairment (DSRS).
- “Need for nasogastric and percutaneous endoscopic gastrostomy or jejunostomy feeding” – we assess that as need for any enteral tube (whether NGT, PEG, RIG, J-tube) and need for any permanent feeding tube (whether PEG, RIG, J-tube).
- “Device-related discomfort or injury” – we assess these through reporting of SADE, USADE or device deficiencies.

27 Analysis methods

27a Methods of analyses: Results will be tabulated as number (%), median [interquartile range] or mean (standard deviation) (*S Table 1.5*). Absolute (binary analyses) and relative measures of effect and 95% confidence intervals will be presented for each analysis. In tabulations, a worst score will be assigned at days 14, 90, 180 and 365 for people who die (e.g. DSRS=13, FOIS=0, mRS=6) to avoid losing participants in analyses (and so reduce statistical power), missing a “kill or cure” effect and to anchor analyses.

S Table 1.5: Summary of descriptive and analytical statistics

	Binary	Nominal	Time to event	Ordinal	Continuous
Central	N (%)	N (%)	N (%)	Median	Mean

tendency and distribution				[interquartile range]	(standard deviation)
Comparisons	Fisher's exact test/Chi-square (2x2)	Fisher's exact test/Chi-square (2x2, or rxc)		Mann-Whitney U	t-test (pooled)
Regression including adjustment (a)	Binary logistic regression (BLR, aBLR)	-	Cox proportional hazards regression (CPHR, aCPHR)	Ordinal logistic regression (OLR, aOLR)	Multiple linear regression (MLR, aMLR) Repeated measures regression (RMR, aRMR)

Analyses of Primary endpoint: DSRS will be analysed by adjusted repeated measures linear mixed effects regression based on data from days 14 and 90 with covariate adjustment. The model will include fixed effects for treatment group, time (categorical), and the treatment-by-time interaction, with baseline outcome value included as a covariate. A random intercept for participant will be included to account for within-subject correlation. An unstructured covariance matrix will be used to model the repeated measurements; alternative covariance structures will be explored in sensitivity analyses. Effect size will be reported as the average treatment effect with confidence intervals and p-value. The interaction between treatment effect and time will also be reported.

Analyses of secondary outcomes: Secondary outcomes will be compared using continuous repeated measures regression, ordinal repeated measures regression (e.g. FOIS), multiple linear regression (e.g. Barthel index), ordinal logistic regression (e.g. mRS), Cox proportional hazards regression (e.g. time to death) and binary logistic regression (e.g. discharge with PEG/RIG).

27b Covariate adjustment: Analyses will be adjusted for baseline DSRS, age, sex, impairment (NIHSS), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior) and time to randomisation (days). Non-binary variables will use raw data.

27c Assumption checking: Since the sample size is considered to be large, we will not test for normality, however plots of the residuals will be reviewed to check for potential violations.. The assumption of proportionality for ordinal logistic regression will be tested with the likelihood test and reported.

27d Alternative methods: We will still use OLR if the test of proportionality fails reporting odds ratio and confidence intervals.

27e Sensitivity analyses: The primary outcome of DSRS will be subject to sensitivity analyses as shown in **S Table 6**. The analyses will assess the impact of imputation of missing DSRS scores at day 14 and/or 90, not adjusting for minimisation and other important baseline variables, only including participants where no key protocol violations occurred (per protocol), accounting for free water administration or EDAR assessing the outcome only at day 14 or day 90, and excluding death.

S Table 6. Sensitivity analyses for the primary outcome, DSRS.

Primary outcome	Day	Population	Imputation	Analysis
With no imputation	14/90	mITT	No	aRMR
<i>Sensitivity analyses</i>				
With imputation	14/90	ITT	Yes	aRMR
Unadjusted	14/90	mITT	No	RMR
Per protocol (#20.4)	14/90	PP	No	aRMR
Accounting for FWP/EDAR	14/90	mITT	No	aRMR
With no imputation	14	mITT	No	aMLR
With no imputation, change	14	mITT	No	aMLR
With no imputation	90	mITT	No	aMLR
With no imputation, change	90	mITT	No	aMLR
Excluding death	14/90	mITT	No	aRMR

a: adjusted; EDAR: eating and drinking with acknowledged risk; FWP: free water protocol; ITT: intention-to-treat; mITT: modified intention-to-treat; PP: per protocol

27f Subgroup analyses: The primary outcome, DSRS, will be assessed in pre-specified subgroups based on minimisation factors using interaction tests to identify responder and non-responder subgroups. Between subgroup effect (95% CI) will be presented:

- Age (years): <75, ≥75
- Sex: female, male
- Reperfusion: yes, no
- Dysphagia severity rating scale (/12): <11, 11, 12
- Functional oral intake scale (/7): 1, 2/3
- Stroke severity, NIHSS (/42): <13, ≥13
- Stroke type: ischaemic, primary intracerebral haemorrhage
- Circulation: anterior, posterior
- Onset to randomisation (days): <12, 12-19, ≥19
- Lesion side: right, bilateral, left
- Stroke size: small, medium, large
- White matter disease: none, mild, moderate/severe
- Prior instrumental testing: yes, no
- Responder, DSRS change day 90-0: >-3.5, ≤-3.5
- Responder, DSRS day 90: <4, ≥4
- Experience with PES prior to trial: yes, no
- Site recruitment: >10, ≤10

Since the trial is powered to detect overall differences between groups rather than subgroup interactions, these analyses will be regarded as hypothesis generating. The importance of some of these subgroup analyses is highlighted by similar findings in PHAST-TRAC and PHADER that PES appears to more effective if treatment is started within the first month after stroke.^{16,20}

A subgroup analysis will be performed in the PES-only group to assess the relationship between stimulation levels (in milliamps) and effect on primary outcome at day 90. The analysis will follow that of PHADER.¹⁶

- Threshold, average: <10, 10-13, ≥13
- Stimulation, average: <30, 30-36, ≥36
- Tolerability, average: <35, 35-45, ≥45
- Delta, tolerability – threshold, average: <24, 24-33, ≥33

28 Missing data: Missing data will be imputed using multiple regression imputation for a sensitivity analysis of DSRS. The multiple imputation will be undertaken using

the PROC MI procedure in SAS, with 20 burn-in iterations and using the default setting for number of imputations. Data will be imputed at days 14 and/or 90 if missing. If participants have definitely died, they will be assigned a value of DSSR=13; if they are definitely alive but their DSSR is unknown, they will be assigned a DSRS value of 0-12 as determined by regression imputation.

29 Additional analyses: No further analyses planned.

30 Harms: These will be presented in Tables describing serious adverse events.

31 Statistical software: Statistical Analysis System (SAS) version 9.4 (or later), SAS Institute Incorporation, Cary, North Carolina.

SECTION 7. ADDITIONAL INFORMATION

32 Governance: The trial is funded by the National Institute for Health and Care Research (NIHR) Health Technology Assessment (HTA) Programme and sponsored by the University of Nottingham. Phagenesis Ltd, the device manufacturer, lend sites a base station and provide catheters and training for free. The trial is managed by a Trial Management Committee (TMC), supervised by a Trial Steering Committee (TSC) and overseen by an independent Data Monitoring Committee (DMC).

33 Minimising bias

Multiple approaches are taken in the design and execution of the trial to minimise bias:

- Central data registration with real-time on-line validation;
- Concealment of allocation;
- Adaptive randomisation/minimisation;
- Inclusion of patients enrolled in other studies (co-enrolment) where feasible;
- Blinded assessment of DSRS at day 14;
- Blinded central telephone (or postal) assessment of outcomes at days 90, 180, 365;
- Analysis by modified intention-to-treat and in pre-specified subgroups.

34 Study within a trial (SWAT)

The SWAT will investigate the effects of a package to improve awareness and communication with investigators to improve PES delivery and especially delivered current. Sites are cluster-randomised to the intervention vs control at ratio 1:1. The intervention comprises the Coordinating Centre's speech & language therapist contacting the treater who under-treated the participant and determining what happened and any root causes. The treater is reminded of the importance of providing adequate treatment in respect of stimulation current and treatment duration.

One interim analysis is planned at 200 PES participants (i.e. 400 total randomised) in the main trial. If there is strong evidence of an effect of the SWAT intervention on treatment current, the strategy showing the greatest treatment current would then be implemented for the remaining participants. Otherwise, the SWAT will continue until the end of the trial. Analyses will include descriptive statistics and between-group comparisons for each strategy using multivariate regression models.

Success criteria:

1. Increased treatment current over first three days in participants randomised to SWAT vs no-SWAT.

2. Increased proportion of participants with mean treatment current >20 mA in hospitals randomised to SWAT vs no-SWAT.
3. Increased proportion of participants who only require one PES catheter in hospitals randomised to SWAT vs no-SWAT.

35 Discrepancies between protocol and statistical analysis plan

Where there is a difference between the protocol (website and published versions) and this statistical analysis plan (SAP), the SAP takes precedence. This is particularly relevant to the change in analysis of the primary outcome.

36 Calculations/definitions

36a FOIS: This is a swallow impairment scale

Level	Item	Notes
1	Nothing by mouth	
2	Tube dependent with minimal attempts of food or liquid	Minimal is defined as ≤15 tea-spoons in total of any consistency per day
3	Tube dependent with consistent oral intake of food or liquid	Consistent is defined as >15 tea-spoons of any consistency per day in total (but not sufficient for nutritional purposes, i.e. still requires some element of tube feeding)
4	Total oral diet of a single consistency	
5	Total oral diet with multiple consistencies, but requiring special preparation or compensations	
6	Total oral diet with multiple consistencies without special preparation, but with specific food limitations	
7	Total oral diet with no restrictions	

36b FSS: The FSS assesses feeding status based on mode of nutrition. The scale is shown here with the scores as coded incorrectly in the REDCap database and the correct scores, as published,⁴⁸ which will be reported in publications.

Correct score	Score in REDCap	Level	Item (adapted from ⁴⁸)
0	1	Oral: Normal	- Regular oral diet (IDDSI L7/L7 easy chew). - Normal oral fluids (IDDSI L0). - Non-oral feeding route may be in situ but is not required to maintain nutrition.
1	2	Oral: Soft	- Modified oral diet (IDDSI L6 or below). - Oral fluids – normal or modified (IDDSI L0-L4). - Non-oral feeding route may be in situ but is not required to maintain nutrition.
2	3	Non-oral: NGT	Nasogastric tube feeding required to

			maintain nutrition. • May receive oral fluids – normal or modified (IDDSI L0-L4). • May receive tastes/oral trials of diet.
3	4	Non-oral: PEG/RIG/J-tube	• Percutaneous endoscopic gastrostomy feeding (or equivalent) required to maintain nutrition. • May receive oral fluids – normal or modified (IDDSI L0-L4). • May receive tastes/oral trials of diet.
4	5	Non-oral: IV/SC	• Non-oral supplemented fluid intake. • May receive oral fluids – normal or modified (IDDSI L0-L4). • May receive tastes/oral trials of diet.
5	6	Non-oral: nothing	• May receive tastes/free water protocol for comfort. • May be awaiting placement of PEG/NGT.
6	0	Died	• Due to any cause

36c IDDSI-FDS: Two IDDSI scores rate swallow impairment on the basis of intake of:⁴⁹

- Food : regular (i.e. normal) score =7; soft & bite sized =6; minced & moist =5; pureed =4; liquidised =3; nil by mouth = 2.
- Drinks : thin (i.e. normal) =0; slightly thick =1; mildly thick =2; moderately thick =3; extremely thick, score =4; nil by mouth =5.

From this, the overall swallow impairment is calculated using the IDDSI functional diet scale (IDDSI-FDS) score which ranges from 1 to 8.⁵⁰ We have modified the IDDSI-FDS score to include values less than 1 to allow scoring of nil by mouth, including minimal and consistent trials. Hence, the modified IDDSI-FDS (m-IDDSI-FDS) ranges from 0 (nil-by-mouth) to 8 (food 7, drinks 0, i.e. normal). Score of 0, 0.25, 0.5 and 0.75 allow differentiation of patients who are nil-by-mouth but may or not be taking trials of feeding:

- 0: NGT, PEG (G-tube), RIG, NJT or PEJ (J-tube) *in situ* with no trials of food or drinks, i.e. equivalent to FOIS=0.
- 0.25: NGT, PEG (G-tube), RIG, NJT or PEJ (J-tube) *in situ* with trials involving <=15 teaspoons of food or fluids (minimal trials), i.e. equivalent to FOIS=2.
- 0.5: NGT, PEG (G-tube), RIG, NJT or PEJ (J-tube) *in situ* with trials involving >15 teaspoons of food or fluids (consistent trials), i.e. equivalent to FOIS=3.
- 0.75: This is calculated as (food=4 AND drink=NA) OR (food=3 AND drink=4) OR (food=NA AND drink=3).

The modification maintains symmetry around the top-right / bottom-left axis.

IDDSI-FDS							m-IDDSI-FDS						
Food scores						Drink score	Food scores						Drink score
7	6	5	4	3	No food		7	6	5	4	3	2 = No food (tube +/- trials)	
3	2	1	X	X	X	No drink	3	2	1	0.75	0.5	0.5 or 0.25 or 0	5 = No drink (tube +/- trials)
4	3	2	1	X	X	4	4	3	2	1	0.75	0.5	4

PhEAST SAP

11/08/2026

Version 0.871

5	4	3	2	1	X	3	→	5	4	3	2	1	0.75	3
6	5	4	3	2	1	2		6	5	4	3	2	1	2
7	6	5	4	3	2	1		7	6	5	4	3	2	1
8	7	6	5	4	3	0		8	7	6	5	4	3	0

36d EAT-10:⁵¹ This 10-item scale assesses swallowing difficulties when swallowing is still possible. Each item scores from 0 = no problem to 4 = severe problem so that the maximum score is 10 x 4 = 40. Since the scale does not account for non-oral feeding, we have extended it by adding a nil by mouth score = 5 resulting in a maximum score of 50. This score includes participants who are on oral trials.

36e Death scores: Several scales already include a value for death, e.g. mRS=6, BI=-5 and EQ-5D health utility = 0. We have extended other scales to include death, usually with a value more extreme than any living value. The aim is to avoid: losing participants in analyses (and so maintain statistical power), missing a “kill or cure” effect and to anchor analyses.^{36,38,39,46} The death values are given here:

- Swallowing/eating: DSRs=13, FOIS=0, FSS=6, IDDSI-food=1, IDDSI-drink=6, IDDSI-FDS=-1, EAT-10=51
- Others: CFS=10, EQ-VAS=-1, NIHSS=43, ZDS=102.5

36f Global analyses: Analyses that include a global assessment of multiple scales are usually more efficient statistically. We will use the Wei-Lachin test to perform these for two groupings of scales:

- Swallowing/feeding: DSRs, FOIS, FSS, IDDSI-FDS and EAT-10.
- Stroke impact scale:⁵² 8 domains – strength, hand function, mobility, activities of daily living, memory, communication, handicap.

36d Composite of poor outcomes at death or discharge: DSRs>6, pneumonia in hospital, PEG *in situ* at discharge, died in hospital or length of stay >upper interquartile range in days.

37 Publications, published and planned

1. Protocol: On trial website and published.²⁴
2. Statistical Analysis Plan (SAP) and Health economics analysis plan: On trial website and this publication.
3. Baseline characteristics.
4. Primary results using data to day 90.
5. Health economics using data to day 90.
6. Prediction of swallowing recovery including responder analysis.
7. Pharyngeal electrical stimulation: experience, safety and device deficiencies.
8. Update to published Cochrane Collaboration systematic review.
9. Update to published individual patient data meta-analysis.¹⁴
10. Design learnings, including training, weekends, consent, challenges, free water, eating and drinking with acknowledged risks (EDAR), dysphasia, co-enrolment.
11. Validation of PRESS score.⁶¹
12. Primary swallowing/dysphagia results using data to day 360.
13. Health economics using data to day 360.
14. Cognition and mood trajectory using data to day 360.
15. Prevalence and use of oral trials.
16. Qualitative assessment of PES delivery.
17. Calculating swallowing and eating scales from clinical information.

38 Data sharing

Data from PhEAST will be added to the Cochrane Collaboration review of PSD⁴ and integrated into an updated individual patient data meta-analysis of PES.⁶² Individual patient data will be shared with the 'Virtual International Stroke Trials Archive' (VISTA)⁶³ and, ultimately, made available over the web.³⁴

SUPPLEMENT 2

Health Economic Analysis Plan (HEAP) for the 'Pharyngeal electrical stimulation for acute stroke dysphagia trial' (PhEAST)

Version 5 26.01.2026

Authors: Marilyn James, Cristina Roadevin. Nottingham Clinical Trials Unit

Signatures

Role	Name	Signature	Date
<i>Author, Professor of Health Economics</i>	Prof Marilyn James		
<i>Author, Health Economist</i>	Dr Cristina Roadevin		

1. Purpose of HEAP

The aim of this HEAP is to outline the design and analytical methods used to evaluate the cost effectiveness of the PhEAST trial intervention compared to usual care. The methods specified are prospective and an indicative guide for the analysis but are subject to change in line with recommendations of official bodies and/or guidelines of best practice. Similarly, completion rates of health economic measures within the trial may require further assumptions or analyses.

2. Economic Perspective

The primary analysis will adopt an NHS and personal social services perspective in accordance with NICE guidance (2022). Parallel analysis will take a broader societal perspective.

3. Economic Data and Management

3.1 Software

Data will be imported from Microsoft Excel and analysed in StataSE (Release 18.5; StataCorp USA).

3.2 Data Cleaning

Plausibility checks will be performed on data fields relevant to the economic evaluation. Any identified issues will be clarified with the data manager or relevant staff.

3.3 Outcomes

Primary outcome (Health economics)

- Health-related quality of life (HRQoL) for participant: EuroQoL-5D-5L (EQ-5D-5L) administered at baseline and days: 14 and 90. We hope to later extend this to days 180 and 365.

3.4 Derivation of Indices of HRQoL and Quality Adjusted Life Years

HRQoL indices will be derived using a relevant population tariff. If unavailable for EQ-5D-5L, responses will be mapped to EQ-5D-3L for analysis. Area under the curve

(AUC) will be used to adjust participant HRQoL for the time spent in their respective health states, constructing quality adjusted life-years (QALYs).

3.5 Health Care and Other Resource Use

3.5.1 Patient level resource use

Patient level resource use data will be collected using a purposely designed patient resource proforma. This will collect health care and out of pocket and patient level data such as medication, hospital visits, rehabilitation, primary care, aids and appliances, social care and time lost from paid employment for the patient or their relatives/carers. The proforma will be used to collect data at day 90. The same is used at day 180 and 365 and these will be used potentially in a secondary analysis.

3.5.2 Potential intervention-specific costs

1. PES Equipment Costs:

- Base Station: The durable component of the PES system, which generates, optimises and monitors the delivery of electrical stimulation.
- Catheter: A single-use sterile disposable component. It incorporates electrodes and wiring for delivering electrical stimulation and also functions as a feeding tube. Each patient may require additional catheters to account for potential dislodgement and the need for replacements.

2. Staff Costs:

- Training: Costs associated with training Nurses and Speech and Language Therapists to administer PES.
- Staff type and grade delivering PES: Costs for the time spent by trained staff to administer PES treatments. This includes six daily 10-minute sessions per patient.
- Monitoring and Support: Costs for ongoing monitoring and support, including troubleshooting and ensuring the correct administration of PES.

3. Swallowing Therapy:

Costs related to swallowing therapy contact time (SLT) which might be impacted by the effectiveness of PES.

4. PEG/RIG tube use

The cost of the PEG/RIG tube itself, including any ancillary supplies required for its placement and maintenance. The cost of tube placement, typically involving an endoscopic (or radiologic) procedure performed by a gastroenterologist. Ongoing costs for maintaining the tube, including supplies for cleaning and feeding, and any additional procedures required to manage complications (e.g., bleeding, infections, displacement). Nursing, dietician and other healthcare professional time required to care for patients with tubes, including feeding and managing the tube.

5. Diet costs

Costs related to specialised diets, thickening powders and nutritional supplements provided to patients with dysphagia. Includes the cost of preparing and delivering modified food and thickened liquid consistencies to ensure safe swallowing. For participants who are receiving food via a tube, includes costs of enteral feeding sets

3.6 Costing of Resource Use

Costs will be adjusted to current values using inflation indices from PSSRU.⁶⁴

Table 1. Costing sources

NHS inpatient admissions	The National Cost Collection for the NHS (https://www.england.nhs.uk/costing-in-the-nhs/national-cost-collection/)
NHS outpatient visits, primary/community services, residential care, nursing care	Unit Costs of Health and Social Care (https://www.pssru.ac.uk/). ⁶⁴
Medication	British National Formulary

	(https://bnf.nice.org.uk/)
Special equipment	https://www.essentialaids.com/mobility/walking-frames-accessories.html
Informal care	The costs of absenteeism will be estimated through the lost wages approach. Reports of health-related time taken off work to help with patient's care will be combined with population-level gross weekly salaries from the Annual Survey of Hours and Earnings (Office for National Statistics, 2024).

Table 2. Costing sources, direct intervention costs

PES Equipment Costs	Company (Phagenyx®, Phagenesis Ltd)
Staff costs(training, administration, monitoring)	Company (Phagenyx®, Phagenesis Ltd)
Speech and Language therapy	Unit Costs of Health and Social Care ⁶⁴ (https://www.pssru.ac.uk). The National Cost Collection for the NHS Duration from sites
PEG/RIG tube	The National Cost Collection for the NHS
Diet costs	Sites/Pharma

We will apply the following assumption when costing the 'other' category, where patients have provided free-text responses: the average cost of items within that category will be imputed. For example, if a drug is listed under 'other', we will assign it the average cost of all medications.

4. Procedures for Missing Data

Handling of missing data will follow guidelines for intention-to-treat analysis with incomplete observations.⁶⁵ Multiple imputation methods will be applied using available case data, such as Multiple Imputation of Chained Equations (MICE), which build into their models the inherent uncertainty associated with the missing data.⁶⁶

5. Within-trial Analysis

5.1 Population and Time Horizon

The economic evaluation will use an incremental approach between the two groups with an intention-to-treat (ITT) population and a 90-day time horizon. We hope to extend this to a 365-day horizon subject to funding.

5.2 Discount Rates

No discounting will be applied to derived QALYs or costs within the 3 month period.

5.3 Analysis of Outcomes and Resource Use

The economic evaluation will adopt both a healthcare perspective (NHS and Personal Social Services) and a broader societal perspective. Costs, QALYs, and cost-effectiveness estimates will be derived using multiply imputed datasets to address missing data and improve analytical robustness.

Cost-effectiveness will be assessed through incremental cost-effectiveness ratios (ICERs) and incremental net monetary benefits (INMBs). In the base-case analysis, differences in costs and outcomes between trial arms will be estimated using seemingly unrelated regression (SUR) models applied to the imputed datasets. This method enables the joint estimation of cost and QALY equations, allowing for

correlation between their error terms while adjusting for baseline characteristics and other relevant covariates.

Regression models will estimate treatment-specific mean costs and QALYs, adjusting for age, sex, and other potentially important trial variables. To account for baseline differences in health-related quality of life, QALY models will also include baseline EQ-5D preference scores as a covariate.

In addition to the cost-utility analysis, we will perform a cost effectiveness analysis focusing on the trial's primary outcome, the Dysphagia Severity Rating Scale (DSRS), provided that the outcome favours the intervention and is statistically significant. Using the clinical outcome analysis provided by the statistics team, we will calculate incremental cost effectiveness ratios (ICERs) by dividing the mean difference in costs by the mean difference in DSRS scores between the PES and no-PES group. If the clinical outcome favours standard care, a cost effectiveness analysis will not be undertaken, as the intervention would be dominated by standard care and the calculation of an ICER would not be informative for decision making.

5.4 Sensitivity Analyses

Uncertainty surrounding the estimated treatment effects will be addressed through probabilistic sensitivity analysis (PSA). This will involve simulating the joint distribution of regression coefficients using 10,000 Monte Carlo iterations. This approach will enable us to quantify how uncertainty in both costs and outcomes translates into the likelihood of overall cost effectiveness.

Cost Effectiveness Acceptability Curves (CEACs) will be generated using the net monetary benefit framework, which expresses the value to the NHS of the intervention for a given willingness to pay threshold (λ) for the outcome of interest. The incremental net monetary benefit is defined as:

$$\text{INMB} = (\lambda \times \Delta\text{Outcome}) - (\text{Cost}_{\text{Intervention}} - \text{Cost}_{\text{TAU}})$$

Under this framework, the decision rule is that an intervention is considered cost effective when the incremental net monetary benefit is greater than zero, which is equivalent to the incremental cost effectiveness ratio being below the willingness to pay threshold. Using joint uncertainty in costs and outcomes derived from the SUR based probabilistic sensitivity analysis, we will calculate the INMB across a range of willingness to pay thresholds for each simulation. The CEAC will then show the probability that the intervention is cost effective at different willingness to pay thresholds per QALY. In addition, sensitivity analysis will be used to explore key cost drivers if necessary.

6. Reporting/Publishing

6.1 Reporting Standards

The final report will be submitted for peer review alongside a CHEERS checklist.

6.2 Reporting Deviations from the HEAP

Any deviations from the HEAP will be reported in full and communicated with the study team.

7. References

National Institute for Health and Care Excellence. NICE health technology evaluations: the manual 2022. <https://www.nice.org.uk/process/pmg36/chapter/introduction-to-health-technology-evaluation>

Office for National Statistics. Dataset: Consumer price inflation tables. 2024 (<https://www.ons.gov.uk/economy/inflationandpriceindices/datasets/consumerpriceinflation>)

PhEAST SAP

11/08/2026

Version 0.871

Office for National Statistics. Dataset: Earnings and hours worked, all employees:

ASHE Table 1. 2024

(<https://www.ons.gov.uk/employmentandlabourmarket/peopleinwork/earningsandworkinghours/datasets/allemployeesashtable1>)

Example tabulated results at 12 months are shown in Supplement 5.

SUPPLEMENT 3

PHARYNGEAL ELECTRICAL STIMULATION FOR ACUTE/SUB-ACUTE POST-STROKE DYSPHAGIA: MAIN RESULTS FROM THE PHEAST TRIAL

1. DESCRIPTION OF MAIN PAPER

Title

Pharyngeal electrical stimulation for acute post-stroke dysphagia: main results from the PhEAST trial

Authors

PhEAST Investigators

Aim

Describe the primary outcome and key secondary outcomes.

Participants

All in PhEAST.

Outcomes

Primary outcome, days 14/90: DSRS.

Secondary outcomes, days 14/90:

Safety, day 14/90: serious adverse events, serious adverse device effects, unexpected serious adverse device effects, device deficiencies.

Analyses

Descriptive statistics: number (%), median [interquartile range] or mean (standard deviation). Comparisons using repeated measures regression, multiple linear regression, ordinal logistic regression, Cox proportional hazards regression, binary logistic regression and Wei-Lachin test.

Table 1. Characteristics of the patients at baseline.

Characteristic	N	All (N=xxx)	PES Group (N=xxx)	No PES Group (N=xxx)
Demographics				
Age (years) #†	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)
Sex, female (%) †	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Country (%)				
Austria	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Denmark	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
UK	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Ethnicity (%)				
Other than white	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
White	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Onset-randomisation (days) #†	xxx	xx.x [xx.x, xx.x]	xx.x [xx.x, xx.x]	xx.x [xx.x, xx.x]
Swallowing				
DSRS (/12) †‡	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)
Fluids [/4]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)
Diet [/4]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)
Supervision [/4]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)
FOIS [/7] #§	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]
1 (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
2 (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
3 (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
IDDSI				
Food (3-7)		x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]
Drinks (0-4)		x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]
Functional diet scale (0-8)		x.x (x.x)	x.x (x.x)	x.x (x.x)
Feeding status scale (/6) %	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]

Characteristic	N	All (N=xxx)	PES Group (N=xxx)	No PES Group (N=xxx)
0. Normal diet		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
1. Soft diet		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
2. NGT-fed		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
3. PEG/RIG-fed		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
4. IV/sc fluids only		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
5. No fluids/feeding		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
EAT-10 (/50)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)
Clinical				
Modified Rankin scale, premorbid [/5]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]
Modified Rankin scale, baseline [/5]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]
Previous stroke (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
NIHSS (/42) †¶	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)
Barthel index (/100)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)
Stroke type, haemorrhage (%) †	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Stroke syndrome (%)				
Total anterior circulation †	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Partial anterior circulation	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Lacunar	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Posterior circulation	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Weight (kg)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)

DSRS (dysphagia rating scale); EAT-10: eating assessment tool-10; FOIS: functional oral intake score; IDDSI: international dysphagia diet standardisation initiative; NIHSS (NIH stroke severity scale); OCSP: Oxfordshire community stroke project; TACS: total anterior circulation syndrome.

Data are number (%), median [interquartile range] or mean (standard deviation).

Inclusion criterion

† Minimisation variables: age (<75/75+), sex (female/male), DSRS (<12/12), impairment (NIHSS <15/15+), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior), time to randomisation (<15/15+ days), country (other than UK/UK).

‡ The dysphagia severity rating scale (DSRS) evaluates swallowing impairment with scores ranging from 0 (normal) to 12 with higher scores indicating more severe swallowing impairment.^{12,26} The total score is the sum of three subscale scores relating to fluids, food and supervision, each ranging from 0 (normal) to 4 (nil by mouth). See table 1.1 for trial specific information on DSRS scoring for patients on oral trials.

§ The functional oral outcome scale (FOIS) evaluates swallowing impairment with scores ranging from 1 to 7 with lower scores indicating more severe swallowing impairment; inclusion criteria were: 1 – nil by mouth, 2 – tube dependent with minimal attempts of food or liquid, 3 – tube dependent with consistent oral intake of food or liquid.²⁵ See 36a for trial specific information on FOIS scoring for patients on oral trials.

¶ The National Institutes of Health stroke scale (NIHSS) evaluates stroke severity/impairment with scores ranging from 0 to 42 with higher scores indicating greater neurological impairment.⁶⁷

Table 2. Primary and secondary swallowing and feeding outcomes at days 14 and 90 with no imputation. Analyses by adjusted continuous repeated measures regression (acRMR), adjusted ordinal repeated measures regression (aoRMR), adjusted Cox proportional hazards regression (aCPHR). Or adjusted binary logistic regression (aBLR)

Outcome	N	PES	No PES	Treatment effect	Test
1ry outcome					
DSRS [1]				x.xx (x.xx, x.xx), p=0.xxx	acRMR
Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
2ry outcomes					
FOIS [2]				x.xx (x.xx, x.xx)	aoRMR
Baseline	xxx	x.x (x.x)	x.x (x.x)		
Day 14	xxx	x.x (x.x)	x.x (x.x)		
Day 90	xxx	x.x (x.x)	x.x (x.x)		
IDDSI FDS [3]				x.xx (x.xx, x.xx)	acRMR
Baseline	xxx	x.x (x.x)	x.x (x.x)		
Day 14	xxx	x.x (x.x)	x.x (x.x)		
Day 90	xxx	x.x (x.x)	x.x (x.x)		
FSS [4]				x.xx (x.xx, x.xx)	aoRMR
Baseline	xxx	x.x (x.x)	x.x (x.x)		
Day 14	xxx	x.x (x.x)	x.x (x.x)		
Day 90	xxx	x.x (x.x)	x.x (x.x)		
EAT-10 [5]				x.xx (x.xx, x.xx)	acRMR
Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
Safety outcomes (%)					
Death from any cause day 90	xxx	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aCPHR
Serious adverse events (SAE) day 14	xxx	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Serious adverse device event (SADE) day 14	xxx	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Unexpected SADE (USADE) day 14	xxx	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR

DSRS: dysphagia severity rating scale; EAT-10: eating assessment tool; FSS: feeding status scale; IDDSI-FDS: international dysphagia diet standardisation initiative – functional diet scale; PES: pharyngeal electrical stimulation.

Data are number (%), median [interquartile range] or mean (standard deviation). Analyses are intention to treat with no imputation of missing data. Generalised linear mixed models (SAS "PROC MIXED") were used to calculate adjusted repeated measures regression (aRMR). The models include randomisation group with adjustment for baseline DSRS, age, sex, impairment (NIHSS), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior) and time to randomisation (days). Participants were included if they had DSRS measured and days 14 and/or 90; participants who died scored 13 on the DSRS. Professional scores are used where participants were eating and drinking with acknowledged risk (EDAR) or on a free water protocol (FWP).

1. The dysphagia severity rating scale (DSRS) evaluates swallowing impairment with scores ranging from 0 (normal) to 12 with higher scores indicating more severe swallowing impairment^{12,26} and death indicated by 13. The total score is the sum of three subscale scores relating to fluids, food and supervision, each ranging from 0 (normal) to 4 (nil by mouth). The DSRS x time interaction p value was 0.xxx.
2. The functional oral outcome scale (FOIS) evaluates swallowing impairment with scores ranging from 1 to 7 with lower scores indicating more severe swallowing impairment²⁵ and death indicated by 0; inclusion criteria were: 1 – nil by mouth, 2 – tube dependent with minimal attempts of food or liquid, 3 – tube dependent with consistent oral intake of food or liquid.
3. The international dysphagia diet standardisation initiative - functional diet scale (IDDSI-FDS) evaluates the severity of oropharyngeal dysphagia represented by diet texture restriction with scores ranging from 0 (nil by mouth) to 8 (normal food texture and drink consistency⁵⁰ and death indicated by -1.
4. The feeding status scale (FSS) is a measure of nutritional status with scores 0 (normal oral diet), 1 (soft diet), 2 (nasogastric tube feeding), 3 (PEG/RIG/J-tube feeding), 4 (parenteral fluids), 5 (no fluids/feeding) and 6 (died).^{47,48} Non-oral feeding is a score that exceeds 1.
5. EAT-10 is a measure of swallowing difficulties comprising 10 questions with each score ranging for 0 (no problem) to 4 (severe problem);⁵¹ the scale is extended to include a score of 5 for nil by mouth hence the total score ranges from 0 to 50 with death assigned 51.

Figure 1. Swallowing impairment at 14 and 90 days in the two groups according to scores on the dysphagia severity rating scale (primary outcome).

Shift	Score	0	1	2	3	4	5	6	7	8	9	10	11	12	13
	Day														
PES	00	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	0
	14	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx
	90	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx
No PES	90	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx
	14	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx
	00	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	xx	0

Scores on the dysphagia severity rating scale range from 0 to 13 with 0 indicating no swallowing impairment/normal diet, higher scores indicating more severe swallowing impairment, 12 indicating nil by mouth and 13 indicating death.^{12,26} The total score is the sum of three subscales relating to fluids, food and supervision, each ranging from 0 (normal) to 4 (nil by mouth).

Figure 2. Cumulative incidence of death out to 90 days with analysis by Cox proportional hazards regression.

Panel shows inset with an enlarged y axis.

Figure 3. Subgroup analysis of the primary outcome.

Shown are the difference in means for swallowing impairment in pre-specified subgroups including minimisation factors. The widths of the confidence intervals have not been adjusted for multiplicity and should not be used in place of hypothesis testing. DSRS scores range from 0 to 12, with 13 indicating death and higher scores indicating worse swallowing impairment. FOIS scores range from 1 to 7, with 0 indicating death and lower scores indicating worse swallowing impairment. NIHSS scores range from 0 to 42, with 43 indicating death and higher scores indicating greater neurological impairment. Age, FOIS, DSRS, NIHSS and time are dichotomised as shown.

- Age (years): <75, ≥75
- Sex: female, male
- Reperfusion: yes, no
- Dysphagia severity rating scale (/12): <11, 11, 12
- Functional oral intake scale (/7): 1, 2/3
- Stroke severity, NIHSS (/42): <13, ≥13
- Stroke type: ischaemic, primary intracerebral haemorrhage
- Circulation: anterior, posterior
- Onset to randomisation (days): <11, 11-20, >20
- Lesion side: right, bilateral, left
- Stroke size: very small/small, medium, large/very large
- White matter disease: none, mild, moderate, severe
- Prior instrumental testing (FEES/VFS): yes, no
- Responder, DSRS change day 90-0: >-3.5, ≤-3.5
- Responder, DSRS day 90: <4, ≥4
- Experience with PES prior to trial: yes, no
- Site recruitment: >10, ≤10

FOREST PLOT WITH INTERACTION TERMS

SUPPLEMENT FOR MAIN PAPER

Supplement to main paper

Table S1. Reasons for excluding from participating in the trial.

Reason †	N (%)
FOIS >3	XXXX (xx.x)
Premorbid dysphagia	XXXX (xx.x)
Palliative/end-of-life care	XXXX (xx.x)
Repatriated	XXXX (xx.x)
Frequent removal of nasogastric tubes	XXXX (xx.x)
Declined consent	XXXX (xx.x)
Pacemaker/cochlear implant/ICD	XXXX (xx.x)
Out of time window	XXXX (xx.x)
No clinical indication for NGT or PEG	XXXX (xx.x)
Non-stroke dysphagia	XXXX (xx.x)
Pre-stroke mRS of 4 or 5	XXXX (xx.x)
Ongoing/planned ventilation, intubation or tracheostomy	XXXX (xx.x)
Planned electrical/magnetic stimulation for dysphagia	XXXX (xx.x)
Clinical indication for >35% oxygen	XXXX (xx.x)
Patient unlikely to tolerate PES catheter/treatment	XXXX (xx.x)
Active malignant MCA syndrome	XXXX (xx.x)
Recruited into another clinical trial	XXXX (xx.x)
No staff available to recruit or complete procedures	XXXX (xx.x)
Patient or next of kin declined	XXXX (xx.x)
Patient unfit for research due to other comorbidities	XXXX (xx.x)
Language barrier	XXXX (xx.x)
Eating & drinking with acknowledged risk (EDAR)	XXXX (xx.x)
Not appropriate for swallow assessment(s)	XXXX (xx.x)
Pregnant	XXXX (xx.x)
Presence of pharyngeal pouch	XXXX (xx.x)
NIHSS score 1a = >2	XXXX (xx.x)
Patient from overseas so unable to follow-up	XXXX (xx.x)
Dysphagia considered to be short-term only	XXXX (xx.x)
Total	XXXX

MCA: middle cerebral artery

† Items are not mutually exclusive, i.e. patients may have been excluded for more than one reason.

TO BE ORDERED BY FREQUENCY

Table S2. Clinical examination of swallowing at baseline and prognostication for recovery .

Characteristic	N	All	PES Group (N=xxx)	No PES Group (N=xxx)
Dysphonia (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Dysarthria (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Abnormal gag reflex (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Abnormal volitional cough (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Cough after swallow (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Voice change after swallow (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Frontal operculum involved †	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Calculated PRESS score ‡	xxx	xx.x (xx.x)	xxx (xx.x)	xxx (xx.x)

† Brain scans before or at baseline were not collected and adjudicated; admission neuroradiology reports (see Table S3 below) were entered by recruiting sites; some of these reported the presence of a stroke lesion (infarct or haemorrhage) at the frontal operculum although this item may have been under-reported and so the total PRESS score will be underestimated by one point for some participants.

‡ The predictive swallowing score (PRESS⁶¹) is a prognostic instrument that predicts swallowing recovery by 30 days after stroke and ranges from 0 (no features present) to 10 (10 features present) based on age ≥ 70 years; NIHSS score; stroke location includes lesion at the frontal operculum; presence of dysphonia, dysarthria, abnormal gag reflex, abnormal volitional cough, cough after swallow and voice change after swallow; and FOIS score. Scoring is given in reference.⁶¹ Brain scans before or at baseline were not collected and adjudicated. Instead, relevant neuroradiology reports were entered by recruiting sites but few made any specific comment about the presence of a stroke lesion at the frontal operculum. Hence, this item may have been under-reported and so the total PRESS score is likely an underestimate.

Table S3. Neuroimaging features as recorded by site's (neuro)-radiologist imaging report between admission to hospital and randomisation.

Feature	N (%)	All (N=xxx)	PES (n1=xxx)	No PES (n2=xxx)
CT scan	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
MR scan	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Reported by site (%)	xxx (xx.x)			
Normal scan		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Infarct		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Infarct with HTI		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
ICH		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
SAH		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Other lesion		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Ischaemic stroke	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Side	xxx (xx.x)			
Right		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Left		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Bilateral/midline		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Size	xxx (xx.x)			
Very large		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Large		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Moderate		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Small		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Very small		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Hyperdense artery	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Site	xxx (xx.x)			
Frontal		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Parietal		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Temporal		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Subcortical		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Basal ganglia		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Internal capsule		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Pontine		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Brain stem		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Cerebellum		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Local sites	xxx (xx.x)			
Insula		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Frontal operculum		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Small vessel disease	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Lacunar infarction		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Lacune(s)		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
White matter change		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
SVD score [/3]	xxx (xx.x)	x.x (x.x)	x.x (x.x)	x.x (x.x)
Brain frailty score [/3]	xxx (xx.x)	x.x (x.x)	x.x (x.x)	x.x (x.x)
ICH	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Side	xxx (xx.x)			
Right		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Left		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Bilateral/midline		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Size	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Very large		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Large		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)

PhEAST SAP

11/08/2026

Version 0.871

Moderate		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Small		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Very small		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Site	xxx (xx.x)			
Basal ganglia		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Brainstem		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Cerebellum		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Centrum semiovale		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Corona radiata		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Frontal		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Midbrain		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Occipital		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Parietal		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Temporal		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Other sites				
Frontal operculum		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Background	xxx (xx.x)			
Atrophy		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Old infarct		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Other diagnoses				
Intraventricular haemorrhage	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Subarachnoid haemorrhage	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Subdural	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Arteriovenous malformation	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Aneurysm	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)

† Participants may have multiple features. Data are n (%)

Radiology reports were not available for all participants so numbers (%) may differ from those reported by investigators.

Imaging report interpreted by algorithmic natural language processing: accuracy depends on the structure, detail, completeness and quality of the radiologist report and robustness of the NLP program to non-standard English and missing data. Hence the numbers should be considered approximate and incomplete.

Table S4. Hospital care before and after randomisation.

Procedure	N	All	PES	No PES	aDIM/acOR/aOR (95% CI) [†]	Analysis
Pre-randomisation (%)		xxx	xxx	xxx	/	/
Intravenous thrombolysis	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Mechanical thrombectomy	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Hemicraniectomy	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Neurosurgery, other	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Vascular surgery	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Intensive care unit admission	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Ventilation	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Days of ventilation	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Tracheostomy	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Days with a tracheostomy	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Post-randomisation (%)		xxx	xxx	xxx		
Neurosurgery	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Vascular surgery	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Intensive care unit admission	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Ventilation	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Tracheostomy	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
At discharge/death in hospital (%)		xxx	xxx	xxx		
Antithrombotic	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Antiplatelet	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Anticoagulant, oral	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Statin	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Antihypertensive	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Gastroprotection	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR

aDIM: adjusted difference in means; acOR: adjusted common odds ratio; aOR: adjusted odds ratio; CI: confidence intervals; PES: pharyngeal electrical stimulation

Table S5. Adherence to trial allocation

Assessment	PES
Participants	xxx
<i>PES doses received (%)</i>	
First dose received	xxx (xx.x)
Doses received, at least	
1	xxx (xx.x)
2	xxx (xx.x)
3	xxx (xx.x)
4	xxx (xx.x)
5	xxx (xx.x)
6, all	xxx (xx.x)
Number of doses, median [IQR]	x.x [x.x, x.x]
<i>Reasons for non-adherence to first three treatments (%)</i>	
Withdrawal from treatment	xx (xx)
Withdrawal from trial	xx (xx)
Clinical decision by hospital staff	xx (xx)
Participant discharged	xx (xx)
Lost to follow-up	xx (xx)
PES failure	xx (xx)
Died	xx (xx)
Other	xx (xx)

Table S6. Treatment with pharyngeal electrical stimulation: threshold, tolerance, calculated stimulation, actual stimulation and delta (tolerability-threshold), all in milliamps (mA), and treatment duration in participants randomised to PES.

Data are number (%), median [interquartile range] or mean (standard deviation).

Treatment	N	Threshold	Tolerance	Calculated stimulation	Stimulation	Delta	Duration (secs)
1	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xxx.x [xxx.x, xxx.x]
2	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xxx.x [xxx.x, xxx.x]
3	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xxx.x [xxx.x, xxx.x]
4	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xxx.x [xxx.x, xxx.x]
5	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xxx.x [xxx.x, xxx.x]
6	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xxx.x [xxx.x, xxx.x]
Average	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xxx.x [xxx.x, xxx.x]

Table S7. Treatment with pharyngeal electrical stimulation: participants receiving adequate stimulation

	N	Stimulation ≥ 20 mA	Delta > 20 mA [†]	Time ≥ 9 m50s	Catheters used (SD)
Participants (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.x (x.x)
Treatments (%)	xxxx	xxxx (xx.x)	xxxx (xx.x)	xxxx (xx.x)	/

[†] Delta = tolerance - threshold

Table S8. Swallowing and feeding outcomes at days 14 and 90. Sensitivity analyses by continuous repeated measures regression (cRMR) or ordinal repeated measures regression (oRMR) with adjustment.

Population	Imputation	Adjusted	Outcome	N	PES	No PES	Average difference	
<i>Primary, DSRS [1]</i>								
ITT	Yes	Yes	DSRS				x.xx (x.xx, x.xx)	acRMR
			Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
mITT	No	No	DSRS				x.xx (x.xx, x.xx)	cRMR
			Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
PP	No	Yes	DSRS				x.xx (x.xx, x.xx)	acRMR
			Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
mITT	No	Yes	DSRS [A]				x.xx (x.xx, x.xx)	acRMR
			Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
mITT	No	Yes	DSRS fluids				x.xx (x.xx, x.xx)	aoRMR
			Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
mITT	No	Yes	DSRS diet				x.xx (x.xx, x.xx)	aoRMR
			Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
mITT	No	Yes	DSRS supervision				x.xx (x.xx, x.xx)	aoRMR
			Baseline	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 14	xxx	xx.x (xx.x)	xx.x (xx.x)		
			Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)		
<i>Secondary</i>								

PhEAST SAP			11/08/2026	Version 0.871				
ITT	Yes	Yes	FOIS [2] Baseline Day 14 Day 90	xxx xxx xxx	x.x (x.x) x.x (x.x) x.x (x.x)	x.x (x.x) x.x (x.x) x.x (x.x)	x.xx (x.xx, x.xx)	aoRMR
ITT	Yes	Yes	IDDSI food [3] Baseline Day 14 Day 90	xxx xxx xxx	x.x (x.x) x.x (x.x) x.x (x.x)	x.x (x.x) x.x (x.x) x.x (x.x)	x.xx (x.xx, x.xx)	aoRMR
ITT	Yes	Yes	IDDSI drinks [3] Baseline Day 14 Day 90	xxx xxx xxx	x.x (x.x) x.x (x.x) x.x (x.x)	x.x (x.x) x.x (x.x) x.x (x.x)	x.xx (x.xx, x.xx)	aoRMR
ITT	Yes	Yes	IDDSI FDS [4] Baseline Day 14 Day 90	xxx xxx xxx	x.x (x.x) x.x (x.x) x.x (x.x)	x.x (x.x) x.x (x.x) x.x (x.x)	x.xx (x.xx, x.xx)	acRMR
ITT	Yes	Yes	FSS [5] Baseline Day 14 Day 90	xxx xxx xxx	x.x (x.x) x.x (x.x) x.x (x.x)	x.x (x.x) x.x (x.x) x.x (x.x)	x.xx (x.xx, x.xx)	acRMR
ITT	Yes	Yes	EAT-10 [6] Baseline Day 14 Day 90	xxx xxx xxx	xx.x (xx.x) xx.x (xx.x) xx.x (xx.x)	xx.x (xx.x) xx.x (xx.x) xx.x (xx.x)	x.xx (x.xx, x.xx)	acRMR

DSRS: dysphagia severity rating scale; EAT-10: eating assessment tool; FSS: feeding status scale; IDDSI-FDS: international dysphagia diet standardisation initiative – functional diet scale; ITT: intention to treat; mITT: modified ITT; PES: pharyngeal electrical stimulation; PP: per protocol.

Data are number (%), median [interquartile range] or mean (standard deviation). Generalised linear mixed models (SAS proc mixed) were used to calculate adjusted repeated measures regression (aRMR). The models include randomisation group with adjustment for baseline score, age, sex, DSRS, impairment (NIHSS), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior) and time to randomisation (days).

A. Assesses participants who are EDAR or using a free water protocol.

1. The dysphagia severity rating scale (DSRS) evaluates swallowing impairment, with scores ranging from 0 (normal) to 12 (nil by mouth) ^{12,26} and death indicated by 13.
2. The functional oral outcome scale (FOIS) evaluates swallowing impairment with scores ranging from 1 to 7 with lower scores indicating more severe swallowing impairment ²⁵ and death indicated by 0.
3. The international dysphagia diet standardisation initiative (IDDSI) is a global standardised framework describing food texture and liquid thickening;⁴⁹ food texture scores range between 7 (regular) and 3 (liquidised) and death indicated by 5; fluid thickening scores range between 0 (thin) and 4 (extremely thick) and death indicated by 5.
4. The international dysphagia diet standardisation initiative - functional diet scale (IDDSI-FDS) evaluates the severity of oropharyngeal dysphagia represented by diet texture restriction with scores ranging from 0 (nil by mouth) to 8 (normal food texture and drink consistency ⁵⁰ with death assigned -1.
5. The feeding status scale (FSS) is a measure of nutritional status with scores 0 (normal oral diet), 1 (soft diet), 2 (nasogastric tube feeding), 3 (PEG/RIG/J-tube feeding), 4 (parenteral fluids), 5 (no fluids/feeding) and 6 (died).^{47,48} Non-oral feeding is a score that exceeds 1.
6. The eating assessment tool-10 (EAT-10) is a measure of swallowing difficulties comprising 10 questions with each score ranging for 0 (no problem) to 4 (severe problem);⁵¹ the scale is extended to include a score of 5 for nil by mouth hence the total score ranges from 0 to 50 with death assigned 51.

Table S9. Secondary outcomes related to dysphagia, swallowing and feeding at day 14, discharge and day 90.

Analyses used adjusted binary logistic regression (aBLR), adjusted ordinal logistic regression (aOLR) or adjusted multiple linear regression (aMLR).

Outcome	N	All (N=xxx)	PES (N=xxx)	No PES (N=xxx)	aDIM/acOR/aOR (95% CI) [†]	Analysis
Primary outcome, sensitivity						
DSRS, day 14 [1]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
DSRS, day 14 change from baseline	xxx	x.x (x.x)	x.x (x.x)	x.x (x.x)	x.x (x.x, x.x)	aMLR
DSRS, day 90 [1]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
DSRS, day 90 change from baseline	xxx	x.x (x.x)	x.x (x.x)	x.x (x.x)	x.x (x.x, x.x)	aMLR
Swallowing/feeding						
FOIS, day 14 [2]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aOLR
FOIS, day 90 [2]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aOLR
IDDSI FDS, day 14 [3]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
IDDSI FDS, day 90 [3]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Feeding status scale, day 14 [4]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.xx (x.xx, x.xx)	aOLR
Normal diet		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Soft diet		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
NGT-fed		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
PEG/RIG fed		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
IV/sc fluids only		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
No fluids/feeding		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Died		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Feeding status scale, day 90 [4]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.xx (x.xx, x.xx)	aOLR
Normal diet		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Soft diet		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
NGT-fed		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
PEG/RIG/J-tube fed		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
IV/SC fluids only		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
No fluids/feeding		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Died		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
Non-oral diet, FSS>1, day 14 (%)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)	x.xx (x.xx-x.xx)	aBLR
Non-oral diet, FSS>1, day 90 (%)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)	x.xx (x.xx-x.xx)	aBLR

PhEAST SAP	11/08/2026	Version 0.871				
EAT-10, day 14 [5]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
EAT-10, day 90 [5]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Global swallowing, day 14 [6]	xxx	/	/	/	x.xx (x.xx, x.xx)	WLT
Global swallowing, day 90 [6]	xxx	/	/	/	x.xx (x.xx, x.xx)	WLT
PEG/RIG/J-tube						
At day 14	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
At discharge	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Instrumental assessment						
Before baseline	xxx					
FEES		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
VFS		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
FEES or VFS		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
PAS score [7]		x.xx (x.xx)	x.xx (x.xx)	x.xx (x.xx)	/	
Before day 14	xxx					
FEES		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
VFS		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
FEES or VFS		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
PAS score [7]		x.xx (x.xx)	x.xx (x.xx)	x.xx (x.xx)	/	
Before death or discharge	xxx					
FEES		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
VFS		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
FEES or VFS		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
PAS score [7]		x.xx (x.xx)	x.xx (x.xx)	x.xx (x.xx)	/	

DD: at death or discharge; DSRS: dysphagia severity rating scale; FEES: fiberoptic endoscopic evaluation of swallowing; IDDSI: International dysphagia diet standardisation initiative; IV: intravenous; NIHSS: National Institute of Health stroke scale; PES: pharyngeal electrical stimulation; SADE: serious adverse device event; SAE: serious adverse event; SC: subcutaneous; USADE: unexpected serious adverse device event; VFS: videofluoroscopy.

Data are number (%), median [interquartile range] or mean (standard deviation). The models include randomisation group with adjustment for baseline score, age, sex, DSRS, impairment (NIHSS), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior) and time to randomisation (days).

1. The dysphagia severity rating scale (DSRS) evaluates swallowing impairment, with scores ranging from 0 (normal) to 12 (nil by mouth)^{12,26} with death assigned 13.

2. The functional oral outcome scale (FOIS) evaluates swallowing impairment with scores ranging from 1 to 7 with lower scores indicating more severe swallowing impairment ²⁵ with death assigned 0.
3. The international dysphagia diet standardisation initiative - functional diet scale (IDDSI-FDS) evaluates the severity of oropharyngeal dysphagia represented by diet texture restriction with scores ranging from 0 (nil by mouth) to 8 (normal food texture and drink consistency ⁵⁰ with death assigned -1.
4. The feeding status scale (FSS) is a measure of nutritional status with scores 0 (normal oral diet), 1 (soft diet), 2 (nasogastric tube feeding), 3 (PEG/RIG/J-tube tube feeding), 4 (parenteral fluids), 5 (no fluids/feeding) and 6 (died).⁴⁷*Ref Non-oral feeding is a score that exceeds 1.
5. The eating assessment tool-10 (EAT-10) is a measure of swallowing difficulties comprising 10 questions with each score ranging for 0 (no problem) to 4 (severe problem);⁵¹ the scale is extended to include a score of 5 for nil by mouth hence the total score ranges from 0 to 50 with death assigned 51.
6. A global analysis of DSRS, FOIS, IDDSI-FDS, FSS and EAT-10 was calculated using the Wei-Lachin test.^{68,69}
7. The penetration aspiration scale (PAS) evaluates aspiration and penetration using either FEES or VFS. Assessment of PAS was not mandated in the protocol and it is likely that instrumental testing was biased by indication, i.e. it was largely performed in participants whose swallowing was not improving.

Table S10. Other secondary outcomes at day 14, discharge or hospital death and day 90. Analyses used adjusted binary logistic regression, adjusted Cox proportional hazards regression, adjusted ordinal logistic regression, adjusted multiple linear regression or Wei-Lachin test.

Outcome	N	All (N=xxx)	PES (N=xxx)	No PES (N=xxx)	aDIM/acOR/aOR/aHR (95% CI) [†]	Analysis
Day 14						
Died (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
NIHSS (0-42) [1]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Barthel index (/100) [2]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Modified Rankin Scale [3]	xxx	x.x (x.x)	x.x (x.x)	x.x (x.x)	x.x (x.x, x.x)	aOLR
EQ-5D-5L health utility [4]	xxx	0.xx (0.xx)	0.xx (0.xx)	0.xx (0.xx)	x.xx (x.xx, x.xx)	aMLR
EQ-VAS (/100) [5]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Pneumonia (%)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)	x.xx (x.xx-x.xx)	aBLR
Global stroke impact scale [6]	xxx	/	/	/	xx.x (xx.x, xx.x)	WLT
Discharge/death in hospital						
Death/discharge (%) [7]	xxx				x.xx (x.xx, x.xx)	OLR
Died in hospital		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aCPHR
Discharge to institution		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
Discharged home		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	/	
Chest infection or pneumonia (%)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)	x.xx (x.xx-x.xx)	aBLR
Chest infection, pneumonia (%) [8]	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)	x.xx (x.xx-x.xx)	aOLR
None	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		
Chest infection, non-fatal	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		
Pneumonia, non-fatal	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		

PhEAST SAP	11/08/2026		Version 0.871			
Fatal	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		
Any antibiotic use (%)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)	x.xx (x.xx-x.xx)	aBLR
Discharged with a PEG/RIG tube (%)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)	x.xx (x.xx-x.xx)	aBLR
Day 90						
Died (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
PEG/RIG/J-tube <i>in situ</i> (from FSS)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
Home time (days) [9]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Barthel index [2]	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Modified Rankin Scale [3]	xxx	x.x (x.x)	x.x (x.x)	x.x (x.x)	x.x (x.x, x.x)	aMLR
EQ-5D-5L health utility [4]	xxx	0.xx (0.xx)	0.xx (0.xx)	0.xx (0.xx)	x.xx (x.xx, x.xx)	aMLR
EQ-VAS (/100) [5]	xxx	xxx.x (xx.x)	xxx.x (xx.x)	xxx.x (xx.x)	xxx.x (xx.x, xx.x)	aMLR
Stroke impact scale [6]	xxx	/	/	/	xx.x (xx.x, xx.x)	WLT
Disposition [7]	xxx				x.xx (x.xx, x.xx)	aOLR
Died		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx, x.xx)	aBLR
In an institution		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		
At home		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		

PES: pharyngeal electrical stimulation.

Data are number (%), median [interquartile range] or mean (standard deviation). The models include randomisation group with adjustment for baseline score, age, sex, DSRS, impairment (NIHSS), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior) and time to randomisation (days).

§ The proportion hazard assumption P-values are: feeding status scale day 14 0.xx, day 90 0.xx; penetration aspiration scale score 0.xx; modified Rankin scale day 14 0.xx, day 90 0.xx.

1. The National Institutes of Health stroke scale (NIHSS) evaluates stroke severity/impairment with scores ranging from 0 to 42 with higher scores indicating greater neurological impairment.⁶⁷ Death is assigned a score -43.
2. The Barthel index (BI) evaluates disability/activities of daily living with scores ranging from 100 (no disability) to 0 (severe disability) with death assigned -5.
3. The modified Rankin scale evaluates dependency with scores ranging from 0 (normal) to 6 (death).
4. The EuroQol 5-dimension 5-level (EQ-5D-5L) patient-reported questionnaire covers 5 domains of health-related quality of life: mobility, self-care, usual activities, pain and discomfort, and anxiety and depression. Each domain has 3 graded levels of response: no problems, moderate problems, or extreme problems. Scores from these levels are combined to provide an overall health utility score that was calculated according to UK population norms. A score of 1 represents perfect health, a score of 0 represents death, and negative scores represent health states considered to be worse than death.
5. The EuroQol-visual analogue scale (EQ-VAS) patient reported current overall health with scores from 100 (perfect health) to 0 (worst health) with death assigned -1.
6. The Stroke impact scale (SIS) is a patient-reported outcome that measures eight domains of health-related quality of life and functional disability.⁵² A global analysis of the eight domains⁷⁰ was performed using the Wei-Lachin test.^{68,69}
7. Discharge to includes another institution (hospital, residential home, nursing home) or home (own or family home with or without carer, warden-aided flat).
8. Chest infection or pneumonia, as determined clinically by hospital investigators.
9. Home time⁵⁴ is time (days) following discharge up to day 90. Participants who died in hospital or who were discharged to and remain in a nursing home score 0.

Table S11. Safety end points: causality, events (up to day 14) and fatal (up to day 90). Data are number (frequency), central tendency (95% confidence intervals) and probability.

End point	N	All	PES Group (N=xxx)	No PES Group (N=xxx)	aDIM/acOR/aOR (95% CI) [†]	Test
Events to day 14		xxx	xxx	xxx		
Relationship to treatment, all					x.xx (x.xx-x.xx)	aOLR
Not related (SAE)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		
Unlikely to be related (SAE)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		
Possibly related (SADE/USADE)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		
Probably related (SADE/USADE)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		
Definitely related (SADE/USADE)	xxx	xxx (x.x)	xxx (x.x)	xxx (x.x)		
Participants		xxx	xxx	xxx		
Event/effect by day 14						
SAE	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx-x.xx)	BLR
Non-fatal	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx-x.xx)	BLR
Fatal	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx-x.xx)	BLR
SADE	xxx	xxx (xx.x)	xxx (xx.x)	/	/	/
Non-fatal	xxx	xxx (xx.x)	xxx (xx.x)	/	/	/
Fatal	xxx	xxx (xx.x)	xxx (xx.x)	/	/	/
USADE	xxx	xxx (xx.x)	xxx (xx.x)	/	/	/
Non-fatal	xxx	xxx (xx.x)	xxx (xx.x)	/	/	/
Fatal	xxx	xxx (xx.x)	xxx (xx.x)	/	/	/
Fatal SAE by day 90	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx-x.xx)	CPHR

SAE: serious adverse event; SADE: serious adverse device effect; USADE: unexpected serious adverse device effect

✕ The proportion hazard assumption P-value is 0.xx.

Table S12. Fatal and non-fatal serious adverse events, serious adverse device events and unexpected serious adverse device events by day 14.

Event	Median time to event (days)	All	PES	No PES	Fatal	PES	No PES
Participants		xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Cardiac	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Atrial fibrillation	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Pulmonary embolism	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Nervous system	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Complication of stroke	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Cerebral oedema	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Deterioration	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Recurrent stroke	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Seizure/convulsion	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Confusion	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Respiratory	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Chest infection	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Pneumonia	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Pleural effusion	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Pneumothorax	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Gastrointestinal	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
GI bleed	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Vomiting	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Genitourinary	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Acute kidney injury	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Haematuria	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Urinary tract infection	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Haematological/immunological	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Metabolic/endocrine	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Electrolyte disturbance	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Musculoskeletal/cutaneous	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Gout	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Phlebitis	x.x [x.x, x.x]	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)

PhEAST SAP

11/08/2026

Version 0.871

Miscellaneous/other	x.x [x.x, x.x]	xxx (xx.x) xxx (xx.x)xxx (xx.x)	xxx (xx.x)xxx (xx.x)xxx (xx.x)
...			

NOTE: OTHERS WILL BE ADDED AS APPROPRIATE

Table S13. Protocol violations

Violation	All	PES	No PES
Participants	N	n1	n2
No consent	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Ineligible	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Non stroke dysphagia	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Pre-stroke dysphagia	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
NIHSS item 1a score 2 or 3	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Ongoing/anticipated ventilation/intubation	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Use of non-trial device for dysphagia	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Malignant MCA syndrome	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Pacemaker, cochlear implant, defibrillator	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
On >35% oxygen	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
To be repatriated to a separate organisation	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
To be rehabilitated at a separate organisation	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Not expected to be at enrolling hospital for 14 days	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Not expected to tolerate PES catheter	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
To be transferred to a non-participating site	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Known to be pregnant at enrolment	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
In another trial with no co-enrolment agreement	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
On palliative care	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Known pharyngeal pouch	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Dysphagia expected to resolve rapidly	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
EDAR at enrolment	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
PES started >=3 days after randomisation	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Non-reported primary outcome	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Early/late reporting of primary outcome	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Non-reported SAE, SADE or USADE	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
...	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Total number of participants with >=1 violation	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)

† Participants may have one or more violations

NOTE: OTHERS WILL BE ADDED AS APPROPRIATE

Table S14. Method of follow-up at day 90

Method	All (N=xxx)	PES (n1=xxx)	No PES (n2=xxx)
Phone, completed by participant	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Phone, completed by carer	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Phone/email, completed by site (for patients still in hospital)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Postal, completed by participant	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Postal, completed by carer	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Refused follow-up	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Lost to follow-up	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
Died	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)
...	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)

NOTE: OTHERS WILL BE ADDED AS APPROPRIATE

Table S15. Analysis of the study within a trial (SWAT) assessment of undertreatment in participants randomised to PES.

PES group only	All	SWAT	No SWAT	Descriptors
Sites				
Sites, UK (N)		S1 xx	S2 xx	
Stroke volume <1000 pa (%)		xx (xx.x)	xx (xx.x)	N (%)
Median [IQR]		xxx.x [xxx.x, xxx.x]	xxx.x [xxx.x, xxx.x]	Median [IQR]
No prior PES experience (%)		xx (xx.x)	xx (xx.x)	N (%)
Participants				
Participants (N)	xxx	N1 xx	N2 xx	-
Age (yrs)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Mean (SD)
Sex, female (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)	N (%)
PES (mA), day 1, mean (SD)				
Threshold (mA)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Mean (SD)
Tolerability (mA)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Mean (SD)
Calculated stimulation (mA)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Mean (SD)
Actual stim, days 1-3 (mA)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Mean (SD)
Stim <20 mA, days 1-3 (%)	xx (xx.x)	xx (xx.x)	x.x (x.x, x.x), p	Unpooled proportion test
Time (secs)	xxx.x [xxx.x, xxx.x]	xxx.x [xxx.x, xxx.x]	xxx.x [xxx.x, xxx.x]	Median [IQR]
Time < 9m 50s, days 1-3 (%)	xx (xx.x)	xx (xx.x)	x.x (x.x, x.x), p	Unpooled proportion test
No. catheters >1 (%)	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x], p	Poisson regression
	xx (xx.x)	xx (xx.x)	x.x (x.x, x.x), p	Unpooled proportion test
Sensitivity				
Stimulation, days 1-6 (mA)	xx.x (xx.x)	xx.x (xx.x)	x.x (x.x, x.x), p	Unpooled t-test
Stim <20 mA, days 1-6 (%)	xx (xx.x)	xx (xx.x)	x.x (x.x, x.x), p	Unpooled proportion test
Time < 9m 50s, days 1-6 (%)	xx (xx.x)	xx (xx.x)	x.x (x.x, x.x), p	Unpooled proportion test
No. of SWAT instances	-	xx	-	-
Participants with a SWAT event	-	xx/N1 (xx.x)	-	-


Table S16. Device deficiencies as reported by site investigators.

Data apply only to PES group and are number (%) or mean (standard deviation) with [range].

PES group only	N (%)
<i>Base-station</i>	
Software issue - system showing incomplete treatments and unable to increase threshold/tolerance	xxx (xx.x)
...	xxx (xx.x)
<i>Catheter insertion</i>	
Guidewire not removed (user error)	xxx (xx.x)
Treatment attempted but catheter in incorrect place (user error)	xxx (xx.x)
...	xxx (xx.x)
<i>Catheter</i>	
Catheter securing plaster found in catheter box	xxx (xx.x)
Catheter losing contact despite being in position	xxx (xx.x)
Naso-gastric tube slippage despite clip in place	xxx (xx.x)
Blocked catheter, needed to be removed	xxx (xx.x)
...	xxx (xx.x)

NOTE: OTHERS WILL BE ADDED AS APPROPRIATE

Figure S1. CONSORT diagram.



Note: Sites variably screened all stroke admissions, only those referred to SLTs with dysphagia or just those where patients were given an information sheet. As a results, screening data are incomplete and an underestimate; of those reported, XXX patients were screened.

Figure S2. Sankey diagram of change in DSRS between days 0 - 14 and days 14 - 90 shown by treatment group.

Figure S3. Swallowing impairment at 14 and 90 days in the two groups according to scores on the functional oral intake scale.

Shift	Score	0	1	2	3	4	5	6	7
	Day								
PES	00	0	xx	xx	xx	xx	xx	xx	xx
	14	xx	xx	xx	xx	xx	xx	xx	xx
	90	xx	xx	xx	xx	xx	xx	xx	xx
No PES	90	xx	xx	xx	xx	xx	xx	xx	xx
	14	xx	xx	xx	xx	xx	xx	xx	xx
	00	0	xx	xx	xx	xx	xx	xx	xx

The functional oral outcome scale (FOIS) evaluates swallowing impairment with scores ranging from 1 to 7 with lower scores indicating more severe swallowing impairment;²⁵ inclusion criteria were: 0 – death, 1 – nil by mouth, 2 – tube dependent with minimal attempts of food or liquid, 3 – tube dependent with consistent oral intake of food or liquid.

Figure S4. Subgroup analyses of the primary outcome within the PES group.

Shown are the difference in means for swallowing impairment at day 90 in pre-specified subgroups of stimulation levels in milliamps. The widths of the confidence intervals have not been adjusted for multiplicity and should not be used in place of hypothesis testing. The analysis follows that of PHADER.¹⁶

- Threshold, average: <10, 10-13, >13
- Stimulation, average: <30, 30-36, >36
- Tolerability, average: <35, 35-45, >45
- Delta, tolerability – threshold, average: <24, 24-33, >33

PhEAST SAP

11/08/2026

Version 0.871

SUPPLEMENT 4

BASELINE PAPER

Title

Baseline characteristics of participants enrolled into the pharyngeal electrical stimulation for acute stroke dysphagia trial (PhEAST) trial.

Authors

TBD

Aim

Describe the baseline characteristics of participants enrolled into the PhEAST trial.

Participants

All in PhEAST

Outcomes

None.

Analyses

Descriptive statistics: number (%), median [interquartile range] or mean (standard deviation).

Table 1. Baseline characteristics for all recorded variables at baseline excluding cognition.

Data are number (%), median [interquartile range] or mean (standard deviation) with minimum/maximum.

Characteristic	N	All	Range
Demographics			
Age (years) #†	xxx	xx.x (xx.x)	xxx, xxx
Sex (%) †			/
Female	xxx	xxx (xx.x)	
Male	xxx	xxx (xx.x)	
Country of recruitment (%) ‡			/
Austria	xxx	xxx (xx.x)	
Denmark	xxx	xxx (xx.x)	
UK †	xxx	xxx (xx.x)	
Ethnicity (%)			/
Asian, south	xxx	xxx (xx.x)	
Asian, east/south-east	xxx	xxx (xx.x)	
Black	xxx	xxx (xx.x)	
White	xxx	xxx (xx.x)	
Onset to randomisation (days) #†	xxx	xx.x [xx.x, xx.x]	xx, xx
Swallowing/feeding scales			
DSRS (/12) † [1]	xxx	xx.x (xx.x)	xx, xx
FOIS [/7] (%) # [2]	xxx	x.x [x.x, x.x]	xx, xx
1 (%)	xxx	xxx (xx.x)	
2 (%)	xxx	xxx (xx.x)	
3 (%)	xxx	xxx (xx.x)	
IDDSI			
Food (3-7) [3]		x.x [x.x, x.x]	x, x
Drinks (0-4) [3]		x.x [x.x, x.x]	x, x
Functional diet scale (0-8) [4]		x.x (x.x)	x, x
Feeding status scale (/6) (%) [5]	xxx	x.x [x.x, x.x]	x, x
Normal oral diet		xxx (xx.x)	
Soft oral diet		xxx (xx.x)	
Nasogastric tube fed		xxx (xx.x)	
PEG/RIG		xxx (xx.x)	
Intravenous/subcutaneous		xxx (xx.x)	
Nothing		xxx (xx.x)	
EAT-10 (/50) [6]	xxx	xx.x (xx.x)	xx, xx
Speech & language therapy assessment			
Clinical assessment (%)			/
Dysphonia	xxx	xxx (xx.x)	
Dysarthria	xxx	xxx (xx.x)	
Abnormal gag reflex	xxx	xxx (xx.x)	

PhEAST SAP

11/08/2026

Version 0.871

Characteristic	N	All	Range
Abnormal volitional cough	xxx	xxx (xx.x)	
Cough after swallow	xxx	xxx (xx.x)	
Voice change after swallow	xxx	xxx (xx.x)	
Calculated PRESS score † [7]	xxx	xx.x (xx.x)	x, x
Calculated aspiration score [6]	xxx	x.x [x.x, x.x]	x, x
Instrumental assessment (%)			/
FEES	xxx	xxx (xx.x)	
Videofluoroscopy	xxx	xxx (xx.x)	
Total assessments	xxx	xxx (xx.x)	
Penetration aspiration scale <i>f</i>	xxx	x.x [x.x, x.x]	x, x
Clinical			
modified Rankin scale, pre-morbid (/6)	xxx	x.x [x.x, x.x]	x, x
modified Rankin scale, baseline (/6)	xxx	x.x [x.x, x.x]	x, x
Previous stroke (%)	xxx	xxx (xx.x)	/
NIHSS (/42) † [8]	xxx	xx.x (xx.x)	xx, xx
Barthel index (/100)	xxx	xx.x (xx.x)	xx, xx
Stroke type (%) †	xxx		/
Ischaemic		xxx (xx.x)	
Haemorrhagic		xxx (xx.x)	
Stroke syndrome (%) ⁷¹	xxx		/
Total anterior circulation (TACS) †		xxx (xx.x)	
Partial anterior circulation (PACS)		xxx (xx.x)	
Posterior circulation (POCS)		xxx (xx.x)	
Lacunar syndrome (LACS)		xxx (xx.x)	
Vascular circulation, posterior (%) †	xxx	xxx (xx.x)	/
Anterior		xxx (xx.x)	
Posterior		xxx (xx.x)	
Lesion location (%)	xxx		/
Right		xxx (xx.x)	
Left		xxx (xx.x)	
Bilateral		xxx (xx.x)	
Infratentorial		xxx (xx.x)	
Weight (kg)	xxx	xx.x (xx.x)	xxx.x, xxx.x
Height (m)	xxx	x.xx (x.xx)	x.xx, x.xx
Body mass index (kg/m/m)	xxx	xx.x (xx.x)	xx.x, xx.x
Neuroimaging, reported by site (%)			
Normal scan		xxx (xx.x)	
Infarct		xxx (xx.x)	
Infarct with HTI		xxx (xx.x)	
ICH		xxx (xx.x)	
SAH		xxx (xx.x)	
Other lesion		xxx (xx.x)	

Characteristic	N	All	Range
<i>In-hospital care (%)</i>	xxx		/
Intravenous thrombolysis (IVT)		xxx (xx.x)	
Mechanical thrombectomy (MT)		xxx (xx.x)	
IVT and/or MT		xxx (xx.x)	
Neurosurgery		xxx (xx.x)	
Vascular surgery		xxx (xx.x)	
ICU admission		xxx (xx.x)	
Ventilation		xxx (xx.x)	
Tracheostomy		xxx (xx.x)	

DSRS (dysphagia rating scale); EAT-10: eating assessment tool-10; FOIS: functional oral intake score; IDDSI: international dysphagia diet standardisation initiative; IQR: interquartile range; IVT: intravenous thrombolysis; MT: mechanical thrombectomy; NIHSS (NIH stroke severity scale); OSCP: Oxfordshire community stroke project; PEG: percutaneous endoscopic gastrostomy; RIG: radiologically inserted gastrostomy; SD: standard deviation; TACS: total anterior circulation syndrome.

Inclusion criterion

† Minimisation variables: age (<75/75+), sex (female/male), DSRS (<12/12), impairment (NIHSS <15/15+), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior), time to randomisation (<15/15+ days), country (other than UK/UK).

‡ Country other than UK vs UK was a stratification variable but this was changed to a minimisation variable once it became clear that recruitment would be low outside of the UK.

1. The dysphagia severity rating scale (DSRS) evaluates swallowing impairment with scores ranging from 0 (normal) to 12 with higher scores indicating more severe swallowing impairment.^{12,26} The total score is the sum of three subscale scores relating to fluids, food and supervision, each ranging from 0 (normal) to 4 (nil by mouth).
2. The functional oral outcome scale (FOIS) evaluates swallowing impairment with scores ranging from 1 to 7 with lower scores indicating more severe swallowing impairment;²⁵ inclusion criteria were: 1 – nil by mouth, 2 – tube dependent with minimal attempts of food or liquid, 3 – tube dependent with consistent oral intake of food or liquid.
3. The international dysphagia diet standardisation initiative (IDDSI) is a global standardised framework describing food texture and liquid thickening;⁴⁹ food texture scores range between 7 (regular) and 3 (liquidised); fluid thickening scores range between 0 (thin) and 4 (extremely thick).
4. The international dysphagia diet standardisation initiative - functional diet scale (IDDSI-FDS) evaluates the severity of oropharyngeal dysphagia represented by diet texture restriction with scores ranging from 0 (nil by mouth) to 8 (normal food texture and drink consistency).⁵⁰
5. The feeding status scale (FSS) is a measure of nutritional status with scores 0 (normal oral diet), 1 (soft diet), 2 (nasogastric tube feeding), 3 (PEG/RIG/J-tube feeding), 4 (parenteral fluids) and 5 (no fluids/feeding).⁴⁷*Ref Non-oral feeding is a score that exceeds 1.
6. The eating assessment tool-10 (EAT-10) is a measure of swallowing difficulties comprising 10 questions with each score ranging for 0 (no problem) to 4 (severe problem);⁵¹ the scale is extended to include a score of 5 for nil by mouth hence the total score ranges from 0 to 50.

7. The predictive swallowing score (PRESS) is a prognostic instrument that predicts swallowing recovery after stroke and ranges from 0 (no features present) to 10 (10 features present) based on age ≥ 70 years; NIHSS score; stroke lesion includes lesion at the frontal operculum; presence of dysphonia, dysarthria, abnormal gag reflex, abnormal volitional cough, cough after swallow and voice change after swallow; and FOIS score.⁶¹ Brain scans before or at baseline were not collected and adjudicated. Instead, relevant neuroradiology reports were entered by recruiting sites but few made any specific comment about the presence of a stroke lesion at the frontal operculum. Hence, this item may have been under-reported and so the total PRESS score is likely an underestimate.
8. The National Institutes of Health stroke scale (NIHSS) evaluates stroke severity/impairment with scores ranging from 0 to 42 with higher scores indicating greater neurological impairment.⁶⁷

Table 2. Sources of consent accounting for country of randomisation and type of consent.

Variable	All
Participants with data	N
Participant consent	xxx (xx.x)
Consultee declaration: England, Denmark, Northern Ireland, Wales	xxx (xx.x)
Personal legal representative: Austria, Scotland	xxx (xx.x)
Independent physician	xxx (xx.x)

Table 3. Co-enrolment into another trial.

Variable	PhEAST order	N
Participants		xxx
ENOS-2 (ISRCTN17654248)	Second	xxx (xx.x)
MAPS-2 (ISRCTN40512746)	Second	xxx (xx.x)
RECAST-3 (ISRCTN63231313)	Second	xxx (xx.x)
TICH-3 (ISRCTN97695350)	Second	xxx (xx.x)
ProFATE ⁷²	Second	xxx (xx.x)
GEKO (ISRCTN11175235)	Second	xxx (xx.x)
...		
...		

Table 4. Univariate correlations between baseline characteristics for age, sex, swallowing severity (DSRS), stroke severity (NIHSS), stroke type (ischaemic, haemorrhagic), time for onset to randomisation (OTR), and year of randomisation.

Correlations by point biserial or Spearman's tests; data are regression coefficient (*P* value).

	Age	Sex	DSRS	FOIS	IDDSI-RDS	EAT-10	NIHSS	IS/ICH	OTR	Year
Age	X	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)
Sex	r (p)	X	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)
DSRS	r (p)	r (p)	X	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)
FOIS	r (p)	r (p)	r (p)	X	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)
IDDSI-RDS	r (p)	r (p)	r (p)	r (p)	X	r (p)	r (p)	r (p)	r (p)	r (p)
EAT-10	r (p)	r (p)	r (p)	r (p)	r (p)	X	r (p)	r (p)	r (p)	r (p)
NIHSS	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	X	r (p)	r (p)	r (p)
IS/ICH	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	X	r (p)	r (p)
OTR	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	X	r (p)
Year	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	r (p)	X

Figure 1. Cumulative recruitment of participants and sites during the trial.

Nb: Place the site line below the participant line.

Figure 2. Monthly recruitment of participants and sites during the trial.

Nb: Bars for each month

Figure 3. Histogram of time from stroke onset to randomisation (days).

Nb: Bars for each day 2-31.

Figure 4. Histogram of recruitment by sites.

Nb: Bars for each site in order of recruitment with sites identified as A, B, C...

SUPPLEMENT 5

Title

Cost effectiveness of pharyngeal electrical stimulation for post stroke dysphagia as compared to usual care. Data from the PhEAST trial.

Authors

Cristina Roadevin, ..., Philip M Bath, Marilyn James.

Aim

Evaluate the cost effectiveness of the PhEAST trial intervention (PES) compared to usual care.

Participants

All in PhEAST

Outcomes

Primary: Health-related quality of life (HRQoL) for participant: EuroQoL-5D-5L (EQ-5D-5L) administered at baseline, and days: 14, 90.

Secondary: *Cristina

Analyses

Descriptive statistics: number (%), median [interquartile range] or mean (standard deviation).

Cost-effectiveness assessed through incremental cost-effectiveness ratios (ICERs) and incremental net monetary benefits (INMBs).

Exemplar tables follow.

Table 1 Estimated and imputed average QALYs per patient at 12 months.

Data are mean (standard error, SE).

	PES Mean (SE)	Usual Care Mean (SE)	Difference in means between groups (SE)
QALYs	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)

SE: standard error

Table 2 Average costs per patient and difference in means between groups at 12 months

	PES + usual care (n=)		Usual Care (n=)		Difference in means between groups (£)
	Mean (£)	SE	Mean (£)	SE	Mean (SE)
Health service perspective costs	xx.x	xx.x	xx.x	xx.x	xx.x
Intervention cost	xx.x	xx.x	xx.x	xx.x	xx.x
Inpatient services	xx.x	xx.x	xx.x	xx.x	xx.x
Outpatient services	xx.x	xx.x	xx.x	xx.x	xx.x
Primary and community care	xx.x	xx.x	xx.x	xx.x	xx.x
Medication	xx.x	xx.x	xx.x	xx.x	xx.x
Aids and Equipment	xx.x	xx.x	xx.x	xx.x	xx.x
Broader Costs	xx.x	xx.x	xx.x	xx.x	xx.x
Informal care	xx.x	xx.x	xx.x	xx.x	xx.x
Aids and Equipment	xx.x	xx.x	xx.x	xx.x	xx.x
Total costs	xx.x	xx.x	xx.x	xx.x	xx.x
Health Service Perspective Costs	xx.x	xx.x	xx.x	xx.x	xx.x
Societal Perspective Costs	xx.x	xx.x	xx.x	xx.x	xx.x

Table 3 Cost Effectiveness Results.

	Incremental costs (95% CI)	Incremental QALYs (95% CI)	ICER (95% CI)
NHS perspective	xx.x (xx.x, xx.x)	xx.x (xx.x, xx.x)	xx.x (xx.x, xx.x)
Societal perspective	xx.x (xx.x, xx.x)	xx.x (xx.x, xx.x)	xx.x (xx.x, xx.x)

Table 4 Net Monetary Benefit Results

WTP thresholds	£20,000	£30,000	£40,000	£50,000	£60,000	£100,000
NMB values	xx.x	xx.x	xx.x	xx.x	xx.x	xx.x

SUPPLEMENT 6

SWALLOWING/DYSPHAGIA UP TO ONE YEAR

Title

Trajectory of swallowing over 1 year in patients with severe ischaemic or haemorrhagic stroke and dysphagia. Data from the PhEAST trial.

Authors

TBD

Aim

Describe the trajectory of swallowing at 5 time points over 1 year in patients with severe ischaemic or haemorrhagic stroke and dysphagia defined by tube-feeding at baseline. Compare swallowing impairment by PES versus no PES.

Participants

All in PhEAST

Outcomes

Primary: DSRS.

Secondary: At all timepoints – FOIS, FSS, IDDSI, IDDSI-FDS, NGT or PEG in situ, death, discharge disposition.

Others: mRS, Barthel, EQ-5D, EQ-VAS, global outcome

Analyses

Descriptive statistics: number (%), median [interquartile range] or mean (standard deviation).

Comparison by adjusted repeated measures regression.

Data will be tabulated in all patients whether alive for not, and in alive patients only.

The analyses will follow those in the main paper but cover outcomes out to one year.

Analyses present in the main paper but not of relevance at one year, e.g. PES stimulation levels, will not be re-presented.

Table 1. Baseline characteristics by baseline FOIS score.

Characteristic	Statistic	N	All (N=xxx)	FOIS=1 (N=xxx)	FOIS=2 (N=xxx)	FOIS=3 (N=xxx)	2p trend
Demographics							
Age (years) #†	Mean (SD)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	Jonckheere test
Sex, female (%)	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Country, other than UK	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Ethnicity, other than white (%)	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Onset to randomisation (days) #†	Median [IQR]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	Jonckheere test
Swallowing							
DSRS (/12) †	Mean (SD)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Jonckheere test
FOIS [/7] #	Median [IQR]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	Jonckheere test
IDDSI							
Food (3-7)	Median [IQR]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	Jonckheere test
Drinks (0-4)	Median [IQR]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	Jonckheere test
Functional diet scale (0-8)	Mean (SD)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Jonckheere test
Feeding status scale, FSS>1 (%)	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
EAT-10 (/50)	Mean (SD)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Jonckheere test
Clinical							
Modified Rankin scale, premorbid [/6]	Median [IQR]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	Jonckheere test
Modified Rankin scale, baseline [/6]	Median [IQR]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	Jonckheere test
Previous stroke	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
NIHSS (/42) †	Mean (SD)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	Jonckheere test
Barthel index (/100)	Mean (SD)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	Jonckheere test
Stroke type, haemorrhage (%) †	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Stroke syndrome, TACS (%) †	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Vascular circulation, posterior (%) †	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2

Characteristic	Statistic	N	All (N=xxx)	FOIS=1 (N=xxx)	FOIS=2 (N=xxx)	FOIS=3 (N=xxx)	2p trend
<i>In-hospital care pre-randomisation</i>	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	
Intravenous thrombolysis	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Mechanical thrombectomy	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Hemicraniectomy	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Intensive care unit admission	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2
Ventilation	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	CAT 3x2

DSRS (dysphagia rating scale); EAT-10: eating assessment tool-10; FOIS: functional oral intake score; FSS: feeding status scale; IDDSI: international dysphagia diet standardisation initiative; NIHSS (NIH stroke severity scale); OSCP: Oxfordshire community stroke project; TACS: total anterior circulation syndrome.

Data are number (%), median [interquartile range] or mean (standard deviation). Comparisons by Cochrane-Armitage test (CAT), or Jonckheere test for ordered alternatives.

Inclusion criterion

† Minimisation variables: age (<75/75+), sex (female/male), DSRS (<12/12), impairment (NIHSS <15/15+), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior), time to randomisation (<15/15+ days), country (other than UK/UK).

Table 2. Outcomes at days 14, 90, 180 and 365 in all participants.

Outcome	N	14	90	180	365	aDIM/aDIMed/ acOR/aOR/aHR (95% CI)+	Analysis
Participants		xxx	xxx	xxx	xxx		
DSRS (/13) ‡ PES No PES	xxx	xx.x (x.x) xx.x (x.x) aMLR	xx.x (x.x) xx.x (x.x) aMLR	xx.x (x.x) xx.x (x.x) aMLR	xx.x (x.x) xx.x (x.x) aMLR	xx.x (xx.x, xx.x)	acRMR / / /
FOIS [/7] ‡ PES No PES	xxx	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	xx.x [xx.x, xx.x]	aoRMR / / /
Feeding status scale [/7] ‡ PES No PES	xxx	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	xx.x [xx.x, xx.x]	aoRMR / / /
IDDSI-FDS [/8] ‡ PES No PES	xxx	x.x [x.x, x.x] x.x [x.x, x.x] aMLR	x.x [x.x, x.x] x.x [x.x, x.x] aMLR	x.x [x.x, x.x] x.x [x.x, x.x] aMLR	x.x [x.x, x.x] x.x [x.x, x.x] aMLR	xx.x [xx.x, xx.x]	acRMR / / /
Global swallowing ✕		/	/	/	/		
NGT or PEG/RIG in situ (%) PES No PES	xxx	xxx (xx.x) xxx (xx.x) aBLR	xxx (xx.x) xxx (xx.x) aBLR	xxx (xx.x) xxx (xx.x) aBLR	xxx (xx.x) xxx (xx.x) aBLR	x.xx (xx.x-xx.x)	aoRMR / / /
Pneumonia, fatal (%) PES No PES	xxx	xxx (xx.x) xxx (xx.x) aBLR	xxx (xx.x) xxx (xx.x) aBLR	xxx (xx.x) xxx (xx.x) aBLR	xxx (xx.x) xxx (xx.x) aBLR	x.xx (xx.x-xx.x)	aCPHR / / /
Hospital length of stay (days) PES No PES	xxx	xxx (xx.x) xxx (xx.x)	xxx (xx.x) xxx (xx.x)	xxx (xx.x) xxx (xx.x)	xxx (xx.x) xxx (xx.x)	x.xx (xx.x-xx.x)	aMLR / /

Death (%) PES No PES	xxx	xxx (xx.x) xxx (xx.x) aBLR	xxx (xx.x) xxx (xx.x) aCPHR	xxx (xx.x) xxx (xx.x) aCPHR	xxx (xx.x) xxx (xx.x) aCPHR	x.xx (xx.x-xx.x)	aCPHR
Modified Rankin Scale [/6] PES No PES	xxx	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.x [x.x, x.x] x.x [x.x, x.x] aOLR	x.xx (xx.x-xx.x)	aoRMR
Barthel index [/100] PES No PES	xxx	x.x [x.x, x.x] x.x [x.x, x.x] aMLR	x.x [x.x, x.x] x.x [x.x, x.x] aMLR	x.x [x.x, x.x] x.x [x.x, x.x] aMLR	x.x [x.x, x.x] x.x [x.x, x.x] aMLR	x.xx (xx.x-xx.x)	acRMR
Clinical frailty index [/9] ⁷³	xxx					x.xx (xx.x-xx.x)	aoRMR
PES		x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]		
No PES		x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]		
		aOLR	aOLR	aOLR	aOLR		
EQ-5D-5L health utility (/1) ¶	xxx	0.xx (0.xx) 0.xx (0.xx) aMLR	0.xx (0.xx) 0.xx (0.xx) aMLR	0.xx (0.xx) 0.xx (0.xx) aMLR	0.xx (0.xx) 0.xx (0.xx) aMLR	xx.x (xx.x, xx.x)	acRMR
EQ-VAS (/100) ¶	xxx	0.xx (0.xx) 0.xx (0.xx) aMLR	0.xx (0.xx) 0.xx (0.xx) aMLR	0.xx (0.xx) 0.xx (0.xx) aMLR	0.xx (0.xx) 0.xx (0.xx) aMLR	xx.x (xx.x, xx.x)	acRMR
Living in an institution (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)		x.xx (xx.x-xx.x)	aBLR

Data are number (%), median [interquartile range or mean (standard deviation)]. Comparisons by binary logistic regression, Cox proportional hazards regression, ordinal logistic regression, multiple linear regression, or repeated measures equivalents. The models include randomisation group with adjustment for baseline score, age, sex, DSRS, impairment (NIHSS), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior) and time to randomisation (days).

✕ Global analysis of DSRS, FOIS, FSS, IDDSI functional diet scale and NGT/PEG placement, using the Wei-Lachin test.

Table 3. Time to event analyses for length of stay and removal of enteral feeding.

Outcome	N	All	PES	No PES	aHR (95% CI)[†]	Analysis
Participants		xxx	xxx	xxx		
Died (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	x.xx (x.xx-x.xx)	aCPHR
Length of stay in hospital (%) ‡	xxx	xxx [xx.x, xx.x]	xxx [xx.x, xx.x]	xxx [xx.x, xx.x]	x.xx (x.xx-x.xx)	aCPHR

‡ To home or institution

PhEAST SAP

11/08/2026

Version 0.871

SUPPLEMENT 7

COGNITION/DEMENTIA UP TO ONE YEAR

Title

Trajectory of cognition over 1 year in patients with severe ischaemic or haemorrhagic stroke. Data from the PhEAST trial.

Authors

TBD

Aim

Describe the trajectory of cognition at 5 time points over 1 year in patients with severe ischaemic or haemorrhagic stroke.

Participants

All in PhEAST

Outcomes

Primary: DSM-5 7-level ordinal cognition.

Secondary: At days 0 and 14 – NIHSS-Cog-4, MoCA and Free-Cog

Secondary: At all timepoints – DSM-5 4-level, t-MoCA, TICS-M, t-Free-Cog, 6-CIT, phenomic fluency and semantic fluency.

Informant: IQCODE.

Others: Clinical frailty index, Zung depression index.

Analyses

Descriptive statistics: number (%), median [interquartile range] or mean (standard deviation).

Data will be tabulated in all patients whether alive for not, and in alive patients only.

Table 1. Baseline characteristics.

Data are number (%), median [interquartile range or mean (standard deviation)].

Characteristic	Statistic	N	All	Cognition data †	No Cognition data †	2p
Participants	N		xxx	With DSM-5	Without DSM-5	
Demographics						
Age (years) †	Mean (SD)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	MLR
Sex, female (%)	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	BLR
Ethnicity, non-white (%)	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	BLR
Onset to randomisation (days) †	Median [IQR]	xxx	xxx [xxx]	xxx [xxx]	xxx [xxx]	OLR
Clinical						
Previous stroke	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	BLR
Dysarthria	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	BLR
NIHSS (/42) ¶	Mean (SD)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	MLR
NIHSS Cog-4 [/9] §	Median [IQR]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	OLR
Modified Rankin Scale [/6]	Median [IQR]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	x.x [x.x, x.x]	OLR
Barthel index (/100)	Mean (SD)	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	MLR
OCSF, TACS (%)	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	BLR
Stroke type, haemorrhage (%)	n (%)	xxx	xxx (xx.x)	xxx (xx.x)	xxx (xx.x)	BLR

NIHSS (NIH stroke severity scale); OCSF: Oxfordshire community stroke project; TACS: total anterior circulation syndrome.

† The group with cognition information are those with data for the DSM-5 7 level scale.

¶ Scores on the National Institutes of Health Stroke Scale (NIHSS) range from 0 to 42, with higher scores indicating greater neurological impairment.

§ Scores on the NIHSS Cog-4 scale range from 0-9, with higher scores indicating worse cognition.⁷⁴

Table 2. Cognition variables at baseline and days 14, 90, 180 and 365.

Results include participants who died with death assigned a score worse than any living score.

Outcome	N	0	14	90	180	365	Treatment effect	Test
Participants		xxx	xxx	xxx	xxx	xxx		
NIHSS COG-4 [/9] ⁷⁴ §	xxx						xx.x (xx.x, xx.x)	aoRMR
PES		x.x (x.x)	x.x (x.x)	x.x (x.x)	NR	NR		
No PES		x.x (x.x) /	x.x (x.x) aMLR	x.x (x.x) aMLR	NR	NR		
MOCA (/30) ⁷⁵	xxx						xx.x [xx.x, xx.x]	acRMR
PES		xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	NR	NR		
No PES		xx.x (x.x) (N) /	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	NR	NR		
t-MOCA (/22) *	xxx						xx.x [xx.x, xx.x]	acRMR
PES		xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)		/
No PES		xx.x (x.x) (N) /	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR		/
Phonemic fluency (Letter F) ⁷⁶	xxx						xx.x [xx.x, xx.x]	acRMR
PES		xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)		/
No PES		xx.x (x.x) (N) /	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR		/
Category/semantic fluency (animal naming) ⁷⁶	xxx						x.xx (xx.x-xx.x)	acRMR
PES		xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)		

PhEAST SAP		11/08/2026		Version 0.871				
No PES		xx.x (x.x) (N) /	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR		
TICS-m (/39) ⁷⁷ PES	xxx	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)	x.xx (xx.x-xx.x)	acRMR /
No PES		xx.x (xx.x) (N) /	xx.x (xx.x) (N) aMLR	xx.x (xx.x) (N) aMLR	xx.x (xx.x) (N) aMLR	xx.x (xx.x) (N) aMLR		/
DSM5-4 level [/3] ⁶⁰ PES	xxx	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.xx (xx.x-xx.x)	aoRMR /
No PES		x.x [x.x, x.x] (N) /	x.x [x.x, x.x] (N) aOLR	x.x [x.x, x.x] (N) aOLR	x.x [x.x, x.x] (N) aOLR	x.x [x.x, x.x] (N) aOLR		/
DSM5-7 level [/6] ⁶⁰ PES	xxx	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.xx (xx.x-xx.x)	aoRMR /
No PES		x.x [x.x, x.x] (N) /	x.x [x.x, x.x] (N) aOLR	x.x [x.x, x.x] (N) aOLR	x.x [x.x, x.x] (N) aOLR	x.x [x.x, x.x] (N) aOLR		/
Free-Cog (/30) ⁷⁸ PES	xxx	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	x.xx (xx.x-xx.x)	acRMR
No PES		xx.x (x.x) (N) /	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR	xx.x (x.x) (N) aMLR		
Tele Free-Cog (/24) ⁷⁸ PES	xxx	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	x.xx (xx.x-xx.x)	acRMR
No PES		xx.x (x.x)	xx.x (x.x)	xx.x (x.x)	xx.x (x.x)	xx.x (x.x)		

PhEAST SAP

11/08/2026

Version 0.871

		(N) /	(N) aMLR	(N) aMLR	(N) aMLR	(N) aMLR		
6-CIT [/28] ⁷⁹ PES No PES	xxx	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) /	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) aMLR	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) aMLR	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) aMLR	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) aMLR	xx.x (xx.x, xx.x)	acRMR
Zung depression scale (/100) ⁸⁰ PES No PES	xxx	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) /	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) aMLR	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) aMLR	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) aMLR	x.x [x.x, x.x] (N) x.x [x.x, x.x] (N) aMLR	xx.x (xx.x, xx.x)	acRMR
In institution or died (%)	xxx						x.xx (xx.x-xx.x)	abRMR
PES		xxx (xx.x) (N)	xxx (xx.x) (N)	xxx (xx.x) (N)	xxx (xx.x) (N)	xxx (xx.x) (N)		
No PES		xxx (xx.x) (N)	xxx (xx.x) (N)	xxx (xx.x) (N)	xxx (xx.x) (N)	xxx (xx.x) (N)		
		/	aBLR	aBLR	aBLR	aBLR		

Indicted scores for death: NIHSS Cog-4 10, MoCA -1, t-MoCA -1, letter F -1, TICS-m -1, DSM-5 4-level 3, DSM-5 7-level 6, animal naming -1, free-cog -1, t-free-cog -1, 6-CIT 29, clinical frailty 10, Zung depression 102.5.

Table 3. Cognition variables at baseline and days 14, 90, 180 and 365. *Philip

Results exclude participants who died by the relevant timepoint.

Measure					
Day	0	14	90	180	365
Participants	N	N	N	N	N
NIHSS COG-4 [/9] ⁷⁴ §	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	/	/	/
MOCA (/30) ⁷⁵	xx.x (x.x) (N)	xx.x (x.x) (N)	/	/	/
t-MOCA (/22) *	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)
Phonemic fluency (Letter F) ⁷⁶	x.x (x.x) (N)	x.x (x.x) (N)	x.x (x.x) (N)	x.x (x.x) (N)	x.x (x.x) (N)
TICS-m (/39) ⁷⁷	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)
Cog-R4VaD-4 level [/6] ⁶⁰	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)
Cog-R4VaD-7 level [/6] ⁶⁰	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)
Category/semantic fluency (animal naming) ⁷⁶	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)
Free-Cog (/30) ⁷⁸	xx.x (x.x) (N)	xx.x (x.x) (N)	/	/	/
Tele Free-Cog (/24) ⁷⁸	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)	xx.x (x.x) (N)
6-CIT [/28] ⁷⁹	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)
Clinical frailty index [/9] ⁷³	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)	x.x [x.x, x.x] (N)
Zung depression scale (/100) ⁸⁰	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)	xx.x (xx.x) (N)
IQCODE	xx.x (x.x)	xx.x (x.x)	xx.x (x.x)	xx.x (x.x)	xx.x (x.x)

Note: Data are median [interquartile range] (N) or mean (standard deviation) (N).

SUPPLEMENT 8

RESPONDER ANALYSIS

Title

Comparison of patients with severe dysphagia who responded to versus those who did not respond to pharyngeal electrical stimulation. Data from the PhEAST trial.

Authors

TBD

Aim

Compare baseline characteristics and outcomes in people who responded versus those who did not respond to PES.

Participants

All those randomised to PES.

Definition of response

At day 90 follow-up: DSRS <4 (vs ≥ 4).

Outcomes

Groups: DSRS at day 90 0 or 1 versus 2 to 12 versus died

Baseline: demographic and other characteristics

Follow-up: Swallowing and functional outcomes

Analyses

Descriptive statistics: number (%), median [interquartile range] or mean (standard deviation).

Data will be tabulated in three groups.

Table 1. Baseline characteristics by responder group at 90 days.

Data are number (%), median [interquartile range] or mean (standard deviation); comparison by unadjusted risk difference, difference in medians (Mann-Whitney u test) or difference in means (Welsh test) with 95% confidence intervals. Information on those who had died by day are given for completeness.

Characteristic	N	Responder DSRS<4 N=xxx	Non-responder DSRS>=4 N=xxx	Difference	Died N=xxx
Demographics					
Age (years) #†	xxx	xx.x (xx.x)	xx.x (xx.x)	DiMean (95% CI) Unpooled z test	xx.x (xx.x)
Sex (%) †					
Female	xxx	xxx (xx.x)	xxx (xx.x)	Chi-squared test	xxx (xx.x)
Male	xxx	xxx (xx.x)	xxx (xx.x)		xxx (xx.x)
Country of recruitment (%) ‡					
Austria	xxx	xxx (xx.x)	xxx (xx.x)		xxx (xx.x)
Denmark	xxx	xxx (xx.x)	xxx (xx.x)	Chi-squared test	xxx (xx.x)
UK †	xxx	xxx (xx.x)	xxx (xx.x)		xxx (xx.x)
Ethnicity (%)					
Asian, south	xxx	xxx (xx.x)	xxx (xx.x)		xxx (xx.x)
Asian, east/south-east	xxx	xxx (xx.x)	xxx (xx.x)	DiMed (95% CI)	xxx (xx.x)
Black	xxx	xxx (xx.x)	xxx (xx.x)		xxx (xx.x)
White	xxx	xxx (xx.x)	xxx (xx.x)		xxx (xx.x)
Onset to randomisation (days) #†	xxx	xx.x [xx.x, xx.x]	xx.x [xx.x, xx.x]		xx.x [xx.x, xx.x]
Swallowing/feeding scales					
DSRS (/12) † [1]	xxx	xx.x (xx.x)	xx.x (xx.x)	DiMean (95% CI)	
FOIS [/7] (%) # [2]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	DiMed (95% CI)	
1 (%)	xxx	xxx (xx.x)	xxx (xx.x)		

Characteristic	N	Responder DSRS<4 N=xxx	Non-responder DSRS>=4 N=xxx	Difference	Died N=xxx
2 (%)	xxx	xxx (xx.x)	xxx (xx.x)	DiMean (95% CI) DiMean (95% CI)	
3 (%)	xxx	xxx (xx.x)	xxx (xx.x)		
IDDSI					
Food (3-7) [3]		x.x [x.x, x.x]	x.x [x.x, x.x]		
Drinks (0-4) [3]		x.x [x.x, x.x]	x.x [x.x, x.x]		
Functional diet scale (0-8) [4]		x.x (x.x)	x.x (x.x)		
Feeding status scale (/6) (%) [5]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]		
Normal oral diet		xxx (xx.x)	xxx (xx.x)		
Soft oral diet		xxx (xx.x)	xxx (xx.x)		
Nasogastric tube fed		xxx (xx.x)	xxx (xx.x)		
PEG/RIG		xxx (xx.x)	xxx (xx.x)		
Intravenous/subcutaneous		xxx (xx.x)	xxx (xx.x)		
Nothing		xxx (xx.x)	xxx (xx.x)		
EAT-10 (/50) [6]	xxx	xx.x (xx.x)	xx.x (xx.x)	DiMean (95% CI)	
Speech & language therapy assessment					
Clinical assessment (%)					
Dysphonia	xxx	xxx (xx.x)	xxx (xx.x)		
Dysarthria	xxx	xxx (xx.x)	xxx (xx.x)		
Abnormal gag reflex	xxx	xxx (xx.x)	xxx (xx.x)		
Abnormal volitional cough	xxx	xxx (xx.x)	xxx (xx.x)		
Cough after swallow	xxx	xxx (xx.x)	xxx (xx.x)		
Voice change after swallow	xxx	xxx (xx.x)	xxx (xx.x)		
Calculated PRESS score † [7]	xxx	xx.x (xx.x)	xx.x (xx.x)		
Calculated aspiration score [6]	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]		

Characteristic	N	Responder DSRS<4 N=xxx	Non-responder DSRS>=4 N=xxx	Difference	Died N=xxx
Instrumental assessment (%)					
FEES	xxx	xxx (xx.x)	xxx (xx.x)		
Videofluoroscopy	xxx	xxx (xx.x)	xxx (xx.x)		
Total assessments	xxx	xxx (xx.x)	xxx (xx.x)		
Penetration aspiration scale <i>f</i>	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]		
Clinical					
modified Rankin scale, pre-morbid (/6)	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]	DiMean (95% CI)	
modified Rankin scale, baseline (/6)	xxx	x.x [x.x, x.x]	x.x [x.x, x.x]		
Previous stroke (%)	xxx	xxx (xx.x)	xxx (xx.x)		
NIHSS (/42) † [8]	xxx	xx.x (xx.x)	xx.x (xx.x)		
Barthel index (/100)	xxx	xx.x (xx.x)	xx.x (xx.x)		
Stroke type (%) †	xxx				
Ischaemic		xxx (xx.x)	xxx (xx.x)		
Haemorrhagic		xxx (xx.x)	xxx (xx.x)		
Stroke syndrome (%) ⁷¹	xxx				
Total anterior circulation (TACS) †		xxx (xx.x)	xxx (xx.x)		
Partial anterior circulation (PACS)		xxx (xx.x)	xxx (xx.x)		
Posterior circulation (POCS)		xxx (xx.x)	xxx (xx.x)		
Lacunar syndrome (LACS)		xxx (xx.x)	xxx (xx.x)		
Lesion location (%)	xxx				
Right		xxx (xx.x)	xxx (xx.x)		
Left		xxx (xx.x)	xxx (xx.x)		
Bilateral		xxx (xx.x)	xxx (xx.x)		
Infratentorial		xxx (xx.x)	xxx (xx.x)		
Weight (kg)	xxx	xx.x (xx.x)	xx.x (xx.x)		

Characteristic	N	Responder DSRS<4 N=xxx	Non-responder DSRS>=4 N=xxx	Difference	Died N=xxx
Height (m)	xxx	x.xx (x.xx)	x.xx (x.xx)		
Body mass index (kg/m/m)	xxx	xx.x (xx.x)	xx.x (xx.x)		
Neuroimaging, reported by site (%)	xxx				
Normal scan		xxx (xx.x)	xxx (xx.x)		
Infarct		xxx (xx.x)	xxx (xx.x)		
Infarct with HTI		xxx (xx.x)	xxx (xx.x)		
ICH		xxx (xx.x)	xxx (xx.x)		
SAH		xxx (xx.x)	xxx (xx.x)		
Other lesion		xxx (xx.x)	xxx (xx.x)		
In-hospital care (%)	xxx				
Intravenous thrombolysis		xxx (xx.x)	xxx (xx.x)		
Mechanical thrombectomy		xxx (xx.x)	xxx (xx.x)		
Combined IVT and MT		xxx (xx.x)	xxx (xx.x)		
Neurosurgery		xxx (xx.x)	xxx (xx.x)		
Vascular surgery		xxx (xx.x)	xxx (xx.x)		
ICU admission		xxx (xx.x)	xxx (xx.x)		
Ventilation		xxx (xx.x)	xxx (xx.x)		
Tracheostomy		xxx (xx.x)	xxx (xx.x)		

DSRS (dysphagia rating scale); EAT-10: eating assessment tool-10; FOIS: functional oral intake score; IDDSI: international dysphagia diet standardisation initiative; IQR: interquartile range; IVT: intravenous thrombolysis; MT: mechanical thrombectomy; NIHSS (NIH stroke severity scale); OSCP: Oxfordshire community stroke project; PEG: percutaneous endoscopic gastrostomy; RIG: radiologically inserted gastrostomy; SD: standard deviation; TACS: total anterior circulation syndrome.

Inclusion criterion

† Minimisation variables: age (<75/75+), sex (female/male), DSRS (<12/12), impairment (NIHSS <15/15+), stroke type (ischaemic/haemorrhagic), circulation (anterior/posterior), time to randomisation (<15/15+ days), country (other than UK/UK).

‡ Country other than UK vs UK was a stratification variable but this was changed to a minimisation variable once it became clear that recruitment would be low outside of the UK.

9. The dysphagia severity rating scale (DSRS) evaluates swallowing impairment with scores ranging from 0 (normal) to 12 with higher scores indicating more severe swallowing impairment.^{12,26} The total score is the sum of three subscale scores relating to fluids, food and supervision, each ranging from 0 (normal) to 4 (nil by mouth).
10. The functional oral outcome scale (FOIS) evaluates swallowing impairment with scores ranging from 1 to 7 with lower scores indicating more severe swallowing impairment;²⁵ inclusion criteria were: 1 – nil by mouth, 2 – tube dependent with minimal attempts of food or liquid, 3 – tube dependent with consistent oral intake of food or liquid.
11. The international dysphagia diet standardisation initiative (IDDSI) is a global standardised framework describing food texture and liquid thickening;⁴⁹ food texture scores range between 7 (regular) and 3 (liquidised); fluid thickening scores range between 0 (thin) and 4 (extremely thick).
12. The international dysphagia diet standardisation initiative - functional diet scale (IDDSI-FDS) evaluates the severity of oropharyngeal dysphagia represented by diet texture restriction with scores ranging from 0 (nil by mouth) to 8 (normal food texture and drink consistency).⁵⁰
13. The feeding status scale (FSS) is a measure of nutritional status with scores 0 (normal oral diet), 1 (soft diet), 2 (nasogastric tube feeding), 3 (PEG/RIG/J-tube feeding), 4 (parenteral fluids) and 5 (no fluids/feeding).⁴⁷ *Ref Non-oral feeding is a score that exceeds 1.
14. The eating assessment tool-10 (EAT-10) is a measure of swallowing difficulties comprising 10 questions with each score ranging from 0 (no problem) to 4 (severe problem);⁵¹ the scale is extended to include a score of 5 for nil by mouth hence the total score ranges from 0 to 50.
15. The predictive swallowing score (PRESS) is a prognostic instrument that predicts swallowing recovery after stroke and ranges from 0 (no features present) to 10 (10 features present) based on age ≥ 70 years; NIHSS score; stroke lesion includes lesion at the frontal operculum; presence of dysphonia, dysarthria, abnormal gag reflex, abnormal volitional cough, cough after swallow and voice change after swallow; and FOIS score.⁶¹ Brain scans before or at baseline were not collected and adjudicated; relevant neuroradiology reports were entered by recruiting sites but they usually do not make any specific comment about the presence of a stroke lesion at the frontal operculum. Hence, this item has been under-reported and so the total PRESS score will be an underestimate.
16. The National Institutes of Health stroke scale (NIHSS) evaluates stroke severity/impairment with scores ranging from 0 to 42 with higher scores indicating greater neurological impairment.⁶⁷

The same table will be presented for participants randomised to no PES.

Table 2. Comparison of DSRS at day 90 between PES and no PES in pre-specified subgroups.

Outcome	N	All (N=xxx)	PES (N=xxx)	No PES (N=xxx)	aDIM/acOR/aOR (95% CI)†	Analysis
Day 90	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Day 90 change from baseline	xxx	x.x (x.x)	x.x (x.x)	x.x (x.x)	x.x (x.x, x.x)	aMLR
Subgroup, change ≥ -1 vs < -1						
≥ -1	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
< -1	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Subgroup, change ≥ -2 vs < -2						
≥ -2	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
< -2	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Subgroup, change ≥ -3.5 vs < -3.5						
≥ -3.5	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
< -3.5	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
Subgroup, absolute ≥ 4 vs < 4						
≥ 4	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR
< 4	xxx	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x)	xx.x (xx.x, xx.x)	aMLR

Figure 1. Subgroup analysis of the primary outcome.

Shown are the difference in means for swallowing impairment assessed using the dysphagia severity rating scale at day 90 in pre-specified subgroups. The widths of the confidence intervals have not been adjusted for multiplicity and should not be used in place of hypothesis testing. FOIS scores range from 1 to 7, with 0 indicating death and lower scores indicating worse swallowing impairment. DSRS scores range from 0 to 12, with 13 indicating death and higher scores indicating worse swallowing impairment.

- Change day 90-0: ≥ -1 , < -1
- Change day 90-0: ≥ -2 , < -2
- Change day 90-0: ≥ -3.5 , < -3.5 ,
- Absolute day 90: < 4 , ≥ 4
- Change day 90-0: mean difference

FOREST PLOT WITH INTERACTION TERMS