#### Correlation


Like covariance, but removes scale.

The correlation coefficient between X and Y is defined by

$$Cor(X, Y) = \rho = \frac{Cov(X, Y)}{\sigma_X \sigma_Y}.$$

#### Properties:

- **1.**  $\rho$  is the covariance of the standardized versions of X and Y.
- **2.**  $\rho$  is dimensionless (it's a ratio).
- **3.**  $-1 \le \rho \le 1$ .  $\rho = 1$  if and only if Y = aX + b with a > 0 and  $\rho = -1$  if and only if Y = aX + b with a < 0.



- Covariance formula is a statistical formula which is used to assess the relationship between two variables.
- In simple words, **covariance** is one of the statistical measurement to know the relationship of the variance between the two variables.
- The covariance indicates how two variables are related and also helps to know whether the two variables vary together or change together.
- The covariance is denoted as Cov(X,Y)

$$Cov(x,y) = \frac{\sum (x_i - \overline{x})(y_i - \overline{y})}{N}$$

$$Cov(x,y) = \frac{\sum (x_i - \overline{x})(y_i - \overline{y})}{N-1}$$

<u>tormula</u>

ξg: >(4ea1)	(Kg)
Age	Weight
20	<b>#</b> 5
18	63
15	45
14	40
25	78

Eq: (Year)	(k	(૧)		
(x) Age	(K Weight	(7)	X1	y1
18 20	63		X↓	71
15	45			
14	41	J X↑	71	(
25	78	J X↑ X↓	71	

Etonomic Gmwth (1/1)	NEFTY	SD Ind x(X)
2.1	8	
2.5	12	
4.0	14	
3.6	10	

Example-2

1 = 3.0 Dy - Z

#### (ov(x,4) = 5(x-7) (y-9 Covariance y-4 X- x 11 31 2.1 -0.6 11 3.1 12 2.5 0.9 11 3-1 14 4.0 11 3.1 10 3.6

### Solution

$$= \frac{3 - 0.6 + 2.7 + 0.5}{3}$$

$$(6v(x, y) = \frac{4.6}{3} = 1.533 \rightarrow +ve^{2}$$

$$(ov(x,y) = (-1)(-3) + (-0.6) \times (1) + (0.9) \times (3) + (0.5)(-1)$$

$$+ (0.5)(-1)$$

$$4 -1$$

## Solution

$$\int (x,y) = \frac{Cov(x,y) = 1.533}{T_{x}^{x}T_{y}} = \frac{(0.8981)(2.58)}{(2.58)}$$

Day	ABC Returns	XYZ Returns
1	1.1%	3.0%
2	1.7%	4.2%
3	2.1%	4.9%
4	1.4%	4.1%
5	0.2%	2.5%

• Daily Return for Two Stocks are given below

Find the covariance between the two stocks (the data is the sample data)

## **Solution-1**

### **Sample Covariance formula**

$$Cov(x,y) = \frac{\sum (x_i - \overline{x})(y_i - \overline{y})}{N-1}$$

- For ABC, the mean would be (1.1 + 1.7 + 2.1 + 1.4 + 0.2) / 5 = 1.30.
- For XYZ, the mean would be (3 + 4.2 + 4.9 + 4.1 + 2.5) / 5 = 3.74.

```
Covariance = [(1.1 - 1.30) \times (3 - 3.74)] + [(1.7 - 1.30) \times (4.2 - 3.74)] + [(2.1 - 1.30) \times (4.9 - 3.74)] + .../(5-1)
```

- = ([0.148] + [0.184] + [0.928] + [0.036] + [1.364]) / (5-1)
- $\bullet$  = 2.66 / (5 1)
- $\bullet$  = 0.665

## **Correlation**

 Definition The correlation coefficient of X and Y, denoted by Corr(X, Y), ρX,Y, or just ρ, is defined by

• It represents a "scaled" covariance – correlation ranges between -1 and 1.

$$r = rac{\sum \left(x_i - ar{x}
ight)\left(y_i - ar{y}
ight)}{\sqrt{\sum \left(x_i - ar{x}
ight)^2 \sum \left(y_i - ar{y}
ight)^2}}$$

r = correlation coefficient

 $oldsymbol{x}_i$  = values of the x-variable in a sample

 $\bar{x}$  = mean of the values of the x-variable

 $y_i$  = values of the y-variable in a sample


 $ar{m{y}}_{-}$  = mean of the values of the y-variable

When two random variables X and Y are not independent, it is frequently of interest to assess how strongly they are related to one another. The covariance between two rv's X and Y is


$$Cov(X,Y)$$
=  $E[(X - \mu(X))(Y - \mu(Y))]$ 

$$= \begin{cases} \sum_{x} \sum_{y} (x - \mu_{X})(y - \mu_{Y})p(x, y) & X, \text{ Y discrete} \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x - \mu_{X})(y - \mu_{Y})f(x, y) \, dx \, dy & X, \text{ Y continuous} \end{cases}$$

- If both variables tend to deviate in the same direction (both go above their means or below their means at the same time), then the covariance will be positive.
- If the opposite is true, the covariance will be negative.
- X and Y are not strongly related, the covariance will be near 0




## 3 types of "co-varying":




(a) positive covariance;

# 3 types of "co-varying":



(b) negative covariance;

## 3 types of "co-varying":



(c) covariance near zero

- An insurance agency services customers who have both a homeowner's policy and an automobile policy. For each type of policy, a deductible amount must be specified.
- For an automobile policy, the choices are \$100 and \$250, whereas for a homeowner's policy, the choices are \$0, \$100, and \$200. Suppose an individual "Rohit" is selected at random from the agency's files.
- Let X = his deductible amount on the auto policy and Y = his deductible amount on the homeowner's policy.
- Suppose the joint pmf is given by the insurance company the accompanying joint probability table:

p(x, y)		0	y 100	200
$\overline{x}$	100	.20	.10	.20
	250	.05	.15	.30

• What is the covariance between X and Y?

### **Correlation**

Definition The correlation coefficient of X and Y, denoted by Corr(X, Y),  $\rho X, Y$ , or just  $\rho$ , is defined by

$$\rho_{X, Y} = \frac{\operatorname{Cov}(X, Y)}{\sigma_X \cdot \sigma_Y}$$

It represents a "scaled" covariance – correlation ranges between -1 and 1.