



# Probability and Statistics

## *Lecture: Conditional probability*

---

Alice and Bob toss two fair coins separately. Denote the event that Alice gets a  $H$  by  $E_A$  and the event that Bob gets a  $H$  by  $E_B$ .

We know that  $P(E_A) = P(E_B) = 1/2$ .

## Example

## Example

Alice and Bob toss two fair coins separately. Denote the event that Alice gets a  $H$  by  $E_A$  and the event that Bob gets a  $H$  by  $E_B$ .

We know that  $P(E_A) = P(E_B) = 1/2$ .

- $P(E_A \cup E_B) = P(E_A) + P(E_B) = 1$ .
- So it's certain that at least one of them will get a head.

## Example

Alice and Bob toss two fair coins separately. Denote the event that Alice gets a  $H$  by  $E_A$  and the event that Bob gets a  $H$  by  $E_B$ .

We know that  $P(E_A) = P(E_B) = 1/2$ .

- $P(E_A \cup E_B) = P(E_A) + P(E_B) = 1$ .
- So it's certain that at least one of them will get a head.
- What's wrong with this argument?

## Example

Alice and Bob toss two fair coins separately. Denote the event that Alice gets a  $H$  by  $E_A$  and the event that Bob gets a  $H$  by  $E_B$ .

We know that  $P(E_A) = P(E_B) = 1/2$ .

- $P(E_A \cup E_B) = P(E_A) + P(E_B) = 1$ . So its certain that at least one of them will get a head.
- What's wrong with this argument?
- But both could get tails!

## Example

Alice and Bob toss two fair coins separately. Denote the event that Alice gets a  $H$  by  $E_A$  and the event that Bob gets a  $H$  by  $E_B$ .

We know that  $P(E_A) = P(E_B) = 1/2$ .

- $P(E_A \cup E_B) = P(E_A) + P(E_B) = 1$ . So it's certain that at least one of them will get a head.
- What's wrong with this argument?
- But both could get tails!
- Are  $E_A$  and  $E_B$  disjoint?

## Example

Alice and Bob toss two fair coins separately. Denote the event that Alice gets a  $H$  by  $E_A$  and the event that Bob gets a  $H$  by  $E_B$ .

We know that  $P(E_A) = P(E_B) = 1/2$ .

- $P(E_A \cup E_B) = P(E_A) + P(E_B) = 1$ . So its certain that at least one of them will get a head.
- Whats wrong with this argument?
- But both could get tails!
- Are  $E_A$  and  $E_B$  disjoint?
- Not quite.
- The sample space you are considering is:  $HT, TH, TT, HH$ .  
So  $E_A = \{HT, HH\}$  and  $E_B = \{TH, HH\}$ .

## Example

Alice and Bob toss two fair coins separately. Denote the event that Alice gets a  $H$  by  $E_A$  and the event that Bob gets a  $H$  by  $E_B$ .

We know that  $P(E_A) = P(E_B) = 1/2$ .

- $P(E_A \cup E_B) = P(E_A) + P(E_B) = 1$ . So it's certain that at least one of them will get a head.
- What's wrong with this argument?
- But both could get tails!
- Are  $E_A$  and  $E_B$  disjoint?
- Not quite.
- The sample space you are considering is:  $HT, TH, TT, HH$ .  
So  $E_A = \{HT, HH\}$  and  $E_B = \{TH, HH\}$ .
- So the union is really  
$$P(E_A \cup E_B) = P(E_A) + P(E_B) - P(E_A \cap E_B) = 3/4.$$



## Example

Alice and Bob toss two fair coins separately. Denote the event that Alice gets a  $H$  by  $E_A$  and the event that Bob gets a  $H$  by  $E_B$ .

We know that  $P(E_A) = P(E_B) = 1/2$ .

- $P(E_A \cup E_B) = P(E_A) + P(E_B) = 1$ . So its certain that at least one of them will get a head.
- What's wrong with this argument? But both could get tails!
- Are  $E_A$  and  $E_B$  disjoint? Not quite. The sample space you are considering is:  $HT, TH, TT, HH$ .

So  $E_A = \{HT, HH\}$  and  $E_B = \{TH, HH\}$ .

- So the union is really  
 $P(E_A \cup E_B) = P(E_A) + P(E_B) - P(E_A \cap E_B) = 3/4$ .
- May be what you are really thinking about is independent and not disjoint events.

# Partial information

- So far, we have assumed we know nothing about the outcome of our experiment, except for the information encoded in the probability law.
- Sometimes, however, we have **partial information** that may affect the likelihood of a given event.
  - The experiment involves rolling a die. You are told that the number is odd.
  - The experiment involves the weather tomorrow. You know that the weather today is rainy.
  - The experiment involves the presence or absence of a disease. A blood test comes back positive.
- In each case, knowing about some event  $B$  (e.g., “it is raining today”) changes our beliefs about event  $A$  (“Will it rain tomorrow?”).
- We want to **update** our probability law to incorporate this new knowledge.

# Conditional Probability

## Original problem:

- What is the probability of some event  $A$ .  
e.g. What is the probability that we roll a number less than 4?
- In other words, what is  $P(A)$ ?
- This is given by our probability law.

# Conditional Probability

## Original problem:

- What is the probability of some event  $A$ .  
e.g., What is the probability that we roll a number less than 4?
- In other words, what is  $P(A)$ ?
- This is given by our probability law.

## New problem:

- Assuming event  $B$  (equivalently given event  $B$ ), what is the probability of event  $A$ ?  
e.g., Given that the number rolled is an odd number, what is the probability that it is less than 4?
- We call this the **conditional distribution** of  $A$  given  $B$ .
- We write this as  $P(A|B)$
- Read  $|$  as “given” or “conditioned on the fact that”.
- Our conditional probability is still describing “the probability of something”, so we expect it to behave like a probability distribution.

# Conditional Probability

- Consider rolling a fair 6-sided die (uniform, discrete probability distribution).
- Let  $A$  be the event “outcome is equal to 1”.
  - What is  $P(A)$ ?
- Let's now assume that the number rolled is an odd number.
  - What is the set,  $B$ , that we are conditioning on?
- What do you think  $P(A|B)$  should be?

# Conditional Probability

- You are conditioning on event  $B$ . So your new sample space is  $\{1,3,5\}$  instead of  $\{1,2,3,4,5,6\}$ .

# Conditional Probability

- You are conditioning on event  $B$ . So your new sample space is  $\{1,3,5\}$  instead of  $\{1,2,3,4,5,6\}$ .
- Each of these are equally likely. So  $P(A|B) = 1/3$ .

# Conditional Probability

- You are conditioning on event  $B$ . So your new sample space is  $\{1,3,5\}$  instead of  $\{1,2,3,4,5,6\}$ .
- Each of these are equally likely. So  $P(A|B) = 1/3$ .

Formally, if all outcomes are equally likely, we have

$$P(A|B) = \frac{\# \text{ elements in } A \cap B}{\# \text{ elements in } B}$$



# Conditional Probability

- You are conditioning on event  $B$ . So your new sample space is  $\{1,3,5\}$  instead of  $\{1,2,3,4,5,6\}$ .
- Each of these are equally likely. So  $P(A|B) = 1/3$ .

Formally, if all outcomes are equally likely, we have

$$P(A|B) = \frac{\# \text{ elements in } A \cap B}{\# \text{ elements in } B}$$

More generally,  $P(A|B) = \frac{P(A \cap B)}{P(B)}$ .

# Conditional Probability

- You are conditioning on event  $B$ . So your new sample space is  $\{1,3,5\}$  instead of  $\{1,2,3,4,5,6\}$ .
- Each of these are equally likely. So  $P(A|B) = 1/3$ .

Formally, if all outcomes are equally likely, we have

$$P(A|B) = \frac{\# \text{ elements in } A \cap B}{\# \text{ elements in } B}$$

More generally,  $P(A|B) = \frac{P(A \cap B)}{P(B)}$ .

A conditional probability is only defined if  $P(B) > 0$ .

# Conditional Probability Axioms

- **Nonnegativity** – Check.
- **Normalization** –
- **Additivity** –

# Conditional Probability Axioms


- **Nonnegativity** – Check.
- **Normalization** – Your new universe is now  $B$ , and we know that  $P(B|B) = 1$ .
- **Additivity** –

# Conditional Probability Axioms


- **Nonnegativity** – Check.
- **Normalization** – Your new universe is now  $B$ , and we know that  $P(B|B) = 1$ .
- **Additivity** –  $P(A_1 \cup A_2|B) = P(A_1|B) + P(A_2|B)$  for two disjoint sets,  $A_1$  and  $A_2$ .

# Conditional Probability Axioms

- **Nonnegativity** – Check.
- **Normalization** – Your new universe is now  $B$ , and we know that  $P(B|B) = 1$ .
- **Additivity** –  $P(A_1 \cup A_2|B) = P(A_1|B) + P(A_2|B)$  for two disjoint sets,  $A_1$  and  $A_2$ .




Conditioning on  $B$   $\rightarrow$



# Conditional Probability Axioms

- **Nonnegativity** – Check.
- **Normalization** – Your new universe is now  $B$ , and we know that  $P(B|B) = 1$ .
- **Additivity** –  $P(A_1 \cup A_2|B) = P(A_1|B) + P(A_2|B)$  for two disjoint sets,  $A_1$  and  $A_2$ .



Using additivity,  $P((A_1 \cup A_2) \cap B) = P(A_1 \cap B) + P(A_2 \cap B)$ , so

$$P(A_1 \cup A_2|B) = \frac{P(A_1 \cap B) + P(A_2 \cap B)}{P(B)} = P(A_1|B) + P(A_2|B)$$

## Properties of conditional probability

If  $P(B) > 0$ ,

If  $A_i$  for  $i \in \{1, \dots, n\}$  are all pairwise disjoint, then

$$P\left(\bigcup_{i=1}^n A_i \mid B\right) = \sum_{i=1}^n P(A_i \mid B)$$



# Properties of conditional probability

If  $P(B) > 0$ ,

If  $A_i$  for  $i \in \{1, \dots, n\}$  are all pairwise disjoint, then

$$P\left(\bigcup_{i=1}^n A_i \mid B\right) = \sum_{i=1}^n P(A_i \mid B)$$

If  $A_1 \subseteq A_2$ , then  $P(A_1 \mid B) \leq P(A_2 \mid B)$ .

# Properties of conditional probability

If  $P(B) > 0$ ,

If  $A_i$  for  $i \in \{1, \dots, n\}$  are all pairwise disjoint, then

$$P\left(\bigcup_{i=1}^n A_i | B\right) = \sum_{i=1}^n P(A_i | B)$$

If  $A_1 \subseteq A_2$ , then  $P(A_1 | B) \leq P(A_2 | B)$ .

$$P(A_1 \cup A_2 | B) = P(A_1 | B) + P(A_2 | B) - P(A_1 \cap A_2 | B)$$

## Properties of conditional probability

If  $P(B) > 0$ ,

If  $A_i$  for  $i \in \{1, \dots, n\}$  are all pairwise disjoint, then

$$P\left(\bigcup_{i=1}^n A_i \mid B\right) = \sum_{i=1}^n P(A_i \mid B)$$

If  $A_1 \subseteq A_2$ , then  $P(A_1 \mid B) \leq P(A_2 \mid B)$ .

$$P(A_1 \cup A_2 \mid B) = P(A_1 \mid B) + P(A_2 \mid B) - P(A_1 \cap A_2 \mid B)$$

$$\text{Union bound: } P(A_1 \cup A_2 \mid B) \leq P(A_1 \mid B) + P(A_2 \mid B)$$

# Properties of conditional probability

If  $P(B) > 0$ ,

If  $A_i$  for  $i \in \{1, \dots, n\}$  are all pairwise disjoint, then

$$P\left(\bigcup_{i=1}^n A_i \mid B\right) = \sum_{i=1}^n P(A_i \mid B)$$

If  $A_1 \subseteq A_2$ , then  $P(A_1 \mid B) \leq P(A_2 \mid B)$ .

$$P(A_1 \cup A_2 \mid B) = P(A_1 \mid B) + P(A_2 \mid B) - P(A_1 \cap A_2 \mid B)$$

Union bound:  $P(A_1 \cup A_2 \mid B) \leq P(A_1 \mid B) + P(A_2 \mid B)$

$$P\left(\bigcup_{i=1}^n A_i \mid B\right) \leq \sum_{i=1}^n P(A_i \mid B)$$

## Example: Coin toss

Consider the experiment of tossing a fair coin three times. What is the probability of getting alternating heads and tails conditioned on the event that your first toss gives a head?

- Notation: Let  $A := \{\text{Tosses yield alternating tails and heads}\}$  and  $B := \{\text{The first toss is a head}\}$ .
- We want  $P(A|B)$ .
- What if you wanted to follow the formula?

## Example: Coin toss

Consider the experiment of tossing a fair coin three times. What is the probability of getting alternating heads and tails conditioned on the event that your first toss gives a head?

- Notation: Let  $A := \{\text{Tosses yield alternating tails and heads}\}$  and  $B := \{\text{The first toss is a head}\}$ .
- We want  $P(A|B)$ .
- What if you wanted to follow the formula?
  - Sample space for three coin tosses  
 $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$ .

## Example: Coin toss

Consider the experiment of tossing a fair coin three times. What is the probability of getting alternating heads and tails conditioned on the event that your first toss gives a head?

- Notation: Let  $A := \{\text{Tosses yield alternating tails and heads}\}$  and  $B := \{\text{The first toss is a head}\}$ .
- We want  $P(A|B)$ .
- What if you wanted to follow the formula?
  - Sample space for three coin tosses  
 $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$ .
  - $A = \{HTH, THT\}$  and  $A \cap B = \{HTH\}$ .

## Example: Coin toss

Consider the experiment of tossing a fair coin three times. What is the probability of getting alternating heads and tails conditioned on the event that your first toss gives a head?

- Notation: Let  $A := \{\text{Tosses yield alternating tails and heads}\}$  and  $B := \{\text{The first toss is a head}\}$ .
- We want  $P(A|B)$ .
- What if you wanted to follow the formula?
  - Sample space for three coin tosses  
 $\{ HHH, HHT, HTH, HTT, THH, THT, TTH, TTT \}$ .
  - $A = \{ HTH, THT \}$  and  $A \cap B = \{ HTH \}$ .
  - We have  $P(B) = 4/8$  and  $P(A \cap B) = 1/8$  and so  $P(A|B) = 1/4$ .



## Example: Coin toss

Consider the experiment of tossing a fair coin three times. What is the probability of getting alternating heads and tails conditioned on the event that your first toss gives a head?

- Notation: Let  $A := \{\text{Tosses yield alternating tails and heads}\}$  and  $B := \{\text{The first toss is a head}\}$ .
- We want  $P(A|B)$ .
- What if you wanted to follow the formula?
  - Sample space for three coin tosses  
 $\{ HHH, HHT, HTH, HTT, THH, THT, TTH, TTT \}$ .
  - $A = \{ HTH, THT \}$  and  $A \cap B = \{ HTH \}$ .
  - We have  $P(B) = 4/8$  and  $P(A \cap B) = 1/8$  and so  $P(A|B) = 1/4$ .
- Your new world/sample space when you condition is  $B = \{ HHT, HTH, HTT, HHH \}$ . Each of these are equally likely. Out of these 1 event satisfies alternating heads and tails.

## Example: Coin toss

Consider the experiment of tossing a fair coin three times. What is the probability of getting alternating heads and tails conditioned on the event that your first toss gives a head?

- Notation: Let  $A := \{\text{Tosses yield alternating tails and heads}\}$  and  $B := \{\text{The first toss is a head}\}$ .
- We want  $P(A|B)$ .
- What if you wanted to follow the formula?
  - Sample space for three coin tosses  
 $\{ HHH, HHT, HTH, HTT, THH, THT, TTH, TTT \}$ .  
 $A = \{ HTH, THT \}$  and  $A \cap B = \{ HTH \}$ .
  - We have  $P(B) = 4/8$  and  $P(A \cap B) = 1/8$  and so  $P(A|B) = 1/4$ .
  - Your new world/sample space when you condition is  
 $B = \{ HHT, HTH, HTT, HHH \}$ . Each of these are equally likely. Out of these 1 event satisfies alternating heads and tails.
  - So,  $P(A|B) = 1/4$ .