

Probability

Measure of uncertainty of events

Axiomatic theory = Classical theory

Equally likely outcomes

Conditional Probability

?

Conditional Probability

$$P(A|B)$$

Conditional Probability

Conditional probability of A given B

$$P(A|B)$$



Given

?

Conditional Probability

Conditional probability of A given B

$$P(A|B)$$



Given

Outcome satisfies **B** \longrightarrow How likely it will satisfy **A**?

Conditional Probability

Conditional probability of A given B

$$P(A|B)$$



Given

Outcome satisfies **B** \longrightarrow How likely it will satisfy **A**?

Chance of occurrence of **A** \longrightarrow **B** has already occurred



Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = ?$$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

*No. of first innings → 76
50 runs → 31
100 runs → 16*

$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = ?$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

No. of first innings → 76

50 runs → 31

100 runs → 16

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = 52\%$$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

*No. of first innings → 76
50 runs → 31
100 runs → 16*

$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = 52\%$

$P(\text{scores 100 runs} \mid \text{scored 50 runs}) \neq P(\text{scores 100 runs})$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

*No. of first innings → 76
50 runs → 31
100 runs → 16*

$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = 52\%$

$P(\text{scores 100 runs} \mid \text{scored 50 runs}) \neq P(\text{scores 100 runs})$

$P(\text{scores 100 runs}) = 16/76 = 21\%$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

No. of first innings → 76

50 runs → 31

100 runs → 16

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = \mathbf{52\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) \neq P(\text{scores 100 runs})$$

$$P(\text{scores 100 runs}) = 16/76 = \mathbf{21\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31$$

A → Pujara scores 50 runs

B → Pujara scores 100 runs

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

No. of first innings → 76

50 runs → 31

100 runs → 16

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = \mathbf{52\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) \neq P(\text{scores 100 runs})$$

$$P(\text{scores 100 runs}) = 16/76 = \mathbf{21\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31$$

A → Pujara scores 50 runs

B → Pujara scores 100 runs

$$P(B|A) = \frac{16}{31}$$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

No. of first innings → 76

50 runs → 31

100 runs → 16

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = \mathbf{52\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) \neq P(\text{scores 100 runs})$$

$$P(\text{scores 100 runs}) = 16/76 = \mathbf{21\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31$$

A → Pujara scores 50 runs

B → Pujara scores 100 runs

$$P(B|A) = \frac{16}{31} \rightarrow n(B \cap A)$$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

No. of first innings → 76

50 runs → 31

100 runs → 16

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = \mathbf{52\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) \neq P(\text{scores 100 runs})$$

$$P(\text{scores 100 runs}) = 16/76 = \mathbf{21\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31$$

A → Pujara scores 50 runs

B → Pujara scores 100 runs

$$P(B|A) = \frac{16}{31} \rightarrow \frac{n(B \cap A)}{n(A)}$$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

No. of first innings → 76

50 runs → 31

100 runs → 16

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = \mathbf{52\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) \neq P(\text{scores 100 runs})$$

$$P(\text{scores 100 runs}) = 16/76 = \mathbf{21\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31$$

A → Pujara scores 50 runs

B → Pujara scores 100 runs

$$P(B|A) = \frac{16}{31} \rightarrow \frac{n(B \cap A)}{n(A)} = \frac{\frac{n(B \cap A)}{n(S)}}{\frac{n(A)}{n(S)}}$$

Cheteshwar Pujara → Scored 50 runs → Chance of scoring 100 runs?

No. of first innings → 76

50 runs → 31

100 runs → 16

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31 = \mathbf{52\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) \neq P(\text{scores 100 runs})$$

$$P(\text{scores 100 runs}) = 16/76 = \mathbf{21\%}$$

$$P(\text{scores 100 runs} \mid \text{scored 50 runs}) = 16/31$$

A → Pujara scores 50 runs

B → Pujara scores 100 runs


$$P(B|A) = \frac{16}{31} \rightarrow \frac{n(B \cap A)}{n(A)} = \frac{\frac{n(B \cap A)}{n(S)}}{\frac{n(A)}{n(S)}} = \frac{P(B \cap A)}{P(A)}$$

Conditional Probability

What is the Probability of
rolling a dice and it's
value is less than 4

$$P(B | A) = \frac{P(A \cap B)}{P(A)}$$

knowing that the value is
an odd number



Example-1: The % of adults who are men and alcoholics is 2.25 %. What is the probability of being an alcoholics, given being man ?



Bayes' theorem

LIKELIHOOD

The probability of "B" being True, given "A" is True

PRIOR

The probability "A" being True. This is the knowledge.

$$P(A|B) = \frac{P(B|A).P(A)}{P(B)}$$

POSTERIOR

The probability of "A" being True, given "B" is True

MARGINALIZATION

The probability "B" being True.

Example-2: What is the probability of two girls given at least one girl?

Example-3: If we randomly draw a blue ball.
What is the probability of being in 1st bucket ?

B1: 3Yellow, 3 Blue

B2: 3Yellow, 2 Blue



Example-4:

B1: 3 Golds

B2: 2 Golds, 1 Silver

B3: 1 Golds, 2 Silver

A person chosen a box at random and takes out a coin. If the coin is Gold. What is the probability that it was drawn from box 3?