

Cambridge IGCSE

Cambridge Assessment International Education

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/42

Paper 4 Theory (Extended)

October/November 2019

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This syllabus is regulated for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

Cambridge Assessment International Education

1 The Periodic Table is very useful to chemists.

Refer only to elements with atomic numbers 1 to 36 in the Periodic Table provided when answering **Question 1**.

a) Us	se information from the Periodic Table provided to identify one element which:	
(i)	has atoms with exactly 9 protons	[1]
(ii)	has atoms with 0 neutrons	[1]
(iii)	has atoms with exactly 23 electrons	[1]
(iv)	has atoms with an electronic structure of 2,8,6	[1]
(v)	forms ions with a charge of 3– containing 18 electrons	[1]
(vi)	forms ions with a charge of 2+ containing 10 electrons	[1]
(vii)	has a relative atomic mass that shows it has at least two isotopes	[1]
b) St	tate which metal in the first 36 elements:	
(i)	is the Group I element which reacts most vigorously with water	[1]
(ii)	reacts with air to form lime.	[1]
c) (i)	ne element in the first 36 elements is used as the fuel in a fuel cell. Name this element.	
		[1]
(ii)	Write the overall chemical equation for the reaction which occurs when the element (c)(i) reacts in a fuel cell.	it in
		[2]
	[Total:	12]

2


The	gases Ar, CO ₂ , N ₂ and O ₂ are in clean, dry air.
CO	, NO, NO ₂ and SO ₂ are gases commonly found in polluted air.
(a)	What percentage of clean, dry air is N ₂ ?
	Give your answer to the nearest whole number.
	% [1]
(b)	Name the process used to separate O ₂ from clean, dry air. [2]
(c)	State one major adverse effect of the pollutant SO ₂ .
	[1]
(d)	NO and NO ₂ are produced in car engines.
	Describe how oxides of nitrogen form in a car engine.
	[2]
(e)	Many cars have catalytic converters in their exhaust systems. In a catalytic converter, most of the CO and NO formed in a car engine is changed into less harmful products.
	Identify these products and state the metal catalyst used.
	products
	catalyst[3]
(f)	CO is formed from the incomplete combustion of fossil fuels such as methane.
	Write a chemical equation to show the incomplete combustion of methane.
	[2]

[Total: 18]

4

(g) The ${\rm CO_2}$ in air is part of the carbon cycle.

The scheme shows a simple representation of part of the carbon cycle.



	in green plants	
(i)	State the scientific terms for each of process A and process B .	
	A	
	В	 [2]
(ii)	Plants convert glucose into complex carbohydrates.	[4]
(11)		
	A unit of glucose can be represented as HO——OH.	
	Complete the diagram to show the complex carbohydrate formed from three units glucose. Show all of the atoms and all of the bonds in the linkages.	of
		[2]
(iii)	Complex carbohydrates break down to form simple sugars.	
(111)		
	State two ways that complex carbohydrates can be broken down into simple sugars.	
	1	
	2	
		[2]
(iv)	Name a suitable technique for separating and identifying the individual sugars form when complex carbohydrates are broken down.	ned
		[1]

3

	3
Ammo	onia is an important chemical.
(a) A	mmonia is manufactured by the Haber process. The reaction is reversible.
(i	What is the sign for a reversible reaction?
	[1]
(ii	State the essential conditions for the manufacture of ammonia by the Haber process starting from hydrogen and nitrogen. Include a chemical equation to show the reaction which occurs.
	[5]
(iii	Name one raw material which is a source of the hydrogen used in the Haber process.
	[1]
(b) A	mmonia is a base and reacts with sulfuric acid to form the salt, ammonium sulfate.
(i	What is meant by the term base?
	[1]
(ii	Name the industrial process used to manufacture sulfuric acid.
	[1]

(iii) Write a chemical equation for the reaction between ammonia and sulfuric acid.

(c)		en aqueous ammonia is added to aqueous iron(II) sulfate a green precipitate is seen. This en precipitate turns red-brown at the surface.
	(i)	Name the green precipitate.
		[1]
	(ii)	Suggest why the green precipitate turns red-brown at the surface.
		ro1
		[2]
((iii)	State what happens when an excess of aqueous ammonia is added to the green precipitate.
		[1]

(d) Ammonia reacts with oxygen as show	(d)	Ammonia	reacts w	vith oxygen	as shown
--	-----	---------	----------	-------------	----------

$$4NH_3(g) + 5O_2(g) \rightarrow 4NO(g) + 6H_2O(g)$$

(i) Calculate the volume of oxygen at room temperature and pressure, in dm³, that reacts with 4.80 dm³ of ammonia.

(ii) The chemical equation for the reaction can be represented as shown.

$$4 \text{ H} - \text{N} - \text{H} + 5 \text{ O} = \text{O} \rightarrow 4 \text{ N} = \text{O} + 6 \text{ H} - \text{O} - \text{H}$$
 \downarrow
 \downarrow

Use the bond energies in the table to calculate the energy change, in kJ/mol, which occurs when **one** mole of NH_3 reacts.

bond	N–H	O=O	N=O	O–H
bond energy in kJ/mol	391	498	587	464

Energy needed to break bonds.

.....kJ

Energy released when bonds are formed.

.....kJ

Energy change when one mole of NH₃ reacts.


[Total: 22]

[4]

4 M	any sub	stances	conduct	electricity	V
------------	---------	---------	---------	-------------	---

(a)	Idei	ntify all the particles responsible for the passage of electricity in:
	•	graphite
	•	magnesium ribbon
	•	molten copper(II) bromide

(b) A student used the following apparatus to electrolyse concentrated aqueous sodium chloride using inert electrodes.




(1)	Suggest the name of a metal which could be used as the inert electrodes.	
		[1]
(ii)	Name the gas formed at the positive electrode.	
		[1]
(iii)	Write an ionic half-equation for the reaction occurring at the negative electrode. Inclustate symbols.	ade
		[3]
(iv)	How, if at all, does the pH of the solution change during the electrolysis? Explain y answer.	our

[Total: 16]

(c) A student used the following electrochemical cell.

The reading on the voltmeter was +1.10 V.



(1)	Draw an arrow on the diagram to show the direction of electron flow.	[1]
(ii)	Suggest the change, if any, in the voltmeter reading if the zinc electrode was replaced van iron electrode. Explain your answer.	with
iii)	The zinc electrode was replaced with a silver electrode. The reading on the voltmeter v –0.46 V.	vas
	Suggest why the sign of the voltmeter reading became negative.	
		[1]

- **5** Methanol, CH₃OH, is a member of the homologous series of alcohols.
 - (a) Methanol can be made from methane in a two-step process.

step 1 Methane is reacted with chlorine gas to produce chloromethane, CH₃C*l*.

step 2 CH₃C*l* is reacted with sodium hydroxide to produce CH₃OH and one other product.

(i) What conditions are needed in step 1?

.....[1]

(ii) Write the chemical equation for the reaction which occurs in step 1.

______[1]


(iii) State the type of organic reaction occurring in **step 1**.

......[1]

(iv) Complete the chemical equation for step 2.

$$CH_3Cl + NaOH \rightarrow CH_3OH + \dots$$
 [1]

(b) Draw a dot-and-cross diagram to show the electron arrangement in a molecule of methanol. Show outer shell electrons only.



[2]

www.dynamicpapers.com

11

(c) Me	ethanol reacts with propanoic acid to form an ester with a molecular formula C ₄ H ₈ O ₂ .
(i)	Name the ester formed when methanol reacts with propanoic acid.
	[1]
(ii)	Name one other substance formed when methanol reacts with propanoic acid.
	[1]
(iii)	Draw the structure of an ester which is a structural isomer of the ester named in (c)(i). Show all of the atoms and all of the bonds.
	[3]
(iv)	State the conditions needed to form an ester from a carboxylic acid and an alcohol.
	[1]
	[Total: 12]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

The Periodic Table of Elements

	 	2	บ L	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	52	Xe	xenon	98	R	radon			
	=				6	ш	fluorine 19	17	Cl	chlorine 35.5	35	ă	bromine 80	53	П	iodine 127	85	¥	astatine -			
	5				8	0	oxygen 16	16	တ	sulfur 32	34	Se	selenium 79	52	<u>е</u>	tellurium 128	84	Po	polonium -	116		vermorium -
	>									hosphorus 31												
	≥									silicon p										114	Ρl	lerovium -
	=	•								luminium 27												
										Ö			zinc 65							112	ت ت	pernicium -
													copper 64									
													nickel 59			_						damstadtium roe
Group													cobalt 59									
		-]		Irogen 1									iron 56									
				hyc									nanganese 55			_						
													hromium mang 52 5						tungsten rher			
				>	ımber	ymbol	e nic mass															m seaborgium
				Key	atomic number	atomic symbo	name relative atomic mass					>	vanadium 51						tantalum 181			mn dubnium
						(0					22	F	titanium 48	40	Zr	zirconiun 91			hafnium 178	104	Ÿ	rutherfordium -
								1			21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89-103	actinoids	
	=				4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium
	_				3	:=	lithium 7	11	Na	sodium 23	19	¥	potassium 39	37	Rb	rubidium 85	55	Cs	caesium 133	87	Ļ	francium -

71 Lu	lutetium 175	103	۲	lawrencium	I
o2 Yb				_	
e9 Tm	thulium 169	101	Md	mendelevium	ı
88 Er	erbium 167	100	Fm	ferminm	1
67 Ho	holmium 165	66	Es	einsteinium	_
66 Dy	dysprosium 163	86	ŭ	californium	ı
65 Tb	terbium 159	97	BK	berkelium	_
64 G d	gadolinium 157	96	Cm	curium	_
63 Eu	europium 152	92	Am	americium	1
62 Sm	samarium 150	94	Pu	plutonium	_
61 Pm	promethium -	93	δ	neptunium	_
90 09	neodymium 144	92	\supset	uranium	238
59 P	praseodymium 141	91	Ра	protactinium	231
Se Ce	cerium 140	06	H	thorium	232
57 La	lanthanum 139	88	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is $24\,\mathrm{dm^3}$ at room temperature and pressure (r.t.p.).