

**QuickRx**

Apps Download From Play Store

Pediatrics Inj. Drug Dose

Drug Name	Formulation	Dose
Ceftriaxone (Traxon, Ceftron, Oricef)	250mg/2.5ml 500mg/5ml 1gm/10ml 2gm/20ml	50-100 mg/kg/day (wt × 1)ml –OD 20-30 ml APN/BS দিয়ে Total 30ml বানাতে হবে। I/V @যত ml তত μ d/min ১ ঘন্টার মধ্যে শেষ করতে হবে।
Ceftazidime (Tazid, Serozid, Cefazid)	250mg/5ml 500mg/10ml 1gm/10ml 2gm/20ml	100-150 mg/kg/day (wt × 1) ml -BD Coverage Gram +ve
Amikacin (Kacin, Amibac)	100mg/2ml 500mg/2ml	15mg/kg/day Wt × 0.15 ml -BD Coverage Gram –ve, Nephrotoxic Drug
Flucloxacillin (Fluclox, Flux, Phylophen)	250mg/2.5ml 500mg/5ml 1gm	25 mg/kg/dose wt × 0.25ml + 5 D/W -QDS
Amoxicillin (Moxin, Moxacil, Fimoxyl)	250mg, 500mg	50-100mg/kg/day- TDS wt × 0.5 ml- TDS
Amoxicillin + Clavunic Acid (Moxaclav, Clacido)	1.2gm/20ml	50-100mg/kg/day- TDS wt × 0.5 ml- TDS
Ampicillin (Pen-A, Ampirex, Ampexin)	250mg, 500 mg	Neonate: 50mg/kg/day-BD Children: 100mg/kg/day-BD
Gentamycin (Gentin, Intamycin, Invigen)	20mg/2ml (Child/Neonate) 80mg/2ml (Adult)	5-6mg/kg/day wt × 0.25 ml- BD wt × 0.15 ml- TDS
Cefotaxime (Maxcef, Cefotax, Taxim)	250mg/5ml 500mg/10ml 1gm/20ml	60mg/kg/dose (wt × 1) <7day's: BD >7day's: TDS
Cefuroxime (Cefotil, Sefur, Kilbac)	750mg/10ml, 1.5gm	Neonate: 40-100mg/kg/day-BD (wt × 0.4 ml) BD. Children: 200-240mg/kg/day-TDS (wt × 0.9 ml) TDS
Meropenem (Meropen, Spacebac, Ropenem)	250mg/5ml 500mg/10ml 1gm (Not Recommended 0-3 mnth)	20-30mg/kg/dose Child: (wt × 0.4 or 0.8ml) + 20 ml B/S – BD/TDS Total 30ml বানাতে হবে। I/V @যত ml তত μ d/min ১ ঘন্টার মধ্যে শেষ করতে হবে।
Vancomycin (Vancomin, Covan)	500mg/10ml 1gm/20ml	60mg/kg/day, Meningitis: 15mg/kg/dose Child: wt × 0.4ml + 20 ml DW– TDS Total 20ml বানাতে হবে। I/V @যত ml তত μ d/min ১ ঘন্টার মধ্যে শেষ করতে হবে।
Ciprofloxacin (Ciprocil, Cipro, Neofloxin)	200mg/100ml	15-20mg/kg/day- BD (wt × 5 ml) BD
Metronidazole (Filmet, Metro, Amodis)	500mg/100ml	15-20mg/kg/day- TDS (wt × 1.5 ml) TDS

Hydrocortisone (Cotson, Glucort)	100mg/2ml	5 mg/kg/dose- TDS (wt × 0.1 ml) – BD/TDS/QDS
Furosemide (Lasix, Fusid)	20mg/2ml	1-2mg/kg/dose (wt × 0.1 ml) – BD (8 AM & 4 PM)
Dexamethasone (Roxadex, Decason)	5mg/ml	0.15-0.4mg/kg/dose –TDS/QDS (Meningitis) (wt × 0.03 ml- TDS/QDS) >3mnth (wt × 0.08 ml- BD) <3mnth
Phenobarbitone (Barbit)	200mg/1ml+9ml D/W	5mg/kg/day BD 12-24hourly LD= wt× 1ml I/V stat MD= wt× 0.125 ml wt× 0.125 ml wt× 0.125 ml or wt÷ 8 ml I/V- BD SOS: LD÷2 ml I/V stat&sos
Fosphenytoin (Fosphen)	150mg/2ml+8ml N/S	LD= wt× 1ml I/V stat (20-30mg/kg/dose over 15min) MD= wt× 0.25 ml I/V- BD (3-4mg/kg/dose over 15min) ** Add Same Amount N/S then give Double Micro Drop/ min.**
Diazepam (Sedil)	10mg/2ml (Not Recommended 0-3 mnth) Cardio-respiratory Distress করে।	0.2-0.3 mg/kg IV, 0.5 mg/kg/dose P/R (Max 10mg) wt × 0.1 ml (+3ml D/W, N/S) (P/R)- Stat Through rubber tube
Ondansetron (Emistat, Anset, Zofra)	8mg/4ml	0.15mg/kg/dose (wt × 0.08 ml) –TDS
Pheniramine Meleate (Avil, Amarin)	45.5mg/2ml	3mg/kg/day (wt × 0.1 ml) I/M Stat/OD
Tiemonium Methylsulphate (Viset, Algin, Visceral)	5mg/2ml	(wt× 0.05 ml) I/V –TDS
Famotidine (Peptid)	10mg/1ml	1-2mg/kg/day (wt× 0.05 ml) IV BD
Calcium gluconate (Hypocalcemia/ater 3 unit blood transfusion)	500mg/5ml 1gm/10 ml	50-70mg/kg/day wt× 1 ml+Same Amount D/W iv very slowly over 15-20 min @ 8µd/min
Vitamin K1 (Konakion) (K-One MM, K-One)	2mg/0.2ml (Neonate) 2mg orally or 1mg IV/IM within 4hours of birth. 10mg/2ml (Adult)	1 amp P/O stat, Then D4,D28 or ½ amp I/V stat, or 1amp I/M stat. (After Birth/if PT>15 sec)
Omeprazole (Seclo, Omep)	20mg/5ml 40mg/10ml	0.6-0.7 mg/kg/day bd (wt× 0.2ml) IV OD/ OD
Aminophylline (Filin)	125mg/5ml	2.5mg/kg/dose LD= (wt× 0.2ml)+3-4cc D/W stat MD= (wt× 0.1ml)+3-4cc D/W BD Apnoea (20sec শ্বাস আটকে থাকলে) Bradicardia,Cyanosis
Dopamin	200mg/5ml	1-20 mcg/kg/min 0.5 ml (in neonate) in each 100 ml fluid Fluid According to age. Then give IV @6-10 µd/min
Adrenalin	1mg/ml	1ml dilute with 9ml D/W then again 1ml dilute with 9ml d/w then give 1ml/kg/day
Sodium bi carbonate (Sodi bi carb)	7.5%	1ml dilute with 9ml amount 10%DA then 1ml/kg/dose
Clindamycin (Clinacyn,Clindax)	300mg/2ml 600mg/4ml	20-40mg/kg/day- TDS/QDS (wt × 0.1 ml) I/V TDS.
Levofloxacin (Levoflox)	500mg/100 ml	10-20 mg/kg/dose-OD (wt × 4 ml) I/V OD. [I/V @যত ml তত µd/min]

Cefepime (Ceftipime, Megapime)	500 mg/5ml 1gm/10ml 2gm/20ml	100 mg/kg/day (wt×0.5ml) IV BD
Acyclovir (virux,xovir)	250mg/10ml	0.4 ml/Kg/Dose –TDS, 10mg/kg/dose- TDS Mixed with (wt×2ml) DW/NS complete within 1 hour
	250mg/50ml 500mg/100ml	(wt×2ml) IV TDS
Antivenom (Polyvalent Antivenom)	6.7ml/vial	10 vial dilute with (10×10 =100 ml D/W) then further dilute with 100 ml DA,DNS,NS Total 200ml, Then give IV over 1 hr. <i>IV @ 40-50 D/min</i>
Pralidoxime (Pralidox)	1000mg/1ml	20-50mg/kg/Dose wt × 0.02-0.03 ml TDS (Mixed with IV fluid & Given Drip)
Atropine	0.6 mg/1 ml	0.05 mg/kg LD:(wt×0.08ml) iv & repeated every 10-15 min to a max 2-3 mg till pupil full dilated (Atropinization) then dose gradually tapered & maintained at least 2-3 days. MD: 0.03- mg/kg
Pralidoxime (Pradox, Pralidot)	(500mg/10ml)	30mg/kg/TDS (wt×0.5ml)+20ml B/S IV over 30 min
Benzathine penicilline	G-Benzathine penicillin:6,12 lac U	<30 kg-6 lac& >30kg:12 lac U IM single dose
Benzyl Penicillin	Penicillin-G Sodium 5,10 lac U	1 lac U/kg/day iv 6hrly for 10-14 d in Tetanus
Hepatitis B Vaccine (Engerix-B)	(10mcg/0.5ml) (20mcg/1ml)	0.5 ml IM anterolateral aspect of (Right thigh) within 12 hr.
Hepatitis B IG (Hepabig)	(100 IU/0.5ml) (200 IU/1ml)	0.5 ml IM anterolateral aspect of (left thigh) within 12 hr.
Immunoglobulin (IVIG)	0.5g(10 ml) 2.5g(50 ml) 5 gm	1ml=50mg plasma protein <i>Neonate</i> : 5 ml/kg - 3 day 1.7 ml/kg/hr over 3hr <i>Children</i> : same but if immunocompromised 3-5 ml/kg weekly; 0.4 ml/kg/hr over 12.5hr
Magnesium (MgSO4) (Nalepsin, Magsulf)	4gm/100ml	25-50mg/kg/dose IV TDS for 2-3 Dose MD: 30-60mg/kg/day Child: 100-200mg/kg/Dose QDS
Mannitol	20% -500ml (20gm in 100ml)	1-2 gm/kg/day (5-7.5 ml/kg/day) (wt×5ml) Complete Within ½ hr
Midazolam (Hypnofast, Dormitol)	5mg/1ml 15mg/3ml	0.2-0.3mg/kg (for sedation: 0.05-0.15 mg/kg) (wt×0.02ml)+ (diluted 3ml D/W) for any procedure.
TIG (Vaxitet-IG)	3000 IU/ml (I/M)	Prophylaxis:4 U/kg Tetanus:Infant-500 U Children:3000-6000 U
TT (TT Vax, Vaxitet)	(40 IU/0.5ml)	1 amp IM Stat
Procaine penicillin (Pronapen)	40 lac	25000-50000 U/kg bd im for 14 days

Diclofenac Sodium	75mg/3ml	3 mg/kg/day (wt×0.04ml) IM-TDS
Ketorolac (Rolac, Torax, Todol)	10mg/ml 30mg/ml 60mg/ml	< 10 kg: 10mg 1 amp (0.5-1mg/kg/dose) 10-20kg: 30mg 1/2 amp >20kg: 30mg 1amp IM/IV-BD/TDS (if given iv must dilute)
Rabies vaccine Rabipur	2.5 IU/ml	0.1 ml/dose in both arm I/D at Day-0,3,7 &28
Rabies IG Rabix IG		Single dose in around the lesion & also IM (it also use 3 rd degree lesion)
Pethidine Pehidine-R	100mg/2ml	(wt×2unit) in 100 unit insulin syringe IM- SOS/BD/TDS
Tranexamic acid Traxyl, Xamic	500mg/5ml	25 mg/kg- BD (wt×0.25 ml) IV BD
Clarithromycin Klaricid	500mg/10ml	15 mg/kg/day (wt×0.15 ml) + 10 ml B/S IV- BD
Teicoplanin Ticoplan	200mg/3ml 400mg/6ml	LD: 10 mg/kg/BD 3dose, MD: 6mg/kg/OD LD: (wt×0.15) + 10 ml B/S IV- BD..3dose MD: (wt×0.1) + 10 ml B/S IV- OD..7days
Cefepime Levoxin, Evo	500mg/100ml	(wt×2 ml) IV OD I/V @ ১০০ ml ত ১০০ μd/min
Human Albumin 25% Albutein 25%,	25gm/100ml	1gm/kg/day (wt×4 ml) IV stat I/V @ ১০০ ml ত ১০০ μd/min
Midazolam Milam	15gm/3ml	0.15-0.2gm/kg/day (wt×0.4 ml) IM stat & SOS
Linezolid Arilin	600mg/300ml	10mg /kg/dose (wt×5 ml) IV TDS
Levetiracetam Iracet	500mg/5ml	10mg /kg (wt×0.1 ml) IV BD

1 TSF per 8 kg weight / dose

Name of drugs	Dose	Strength/tsf	Frequency
Amoxicillin	20-50 mg/kg/day	125 mg	TDS
Amoxicillin + Clavulanic Acid	20-50 mg/kg/day	125+31.25mg	TDS
Cefpodoxime	10 mg/kg/day	40 mg	BD
Cefuroxime+ Clavulanic Acid	20-30 mg/kg/day	125 mg	BD
Cephadrine	50-100 mg/kg/day	125 mg	TDS
Chloroquine	10 mg/kg D1 & D2 then 5mg/kg in D3	80 mg	OD
Co-trimoxazole	10 mg/kg/day	40mg	BD
Flucloxacilline	50-100 mg/kg/day	125 mg	QDS
Paracetamol	15mg/kg/dose	120 mg	TDS/QDS
Phenobarbitone	5 mg/kg/day	20 mg	BD
Potassium (KT)	1-2 mg/kg/day ½ TSF+5 TSF water (If age <1 year) 1 TSF+10 TSF water (If age >1 year)	6.7 mmo/c	BD/TDS
Cotrim	10mg/kg/day	240mg	BD
Cefpodoxime	10mg/kg/day	40mg	BD
Mefenamic Acid	8mg/kg/dose	50 mg	BD/TDS
Cefpodoxime	10 mg/kg/day	40mg	BD
Lactulose		40mg	TDS
Cefadroxil	30-50 mg/kg/day	125mg	BD

½ tsf per 10kg weight/dose

Name of drugs	Dose	Strength/tsf	Frequenay
Azithromycin	10mg/kg/day (Enteric Fever 20mg/kg/day)	200 mg	OD
Carbamazepine	10 mg/kg/day	100 mg	BD
Cefixime	10mg/kg/day (Enteric Fever 20mg/kg/day)	100 mg	BD
Ciprofloxacin	15-30 mg/kg/day	250 mg	BD
Clarithromycin	15 mg/kg/day	125 mg	BD
Clindamycin	8-20 mg/kg/day	75 mg	TDS/QDS
Doxophylline	12 mg/kg/day	100 mg	BD
Iron	3-6 mg/kg/day	50 mg	OD
Metronidazole	15-30 mg/kg/day	200 mg	TDS
Naproxen	5 mg/kg/dose	125 mg	BD
Nitrofurantoin	3-7 mg/kg/day	25 mg	QDS
Piracetam	50 mg/kg/day	500 mg	BD/TDS
Sodium valproate	10-20 mg/kg/day	200 mg	BD
Multivitamin (Filwel Kids, Pogo) Naunehal (Hamdard)		1tsf/5ml	OD
Ambroxol HCL		15mg	BD/TDS
Salbutamol	0.4 mg/kg/day	15mg	TDS
Levosambutamo	0.1 mg/kg/day	1mg	TDS
Ketotifen		1mg	BD
Chlorpheniramine Maleate		2mg	BD/TDS
Desloratidine		2.5mg	OD
Rupatidine		5 mg	OD
Theophylline	10-20 mg/kg/day	120 mg	BD
Sodium valporate	10-20 mg/kg/day	200mg	BD
Sodium Picosulphate	0.25 mg/kg/day	5mg	OD
Butamirate Citrate		7.5mg	TDS
Phenoxymethyl Penicillin	50 mg/kg/day	250mg	QDS
Baclofen	0.25 mg/kg/dose	5mg	TDS
Desloratadine		2.5mg	OD
Rupatadine		5mg	OD

Ferrous fumarate		100mg	OD/BD
Iron Polymaltose Complex	3.33mg/kg/day	50mg	OD/BD
Ferrous sulphate		200mg	OD/BD
Deflazacort	0.6 mg/kg/day	6mg	BD

1 TSF per 10 kg weight / dose:

Name of drugs	Dose	Strength/tsf	Frequenay
Acyclovir	20 mg/kg/day	200 mg	QDS
Ceftibuten	9 mg/kg/day	90 mg	OD
Erythromycin	40-50 mg/kg/day	125 mg	QDS
Ibuprofen	15-20 mg/kg/day	100 mg	BD/TDS
Levofloxacin	10-20 mg/kg/day	125 mg	OD
Nitazoxanide	10mg/kg/dose	100 mg	BD
Penicillin	25-50 mg/kg/day	125 mg	QDS
Prednisolone	1-2 mg/kg/day	5 mg	BD/TDS
Zinc	0.5-1 mg/kg/day	10 mg	BD
Fluconazole	3-6 mg/kg/day	50 mg	OD
Fexofenadine HCL		30 mg	OD
Cetirizine hydrochloride		5mg	OD
Domperidone	0.2-0.4 mg/kg/dose	5 mg	TDS
Linezolid	10 mg/kg/dose	100mg	BD/TDS
Dicycloverine		10mg	TDS
Cephalexin		125mg	BD

Download **QuickRx** Apps from PlayStore

QuickRx is a complete Patient Management Guideline Software

If any Typing Mistake Please contact with Us via Whats APP/ Call: 01623-875729

Other's Syrup Dose

Name of drugs	Dose	Strength/tsf	Frequenay
Pyrantel Pamoate (Delentin, Melphin)	½ TSF/12 kg before meal 1 TSF/25 kg before meal	50mg	Single Dose
Levamisole (Etrax, Vermicom)	1 TSF/13 kg before meal 3 mg/kg single dose	40mg	Single Dose
Albendazole (Alben, Alphin)	1 TSF/13 kg >2years before meal	200mg	Single Dose
Ondansetron (Emistat, Anset)	0.15 mg/kg/dose 1 TSF/25 kg before meal	4 mg	TDS
Famoditine (Famotack, Servipec)	0.5 mg/kg/dose ½ TSF/20 kg before meal	40 mg	BD
Tiemonium Methylsulphate (Algin, Viset)	2 mg/kg/dose 1 TSF/ 5kg	10 mg	TDS
Folic Acid (B-9, Folic Kidz)	0.5 mg/kg/dose 1 TSF/ 5kg or 1ml/kg	2.5mg	OD

Download **QuickRx** Apps from PlayStore

QuickRx is a complete Patient Management Guideline Software

If any Typing Mistake Please contact with Us via Whats APP/ Call: 01623-875729

Drop per kg weight/ Dose

Name of drugs	Dose	Amount (drop) Per Kg	Strength (per ml)	Frequenay
Cefaclor (Drop)	20-40 mg/kg/day	2	100 mg	BD/TDS
Cefpodoxime (Drop)	10 mg/kg/day	4	20 mg	BD
Domperidone (Drop)	0.2-0.4 mg/kg/day	1	5 mg	TDS
Frusemide (Syp)	1-4 mg/kg/day	8 (.5 ml)	8 mg	OD
Iron Polymaltose Complex	1-3 mg/kg/day	1	50 mg	OD
Levetiracetam Syp	20 mg/kg/day	1.5 (0.1ml)	100 mg	BD
Multivitamin drop	10-15 drop	2		OD/BD
Vitamin B Complex		1		BD
Oxcarbazepine (Syp)	10-20 mg/kg/day	1.5 (0.1 ml)	60 mg	BD
Paracetamol (Drop)	10-15 mg/kg/day	3 (0.2ml)	80mg	TDS/QDS
Phenobarbitone (Syp)	5 mg/kg/day	1	4 mg	OD /BD
Ranitidine(Syp)	2-4 mg/kg/day	3 (0.2ml)	15 mg	BD
Cefixime	10 mg/kg/day (Enteric Fever 20mg/kg/day)	3 (0.2ml)	25mg	BD
Simethicone (Flacol, Pedigas)	<2years: 20mg/dose (0.3ml/5drops) >2years: 40mg/dose (0.6ml/10drops)	2	67mg	TDS/QDS
Amoxicillin	20-50 mg/kg/day	(wt× 2) +2	125mg/1.2ml	TDS
Normal Saline N/D (Norsol, Happysol)	0.9% Sodium Chloride	2	0.9gm/10 ml	TDS/QDS
Lomefloxacin E/D (Mexlo)		1 drop Each Eye	0.3%	TDS
Moxifloxacin E/D		1 drop Each Eye	0.5%	TDS/QDS
Chloramphenicol E/D		1 drop Each Eye	0.5%	TDS
Sodium Picosulphate (Syp)	0.25ml/kg/day	4		OD
Butamirate Citrate		2	5mg	TDS/QDS
Desloratadine		3	0.5mg	OD
Cetirizine		3	2.5mg	OD
Erythromycin	40-50 mg/kg/day	3 (0.3ml)	50mg/1.25ml	QDS
Cephadrine		2	125mg/1.2ml	QDS
Cefadroxil		3	125mg/1.25ml	BD
Cephalexin		2	125mg/1.2ml	BD
Ambroxol HCL		2	6mg	BD
Normal Saline Nasal Drop		2 Drops Each Nostril		TDS/QDS

❑ Syrup according to Age/ Wt:

Vit B complex :

Upto 1yr: ½ tsf BD

1-5yrs: 1 tsf BD

>5yrs: 1 tsf TDS

Cetirizine Dihydrochloride:

Upto 1yr: ½ tsf BD

1-5yrs: ½ tsf TDS

>5yrs: 1 tsf TDS

Levamisole:

1-4yrs : 1 tsf (1tab) OD

5-12yrs: 2 tsf (2tab) OD

>12yrs: 3 tsf (3tab) OD

Albendazole: National Guideline > 2years safe.

>1 <2yrs: 1 tsf OD

>2yrs: 2tsf OD

2yrs –adult: 400mg single dose

<2yrs : 200mg single dose

<1yrs :not recommended

Tab.estazol 400mg

Syp.estazol 200mg/tsf (10ml bottle)

Fexofenadine:

6mt-2yrs: ½ tsf BD

2yr-12yrs: 1 tsf BD

>12yrs: 2 tsf OD or 60mg Tab OD

Bromhexine HCl:

6mt to 1yr: 2-3ml BD.

1-5yrs: 1 tsf BD.

5-10yrs: 1 tsf TDS/QDS.

Iron Hydroxide polymaltose :

Paediatrics drop in Infants (5-10kg) : 8 to 15drops Daily.

Infants (5-10kg) : ½ to 1tsf daily.

Child 2-6yrs : 1tsf OD/BD

Child 6-12yrs: 1 to 2tsf OD/BD

Acyclovir:

Upto 2yr: ½ tsf 5times daily.

Child >2yr: 1 tsf 5times daily.

Herbisol/Pepnor:

Upto 6mts: ½ tsf BD/TDS.

6-12mts: 1tsf BD/TDS/QDS.

>1yr: 1-2 tsf TDS/QDS.

Tifal:

Upto 6mts: ½ tsf BD/TDS.

6-12mts: 1 tsf TDS.

>1yr: 1-2 tsf TDS/QDS.

Alkuli:

Upto 6mts: ½ tsf BD.

6-12mts: 1 tsf BD.

>1yr: 1-2 tsf BD/QDS.

Vitamin A :

Upto 6mt: 50,000IU/dose.

6mt-1yr: 1 Lac IU/dose.

>1yr: 2 Lac IU/dose.

It should be given orally on 2 successive days & a 3rd dose to be given at least 2 weeks later.

Vitamin C: Scurvy: 250-500mg/day, **Thalassemia:** 100mg Daily.

Upto 4yrs: 1 TSF BD

4-12yrs: 2 TSF BD

>12yrs: 1 tablet BD

Tab. Ceevit 250mg, 500mg, 1gm, Syp. 100mg/5ml, inj, 500mg/5ml

Vitamin D:

Child in simple Vit-D deficiency- 400IU daily.

Vitamin E:

Child: 1-10mg/kg/day.

Vitamin K:

Dose: 0.2mg/kg/dose

2mg orally or 1mg IV/IM within 4hours of birth. If given orally repeat the dose on day₄ & day₂₈.

Vitamin A+D+E (Cod liver oil):

Upto 1yr: ½ tsf OD

1-5yrs: 1 tsf OD

5-12yrs: 1½ tsf OD

Vitamin B+Zinc:

Upto 1yr: ½ tsf BD

1-5yrs: 1 tsf BD

>5yrs: 1 tsf TDS

Vit C+Vit D+Calcium:

Infants: As advised by physician.

Child 3-7yrs: ½ Tab daily.

Folic Acid:

1st day-1Tab Then 1/4th Tab OD. Another Dose: 0.2mg/kg/day PO (single dose)

Riboflavin: Child : 5-10mg 2-3times daily.

Thiamine: Child: 5-20mg daily.

Pyridoxine: Child: 20-25mg upto 3times daily

Zinc: 0.5-1mg/kg/day bd

<6mt: 10mg/day BD for 14days.

>6mt: 20mg/day BD for 14 days.

Artimether+Lumefantrine:Lumertam/Coartem;20/120 mg

5-14 kg: 1 tab

15-24 kg:2 tab

25-34 kg:3 tab

>35 kg:4 tab

1st dose:0hr ; 2nd dose: after 8 hr ; 3rd dose: after 16 hr ;next 3 doses:12 hrly.

***Contra indication : <5kg &1st trimester

Aspirin: Tab. Ecoprin 75,150,300 mg

30-60mg/kg/day 6hrly

In JRA-60-100mg/kg/day 6hrly

Baclofen: 1 tab=10mg

0.25mg/kg/dose–t.d.s [upto 40mg/day]

Captopril: 25mg & 50 mg tab

Infant: 0.15-0.3 mg/kg/day 8-12 hourly

Children: 0.1-0.5 mg/kg/day 8-12 hourly

Clonazepam: 1tab=0.5mg, 1mg, 2mg

0.01-0.03 mg/kg/day 2-3 divided Dose. (Max dose: 0.5 mg/kg/day)

Cyclophosphamide: Endoxan 50mg tab; 250, 500mg, 1gm inj

3mg/kg/day od-12 wks

[with alternate day prednisolone (.5mg/kg/day-1 yr)]

Deferiprone: Kelfer 250, 500 mg cap

25mg/kg TDS

Dicycloverine HCL:

6 month to 1 years: ½ tsf –t.d.s

1-10 years: 1 tsf –t.d.s

>10 years: 2 tsf –t.d.s

Digoxin: 0.04mg/kg/dose –o.d (1tab=0.25mg)

Diphenhydramine: 6.25—25mg 3-4 times daily

1-5 years: ½ tsf- t.d.s

5-12 years: 1 tsf- t.d.s

Enalapril: 5mg & 10 mg tab

0.08 mg/kg/day 12-24 hourly (up to max 1mg mg/kg/day)

Factor VIII: Aphanate 250, 500 IU; 10-15 IU/kg in mild hge/in prophylaxis

Hydrochlorthiazide: 25mg & 50 mg tab

<6 month: Upto 3 mg/kg/day 12 hourly

>6 month: 1-4 mg/kg/day 12 hourly

Imotil-2mg: >8 years 1 cap –o.d

Lactulose:

<1 years: ½ tsf –b.d

1-5 years: 1 tsf –b.d

5-12 years: 2 tsf –b.d

Mebendazole:

1 tsf b.d for 3 days

2yrs –adult: 100mg twice daily for 3 consecutive days.

<2yrs :not recommended

Tab. 100mg

Syp.ermox 100mg/tsf (30ml bottle)

Miconazole:

<2 years: ½ tsf –b.d
2-6 years: 1 tsf b.d
>6 years: 1tsf –q.d.s

Monera:

2-5 years: ½ tsf -b.d or t.d.s
6-12 years: 1 tsf –b.d or t.d.s

Montelukast:

6month-5yrs : 4mg –o.d
5-12 years: 5 mg –o.d
>12 years: 10 –o.d
Tab.montene 4mg.5mg.10mg.

Nifedipine: 0.2-0.5 mg/kg/dose (1 tab=10mg)

Nystatin drop: Nystat Oral Drop

Prophylaxis in neonates- 1ml Daily (5drops TDS)
Child- 1ml QDS for 3day (upto 12 years)

Pivmecillinam: 20-60mg/kg/day- t.d.s (1tab=200mg)

Falciparum: At the end of Rx single dose 0.75 mg/kg
Vivax: 0.3 mg/kg daily for 14 days.

Probio capsule: Cap.Acteria, Cap.GoodGut 1-2 cap t.d.s

Propanolol: 10 mg, 40 mg & 80 mg tab
0.5-1 mg/kg/day 6-8 hourly; May up to 2.5 mg/kg/day

Pyrental pamote : 6month – 12yrs :10mg/kg single dose
Adult: 500mg single dose
<6month :not recommended
Syp.delentin 50mg/ml .250mg/tsf(10ml bottle)
Tab.Melphin 125mg

Note : below 2yrs anti-helminthic drug should be use in the known case of helminthiasis.

Tiemonium Methylsulphate: 3-6 mg/kg/day TDS

½ TSF age <2years
1 TSF age >2years
Tab.viset 50mg, **Syp.**viset 10mg/tsf, **Inj.**viset 5mg/2ml

Tinidazole:Tinizol 1 gm tab

50 g/kg/day od for 3-5 days

Frusamide: 1-2mg/kg/dose Tab.Lasix 40mg

If 20 kg:1+1+0
If 10 kg: ½+½+0

Nifedipine: 0.25-0.50 mg/kg/day Tab. Nifin 10mg Tab. Nifin SR 20mg

If 20 kg: 1+1+0

If 10 kg: $\frac{1}{2} + \frac{1}{2} + 0$

Phenoxymethyl Penicillin: 50 mg/kg/day Tab. Penvik 250mg Tab. Penvik DS 500mg

If 20 kg: 1+1+1+1

If 10 kg: $\frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2}$

Rececadotril: 1.5 mg/kg/dose TDS. Cap. Receca 100mg, Sachet 10mg, 30mg

Infants less than 9 kg: one 10 mg sachet 3 times daily.

Infants from 9 kg to 13 kg: two 10 mg sachets 3 times daily.

Children from 13 kg to 27 kg: one 30mg sachet 3 times daily.

Children of more than 27 kg: two 30mg sachets 3 times daily.

Anti TB drugs

- **INH:** 10-15mg (10),
Rifampicin: 10-20mg (15),
Pyrazinamide: 30-40mg (35),
Ethambutol: 15-25mg (20),
Streptomycin: 12-18mg (15).

Note: Once daily at morning empty stomach (half an hour before breakfast)

A. Tab. Rimcure 3FDC

- Rifampicin-150,
- INH-75,
- Pyrazinamide-400mg.

B. Tab. Rimstar 4FDC

- Rifampicin-150mg,
- INH-75mg,
- Pyrazinamide-400mg,
- Ethambutol-275mg.

Ampicillin :

100mg /kg/day qds

cap. ampicillin 250mg, 500 mg

inj. ampicillin 250mg, 500mg vial

syp. ampicillin 125mg/tsf.

Amoxicillin:

PO: 20-50mg /kg/day tds, I/V, I/M: 50-100mg/kg/day tds (Septicaemia & Meningitis upto 100mg/kg/day)

cap. amoxicil 250mg, 500mg

Syp. amoxicil 125mg/tsf

Moxacil drop 100mg/ml

Inj. amoxicil 250mg, 500mg vial

Amoxicillin + clavulanic acid :

30mg/kg/day tds

>12yr: 375mg/625mg TDS

6-12yr : 2 tsf TDS

1-6yr: 1 tsf TDS

Syp.Moxaclav(125mg+31mg)

Tab.Moxaclav 375, 625, 1gm

Inj. Moxaclav 1.2gm

Ciprofloxacin :

15-30 mg/kg/day bd

tab.ciprocin 250mg,500mg,1gm

Inj.ciprocin 200mg/100ml bag

Syp.ciprocin 250mg/tsf

Levofloxacin:

PO: 5month-5yrs: 10mg/kg /dose bd

>5yrs: 10mg/kg/day daily

Tab.trevox 500mg.750mg

Syp.trevox 125/tsf

Inj.levoxin 500mg/100ml bag

Doxycycline:

PO: 5 mg/kg /day BD followed by 2.5mg/kg/day qds.

Cap.doxin 100mg

Flucloxacillin: 12.5 -25mg/kg/dose qds

cap.fluclox 250mg,500mg

Syp.fluclox 125mg/tsf

Inj.fluclox 250mg,500mg vial

Tetracycline :

25-50mg/kg/day qds

Cap.tetracyna 250mg,500mg

Erythromycin:

PO : 50mg/kg/day qds

Tab.eromycin 250mg,500mg

Syp.eromycin 125mg/tsf

Azithromycin :

10mg/kg/day once daily(enteric fever 20g/kg/day once daily)

Tab.zimax 250mg,500mg

Syp.zimax 200mg/tsf

Inj.zimax 500mg/bottle

Clarithromycin:

15mg/kg/day bd.

Tab.binoclar 250mg,500mg

Syp.binoclar 125 mg/tsf

Inj.klaricid 500mg

Amikacin:

Neonate: I/V : 10mg/kg/day bd

Child: I/V : 15-25mg/kg/day bd-tds

Inj.kacin 100mg/2ml, 200mg/2ml amp

Clindamycin:

Neonate: I/m,I/V : 10-15mg/kg/day bd

Child: 8-20 mg/kg/day TDS

Cap.qcin 150mg,300mg

Syp.clidax 75mg/tsf

Inj.qcin 300mg/2ml amp. 600mg/4ml amp

Cotrimoxazole:

PO: 8mg/kg/day bd

Tab.cotrim 480mg,960mg

Syp.cotrim 200mg/tsf

Cephadrine:

25-50mg/kg/day qds

Cap.sefrad 250mg,500mg

Syp.sefrad 125mg/tsf

Sefrad drop 100mg/tsf

Inj.sefrad 250mg,500mg,1gm vial

Cefodroxil:

PO: 30mg/kg/day bd

Cap.arocef 250mg,500mg

Syp.arocef 125mg/tsf

Cefaclor:

PO: 20-40mg/kg/day tds

Cap.oticlор 250mg,500mg

Syp.oticlор 125mg/tsf

Oticlor drop 100mg/ml

Cefixime:

PO: 10mg/kg/day bd (enteric fever 20mg/kg/day bd)

Cap.cef 3 200mg,400mg

Syp.cef 3 100mg/tsf

Ceftid paediatrics drop 25mg/ml

Cefdoxime:

PO: 10mg/kg/day divided bd

Tab.ximeprox 200mg

cap.ximeprox 100mg

syp.ximeprox 40mg/tsf

Ximeprox drop 20mg/ml

Chloroquine:

Malaria prophylaxis: 5mg/kg/wk PO (max dose 300mg/dose).

Uncomplicated malaria: 10mg/kg PO on D1, then 7.5mg/kg on D2-D3 + Primaquine on D4, 3Tab - stat (1Tab- 15mg);

Or 10mg/kg PO on D1 then 5mg/kg on D2-D4.

Tab. 250mg(150mg base), Syp. 80mg(50mg base)/5ml.

Quinine:

child: po:10mg/kg /dose tds for 7days.

In Short:3-9 kg :1/4 ; 10-19 kg: ½ ; 20-29 kg: 1 ;30-39 kg :1 ½ &>40 kg :2 tab tds

Adult: 600mg tds for 7 days.

I/V: 20mg/kg infuse over 4 hours then 10mg/kg over 4hours,continue every 8-12hourly followed by oral quinine.

Tab.jasoquine 300mg

Inj.jasoquine 300mg /5ml ampule

Sulfadoxine+pyrimethamine:

adult: over 14yrs: 2-3 tab single dose

Child : under 4 yrs ½ tab single dose

4-8 yrs 1 tab single dose

9-14 yrs 2 tab single dose

Tab .malacide

Primaquine:

adult: 15mg daily for 14-21 days.

Child : ½ of the adult dose.

Tab.jasoprim 15mg

Artemether + lumefantrine :

D1=4tab stat and 4 tab after 8hours

D2+D3= 4 tab 12 hourly

Tab.coartem

Ivermectin:

150 microgm/kg single dose and repeate after 14 day.

Tab.scabu 3mg ,6mg.

Fluconazole:

5mg/kg/day ---daily.

Cap.flugal 50mg.150mg.200mg.

Syp.flugal 50mg/tsf

Nitazoxanide:

8.5mg/kg/dose bd

<1yrs not recommended

1-4yrs –1tsf bd

4-11yrs—2tsf bd

Syp.zox 100mg/tsf

Tab.zox 500mg

Nalidixic acid:

55mg/kg/day tds

Tab.nalid 500mg.

Syp.nalid 300mg/tsf

Paracetamol:

15mg/kg/Dose tds or qds (In the case of *liver causes* given lowest dose)

Tb.ace 500mg

Tab.ace xr 665mg

Syp.ace 120mg/tsf

Ace drop 80mg/ml

Ace suppository 60mg.125mg.250mg500mg.

Inj.napa 1000mg/100ml bottle

Domperidon:

0.2-0.4mg/kg/dose –qds

Tab.omidon 10mg

Syp.omidon 5mg/tsf

Drop omidon 5mg/ml

Don a suppository 15mg.30mg

Ketoconazole:

5mg/kg/day----daily

Tab.ketoral 200mg.

Ondansetron:

I/V: each 10kg/0.5ml tds

Po: <15kg—1/2 tsf tds

>15kg---1 tsf tds

Inj.emistat 8mg/4ml amp

Syp.emistat 4mg/tsf

Tab.emistat 4mg,8mg

Frusemide:

1-3mg/kg/day divided bd

Tab.frusid 40mg

Inj.frusid 20mg/2ml

Spirinolactone:

3mg/kg/day divided tds

Tab.verospiron 25mg.

Cetirizine:

>6yrs: 5mg twice daily or 10mg once daily.

2-6yrs : 5mg daily

6month-23month: 2.5mg daily

Syp.alatrol 5mg/tsf

Tab.alatrol 10mg

Levocetirizine:

6yrs-11yrs : 2.5mg daily

6month-5yrs: 1.25mg once daily.

Tab.alcet 5mg

Syp.alcet 2.5mg/tsf

Loratidine:

>3yrs: >30kg :10mg/day

<30kg :5mg/day

<3yrs : 5mg daily

<2yrs :not recommended

Tab.loratin fast 10mg

Syp.loratin 5mg/tsf

Desloratidine:

6month-11month :1mg daily
2-5 yrs: 1.25mg once daily.
6-11mg: 2.5mg once daily
Syp.deslor 2.5mg/tsf
Tab.deslor 5mg

Mebhydrolin :

up to 2yrs :1-2 tab daily
2-5yrs:1-3 tqb daily
5-10yrs 2-4 tab daily
Tab.bexidal 50mg

Pheniramine :

1-3yrs: ½ tsf 2-3times daily
4-12yrs:1tsf 2-3 times daily
Syp.avil 15mg/tsf
Tab.avil 22.5mg
Tab.avil retard 75mg
Inj.avil 45.5/2ml

Promethazine:

2-5yrs :5-15mg daily 1 or 2 divided dose.
5-10yrs:10-25mg daily 1 or 2 divided dose.
Syp.Phenexerex 5mg/5ml
Inj.phenexerex 50mg/2ml.25mg/1ml.
Tab.phenexergan 10mg,25mg

Prednisolone :

1-2mg/kg/day divided bd-qds.
Tab.cortan 5mg,10mg,20mg
Syp.cortan 5mg/tsf

Dexamethasone :

0.2mg/kg/dose qds(iv,im,po)
Tab.oradexon 0.5mg
Inj.oradexon 5mg/1ml amp

Hydrocortisone :

I/V: 10-20mg/kg/day divided bd-qds
Shock: 50mg/kg/dose
Inj.anacort 100mg vial
Tab.cotson 10mg

Salbutamol:

0.4mg/kg/day divided tds-qds.

Tab.brodil 2mg.4mg

Syp.brodil 2mg/tsf

Levosalbutamol:

<6yrs :1/2 tsf tds

6-11yrs: 1mg tds

Tab.brodil levo 1mg.2mg

Syp.brodil levo 1mg/tsf

Aminophylline :

20mg/kg/day tds

Tab.aminophylline 100mg

Inj.aminophylline 125mg/5ml amp

Theophylline:

20mg/kg/day tds

Syp.contine 120mg/tsf

Tab.contine 200mg,300mg,400mg

Doxophylline:

6mg/kg/day divided bd

Syp.doxiva 100mg/tsf

Tab.doxiva 200mg,400mg

carbamazepine :

PO: 10mg/kg/day divided tds

tab.tegretol 200mg

syp.tegretol 100mg/tsf

Ambroxol HCL:**drop:**

0-6month: 7drop bd

6-12month:15drop bd

1-2 yrs: 20drop bd

Syp:

2-5yrs:1/2 tsf bd-tds

5-10yrs :1tsf bd-tds

>10yrs: 2tsf tds

Cap:

> 10yrs + 1 cap daily

Strength:

Cap.ambrox 75mg SR

Syp.ambrox 15mg/tsf

Drop.ambrox 6mg/ml

Ketotifen :

>2yrs : 1mg twice daily
<2yrs: ½ tsf twice daily
Syp.tofen 1mg/tsf
Tab.tofen 1mg

Diazepam:

PO: 0.5-1mg/kg/daily divided bd-tds
P/R: 0.5mg/kg/dose
I/V: 0.2-0.3mg/kg/dose bd-tds
Tab.sedil 5mg
Inj.sedil 10mg/2ml amp
Easium suppository 10mg/stick

Phenobarbitone:

I/V.PO: 15-20mg/kg/stat followed by 3-4mg/kg/day neonate (bd) & 5- 6mg/kg/day for child (bd)
For sedation :2mg/kg/dose
tab.barbit 30mg
inj.barbit 200mg/1ml amp(inj should be dissolved in 9ml distill water)

phenytoin:

PO: 15-20mg/kg stat followed by 10mg/kg/day neonate & 10mg/kg/day child (bd)
tab.diphedan 100mg
syp.diphedan 60mg/tsf

sodium valproate :

10mg/kg/day divided BD
tab. Valex CR 200mg
syp.valex 200mg/tsf

Midazolam:

0.15-0.2mg/kg/dose for any procedure.
Tab.dormicum 7.5mg
Inj.dormicum 5mg/5ml.15mg/3ml.

Indomethacin:

2.5mg/kg/day divided bd
Cap.indomet 25mg
Cap.indomet SR 75mg
Indomet supposit0ry 100mg/stick

Ibuprofen:

15mg/kg/day divided bd
Tab.inflam 200mg.400mg
Syp.inflam 100mg/tsf

Naproxen:

15mg/kg/day bd
Syp. Ticoflex 125mg/tsf
Tab.naprosyn 250mg.500mg

Ketorolac:

2-16yrs: 0.4-1mg/kg/dose BD/TDS
Tab.torax 10mg

Diclofenac:

2-3mg/kg/day BD/TDS
Tab.clofenac 25mg.50mg.100mg
Inj.clofenac 75mg/3ml amp
Clofenac supp 12.5mg.25mg.50mg

K+T:

3mmol/kg/day divided bd
Syp.electro k 10mmol/tsf (mixed with 1galss water)

Ranitidine:

PO: 7.5mg/kg/day bd
Neonate:PO.I/V: 1.5-2mg/kg/day bd
Child:PO.I/V: 1-5mg/kg/day tds-qds
Syp.gepin 75mg/tsf
Tab.gepin 150mg.300mg
Inj.gepin 50mg/2ml

Aminocaproic acid :

100-200mg/kg bolus followed by 100mg/kg/dose qds
Inj.caprolisin 1gm/5ml. 2gm/10ml amp.

Chlorpromazine:

PO: 0.3mg/kg/day tds
I/M: 0.1mg/kg/dose
Tab.largactil 25mg.50mg.100mg
Syp.largactil 25mg/tsf
Inj.largactil 50mg/2ml amp

Calcium gluconate :

PO: 200-400mg/kg/day tds-qds
I/V: 1-2mg/kg/dose
Tab.calcium gluconate 600mg
Inj.calcium gluconate 500mg/ 5ml.1gm/10ml amp

Piracetam :

50mg/kg/day tds

tab.neulep 800mg.

syp.neurolep 500mg/tsf

Mannitol :

0.5-1gm/kg bolus followed by 0.25-0.5/kg /dose 4-6hourly.

Inf. Manisol 20% IV 500ml

Tranexamic acid:

25mg/kg/dose bd-tds

Cap.frabex 500mg

Inj.frabex 250mg/5ml.500mg/5ml

Clarithromycin:

15mg/kg/day bd.

Tab.binoclr 250mg,500mg

Syp.binoclar 125 mg/tsf

Inj.klaricid 500mg

Famotidine:

0.5 mg/kg/dose bd. (Max Dose: 40mg/day)

Tab. Famotack 20 mg

Syp. Famotack 40mg/tsf

Hydroxyzine HCL:

1-5yr: ½ TSF BD

>5yr: 1 TSF TDS

Syp. Xyiril 10mg/5ml

Tab. Xyiril 25mg

Probiotics/ Acteria Sachet

1 Sachet mixed with ½ Cup Water..OD

Chlorpheniramine Malate

Upto 1yr: 1mg/2.5ml BD,

1-5yr: 2mg/2.5-5ml TDS,

>5yr: 2-4mg/5-10ml TDS.

Tab.2mg,

Syp. 2mg/5ml

Fluid Requirement:

Age:	Req: (ml/kg/day)
• Day 1	• 60ml
• Day 2	• 80ml
• Day 3	• 100ml
• Day 4	• 120ml
• Day 5	• 140ml
• Day 6 to 9m	• 150ml
• 9m- 1 yr	• 120-150ml
• 1-2 yr	• 100-120ml
• 2-4 yr	• 90-100ml
• 4-8 yr	• 70-90ml
• 8-12 yr	• 60-70ml

For Infant & Children: >9month

- ❖ For 1st 10kg body wt: 100ml/kg/day.
- ❖ For next 10kg body wt: add 50ml/kg/day.
- ❖ Everything over 20kg body wt: add 20ml/kg/day.
[max: 2400ml/day]

Example: Maintenance fluid requirement for 28kg child will be as follows-
 $(100 \times 10) \text{ml/day} + (50 \times 10) \text{ml/day} + (20 \times 8) \text{ml/day} = (1000 + 500 + 160) \text{ml/day} = 1660 \text{ml/day}$.

Fluid to be given:

Age:	Fluid:	Content:
• Day 1	If wt: >1kg 10% DA	10% DA: 10% glucose
	If wt: <1kg 5% DA	5% DA: 5% glucose
• Day 2- 2years	If wt: >1kg 10% DBS /Electrodex 10/ APN	APN: 10%DA + 0.225% NaCl
	If wt: <1kg Baby saline	B/S: 5%DA + 0.225% NaCl
• 2-5 Years	Baby Saline	5%DA + 0.225% NaCl
• >5 years	Libott's Junior	5% glucose + 0.45% Na
• >11years	Adult fluid: 5%DNS, 5% DA, H/S, N/S)	

1. Condition requiring 20% more fluid:

- Fever
- Phototherapy
- Burn or open wound

2. Condition requiring 20% less fluid: (CNS, CVS, Respiratory Cause)

- Sepsis
- PNA
- NEC
- SIADH
- Meningitis

**In Head Injury

- <25 kg- Baby saline
- >25 kg- N/S

Note: Glucose Causes Cerebral Edema in Head injury.

❑ Composition of ORS: (Dissolved in 500ml clean water)

Components:	Grams/500ml:
Sodium Chloride	1.30 gm
Potassium Chloride	0.75 gm
Tri sodium citrate dihydrate	1.45 gm
Glucose Anhydrous	6.75 gm

❑ Composition of Cholera Saline:

Components:	Grams/Litre	mmol/Litre
Na ⁺	3.0	134
Cl ⁻	3.8	98
K ⁺	0.5	13
Acetate	2.8	84

❑ Some Important formula:

→ 24 Hours Fluid Requirement:

- Previous day output + Insensible loss of Fluid

→ Blood Transfusion (Transfusion should be given over 4 hours)

- PCV: (wt in kg × 4 × Hb Deficit) {Hb Deficit= Required Hb (10gm/dl) – Patient Hb}
- Or Randomly: PCV= wt in kg × 10ml/kg
- FHB: (wt in kg × 6 × Hb Deficit) {Hb Deficit=Required Hb (10gm/dl) – Patient Hb}
- Or Randomly: FHB= wt in kg × 20ml/kg (Anti-coagulant 3cc/20ml blood)

→ Body surface area(BSA):

- Height in cm × Wt in Kg ÷ 3600

→ Estimated Glomerular Filtration rate(GFR):

- $K \times \text{height (cm)} \div \text{Serum creatinine (mg/dl)}$
[K= 0.43]

→ Spot Protein Creatinine Ratio. (If >2 suggest NS)

- Urine Protein (mg) ÷ Urine Creatine (mg)

→ **Serum Anion gap:**

- $\text{Na}^+ - (\text{Cl}^- + \text{HCO}_3^-)$
[Normal value: 6-12 mEq/L]

→ **Free water deficit:**

- Total body water $\times (\text{Na}^+ \text{ concentration } / 140) - 1$
[Total body water: $0.6 \times \text{wt(kg)}$]

→ **Expected bladder capacity:**

- <2yr: $[\text{wt in kg} \times 7] \text{ ml}$
- >2yr: $[\text{Age in yrs} + 2 \times 30] \text{ ml}$

→ **Plasma osmolarity:**

- $2(\text{Na}^+) + \text{BUN(mg/dl)} \div 2.8 + \text{glucose} \div 18$
[Normal : 280-285 mOsm/kg]

→ **Amount of fluid for drop:**

- Amount of fluid $\div (4 \times \text{hour})$
- For Maintain 24 hours: যত ml তত Drop/min (Ex: 1000ml @10d/min)

→ **Amount of fluid for micro drop:**

- Amount of fluid $\div \text{hour}$.

→ **Nebulization with Windel Plus (2.5mg+500mcg) 2.5ml**

- $\text{Wt} \times 0.1 \text{ ml TDS/QDS}$
<1year = 0.5ml+1.5ml N/S stat & QDS
1-4 years = 1ml+ 2ml N/S stat & QDS
>4 years = 1amp stat & QDS

→ **Nebulization with Budicort 0.5mg/2ml**

- Neonate(<2month)
 $\text{Wt} \times 0.5 \text{ ml} + 1 \text{ ml N/S stat \& BD/QDS}$

❏ Easy to count weight:

- ❖ **3mt-12mt:** Age in month $+ 9 \div 2$
- ❖ **1yr-10yrs:** Age in year $+ 4 \times 2$

Nebulition with

WindelPlus	: $\text{Wt} \times 0.1 \text{ ml}$
Budicort	: $\text{Wt} \times 0.5 \text{ ml}$
N/S	: 1-2 ml

Stat and 8 hourly

Or

Nebulition with

Sultolin Solution	: $\text{Wt} \times 0.03 \text{ ml}$
Iporex	: $\text{Wt} \times 0.1 \text{ ml}$
N/S	: 1-2 ml

Stat and 6 hourly

Note: Make Total 3ml fluid

❏ Formula Feeding

(Biomeal, Lactogen)

Upto 6 month : I

Upto 1 yrs : II

>1 yrs : III

In case of acute watery diarrhoea give lactogen

free milk (Gastro-fix, Baby Luck FL)

নিয়ম: ৬ চামচ পানির সাথে ১ চামচ দুধ (বাচ্চার প্রয়োজন অনুযায়ী)

❏ Burn Fluid Management

Parkland Formula: $4 \times \text{Percentage of burn} \times \text{Weight}$

Then give Half Amount 1st 8 Hour's


Then Give another Half amount next 16 Hour's

For baby/child (weight <30kg): Parkland Formula+ Maintenance fluid (Baby Saline)

For Baby/child: Baby saline

For Adult: H/S

** Randomly: রোগীর হাতের তালুর অংশ ১% ধরা হয়।



Download **QuickRx** Apps from PlayStore

QuickRx is a complete Patient Management Guideline Software

If any Typing Mistake Please contact with Us via Whats APP/ Call: 01623-875729

শিশুদের নিয়মিত টিকাদান সময়সূচি: **EPI Schedule in Bangladesh**

রোগের নাম	টিকার নাম	টিকার ডোজ	ডোজের সংখ্যা	ডোজের মধ্যে ন্যূনতম বিরতি	টিকা দেয়ার সঠিক সময়	টিকাদানের স্থান	টিকার প্রয়োগ পথ
যক্ষ্মা	বিসিজি	০.০৫ এম এল	১	-	জন্মের পর থেকে	বাম বাহুর উপরের অংশে	চামড়ার মধ্যে
ডিফথেরিয়া, হুপিংকাশি, ধনুষ্ঠংকার, হেপাটাইটিস-বি, হিমোফাইলাস ইনফ্লুয়েঞ্জা বি	পেন্টাভ্যালেন্ট টিকা (ডিপিটি, হেপাটাইটিস-বি, হিব)	০.৫ এম এল	৩	৪ সপ্তাহ	৬ সপ্তাহ ১০ সপ্তাহ ১৪ সপ্তাহ	বাম উরুর মধ্যভাগের বহিরাংশে	মাংসপেশী
নিউমোকোকাল নিউমোনিয়া	পিসিভি টিকা	০.৫ এম এল	৩	৪ সপ্তাহ	৬ সপ্তাহ ১০ সপ্তাহ ১৪ সপ্তাহ	ডান উরুর মধ্যভাগের বহিরাংশে	মাংসপেশী
পোলিওমাইলাইটিস	বিওপিভি	২ ফোঁটা	৩	৪ সপ্তাহ	৬ সপ্তাহ ১০ সপ্তাহ ১৪ সপ্তাহ	মুখে	মুখে
	আইপিভি (ফ্রাকশনাল)	০.১ এম এল	২	৮ সপ্তাহ	৬ সপ্তাহ ১৪ সপ্তাহ	ডান বাহুর উপরের অংশে	চামড়ার মধ্যে
হাম ও রুবেলা	এমআর টিকা	০.৫ এম এল	২	-	৯ মাস ও ১৫ মাস বয়স পূর্ণ হলে	ডান উরুর মধ্যভাগের বহিরাংশে	চামড়ার নীচে

১৫-৪৯ বছর বয়সের মহিলাদের টিকাদান সময়সূচি

রোগের নাম	টিকার নাম	টিকার ডোজ	ডোজের সংখ্যা	টিকা শুরু করার সঠিক সময়	টিকাদানের স্থান	টিকার প্রয়োগ পথ
ধনুষ্ঠংকার	টিটি (টিটেনাস টক্সয়েড)	০.৫ এম এল	টিটি-১	১৫ বছর বয়স হলেই যথাশীঘ্র	বাহুর উপরের অংশে	মাংশপেশী
			টিটি-২	টিটি-১ পাওয়ার কমপক্ষে ২৮ দিন পর		
			টিটি-৩	টিটি-২ পাওয়ার কমপক্ষে ৬ মাস পর		
			টিটি-৪	টিটি-৩ পাওয়ার কমপক্ষে ১ বৎসর পর		
			টিটি-৫	টিটি-৪ পাওয়ার কমপক্ষে ১ বৎসর পর		

সম্প্রসারিত টিকাদান কর্মসূচি (ইপিআই)

স্বাস্থ্য অধিদপ্তর

গণপ্রজাতন্ত্রী বাংলাদেশ সরকার