

Tutorial-03

1. Let x be the weights of frying chicken.
 $x \sim N(500, 100^2)$ pdf of Normal distribution

$$f_x(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

$$x \sim N(\mu, \sigma^2)$$

a) $P(x < 650)$

$$= P\left(\frac{x-\mu}{\sigma} < \frac{650-500}{100}\right)$$

$$= P(z < 1.5)$$

$$= 0.9332$$

b) $P(x < 345)$

$$= P\left(\frac{x-\mu}{\sigma} < \frac{345-500}{100}\right)$$

$$= P(z < -1.55) \quad \text{or} \quad 1 - P(z < 1.55)$$

$$= 0.0606$$

c) $P(480 < x < 710)$

$$= P\left(\frac{480-500}{100} < z < \frac{710-500}{100}\right)$$


$$= P(-0.2 < z < 2.1)$$

$$= 1 - [P(z < -0.2) + P(z > 2.1)]$$

$$= 1 - [P(z < -0.2) + P(z < -2.1)]$$

$$= 1 - [0.4207 + 0.0179]$$

$$= 0.5614$$




d) $P(x > 590)$

$$= P\left(\frac{x-\mu}{\sigma} > \frac{590-500}{100}\right)$$

$$= P(z > 0.9) = 1 - P(z < 0.9)$$

$$= P(z < -0.9) = 0.1841$$



2) Let X be the IQ scores.

$$X \sim N(125, 10^2)$$

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

$$X \sim N(\mu, \sigma^2)$$

a) $P(X < 142)$

$$= P\left(\frac{X-\mu}{\sigma} < \frac{142-125}{10}\right)$$

$$= P(Z < 1.7)$$

$$= 0.9554$$



b) $P(X > 108)$

$$= P\left(\frac{X-\mu}{\sigma} > \frac{108-125}{10}\right)$$

$$= P(Z > -1.7)$$

$$= 1 - P(Z < -1.7)$$

$$= 1 - 0.0446 = 0.9554$$



c) $P(X < x) = 0.88$

$$P(Z < z) = 0.88$$

$$z = 1.175$$


$$\frac{x-\mu}{\sigma}$$

$$z = \frac{x-\mu}{\sigma}$$

$$x = \mu + \sigma z$$

$$= 125 + 10 \times 1.175$$


$$= 136.75$$



d) $P(X < x) = 0.36$
 $P(Z < z) = 0.36$
 $z = -0.36$
 $x = \mu + \sigma z$
 $= 125 + 10 \times -0.36$
 $= 121.4$

e) $P(X < 100)$
 $= P\left(\frac{X - \mu}{\sigma} < \frac{100 - 125}{10}\right)$
 $= P(Z < -2.5)$
 $= 0.0062$

Then the no. of applicants will not be admitted is $1500 \times 0.0062 = 9.3 \approx 9$



$z_{0.025} = -1.96$
 $z_{0.975} = 1.96$

$P(x_1 < X < x_2) = 0.95$

$P(-1.96 < Z < 1.96) = 0.95$

$x_1 = \mu + \sigma z$
 $= 125 + 10(-1.96)$
 $= 105.4$

$x_2 = \mu + \sigma z$
 $= 125 + 10(1.96)$
 $= 144.6$

3) Let x be the annual rainfall.

$$X \sim N(40, 5^2)$$

a) $P(X \geq 36)$


$$= P\left(\frac{X-\mu}{\sigma} > \frac{36-40}{5}\right)$$

$$= P(Z > -0.8)$$

$$= 1 - P(Z < -0.8)$$

$$= 1 - 0.2119$$

$$= 0.7881$$



b) $P(X < 38) + P(X > 50)$


$$= P\left(\frac{X-\mu}{\sigma} < \frac{38-40}{5}\right) + P\left(\frac{X-\mu}{\sigma} > \frac{50-40}{5}\right)$$

$$= P(Z < -0.4) + P(Z > 2)$$

$$= P(Z < -0.4) + P(Z < -2)$$

$$= 0.3446 + 0.0228$$

$$= 0.3674$$



4) ^ea) Let x be the ~~length~~ length of pregnancies

$$X \sim N(268, 15^2)$$


a) $P(X > 308)$

$$= P\left(\frac{X-\mu}{\sigma} > \frac{308-268}{15}\right)$$

$$= P(Z > 2.667)$$

$$= P(Z < -2.667)$$

$$= 0.0038$$



$$b) P(X < x_1) = 0.04$$

$$P(Z < z) = 0.04$$

$$z = -1.75$$

$$x_1 = \mu + \sigma z$$

$$= 268 + 15(-1.75)$$

$$= 241.75$$

5) Let x be ages of the residents.

$$x \sim N(70, 4^2)$$

$$a) P(60 < x < 80)$$

$$= P\left(\frac{60-70}{4} < Z < \frac{80-70}{4}\right)$$

$$= P(-2.5 < Z < 2.5)$$

$$= 1 - 2 \times P(Z < -2.5)$$


$$= 1 - 2 \times 0.0062$$

$$= 0.9876$$

$$b) n = 9$$

Sample mean of the means of the sample = 70
 $\mu_{\bar{x}} = 70$

$$c) \frac{s.d}{\sigma_{\bar{x}}} = \frac{\sigma}{\sqrt{n}} = \frac{4}{\sqrt{9}} = 1.333$$



$$d) P(68.5 < \bar{x} < 71.75)$$


$$= P\left(\frac{68.5 - 70}{\frac{4}{\sqrt{9}}} < Z < \frac{71.75 - 70}{\frac{4}{\sqrt{9}}}\right)$$

$$= P(-1.125 < Z < 1.3125)$$

$$= 1 - [P(Z < -1.13) + P(Z < -1.31)]$$

$$= 1 - [0.1292 + 0.0951]$$

$$= 0.7757$$



e)


$$P(\bar{x}_1 < \bar{x} < \bar{x}_2) = 0.95$$

$$P(-1.96 < Z < 1.96) = 0.95$$

$$\bar{x}_1 = \mu_{\bar{x}} + \sigma_{\bar{x}} Z$$

$$= 70 + 1.33(-1.96)$$

$$= 67.39$$



$$P(Z < z_1) = 0.025$$

$$z_1 = -1.96$$

$$\bar{x}_2 = \mu_{\bar{x}} + \sigma_{\bar{x}} Z$$

$$= 70 + 1.33(1.96)$$

$$= 72.61$$

6


$$* P(\text{Cost is less than } \$11.5) \\ = P(\text{No. of copies below } \overset{\text{total cost value}}{\$11.85 / \$0.05 = 237})$$

$$= P(X < 237)$$

$$= P\left(\frac{X - 250}{50} < \frac{237 - 250}{50}\right)$$

$$= P(Z < -0.26)$$

$$= 0.3974$$



$$\begin{aligned} &= P(Z < -0.26) \\ &= 1 - P(Z < 0.26) \\ &= 1 - 0.6026 \\ &= 0.3974 \end{aligned}$$

7) Let x be the net weights of boxes filled from the machine.

$$x \sim N(16, 0.3^2)$$

a) $P(x > 15)$


$$= P\left(\frac{x - \mu}{\sigma} > \frac{15 - 16}{0.3}\right)$$

$$= P(z > -3.333)$$

$$= 1 - P(z < -3.33)$$

$$= 1 - 0.0005$$

$$= 0.9995$$



b) $P(x > 16.5)$


$$= P\left(\frac{x - \mu}{\sigma} > \frac{16.5 - 16}{0.3}\right)$$

$$= P(z > 1.6667)$$

$$= 1 - P(z < 1.67)$$

$$= 1 - 0.9525$$

$$= 0.0475$$




c) $P(x > 16.4)$

$$= P\left(\frac{x - \mu}{\sigma} > \frac{16.4 - 16}{0.3}\right)$$

$$= P(z > 1.3333)$$

$$= P(z < -1.333)$$

$$= 0.0918$$



$$d) \quad n = 32$$

$$P(\bar{x} > 16.3)$$

$$= P\left(\frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}} > \frac{16.3 - 16}{\frac{0.3}{\sqrt{32}}}\right)$$

$$= P(Z > 5.6568)$$

$$= 0.0$$

$$=$$



$$e) \quad n = 32$$

$$P(\bar{x} < 15.8)$$

$$= P\left(\frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}} < \frac{15.8 - 16}{\frac{0.3}{\sqrt{32}}}\right)$$

$$= P(Z < -3.7712)$$

$$= 0.0001$$

8) $P(X \leq 90) \approx P(X < 90.5)$, a continuity correction is needed as the responses are binary & the counts have a binomial distribution with $n = 400, p = 0.20$
 $\mu = np$
 $= 400 \times 0.2$
 $= 80$

$$\sigma = \sqrt{npq}$$

$$= \sqrt{400 \times 0.2 \times 0.8}$$

$$= 8$$

$$np \geq 5, nq \geq 5$$

we can use normal approximation

$$P(X \leq 90) \approx P(X < 90.5)$$

$$= P\left(\frac{X - \mu}{\sigma} < \frac{90.5 - 80}{8}\right)$$

$$= P(Z < 1.31)$$

$$= 0.9049$$

The sample event is not unlikely assuming the claimed percentage of 20%, hence the claim is not unreasonable.

9

$np = 45 \geq 5$ & $nq > 5$ So we can use normal approximation.

$P(X \geq 57) \cong P(X > 56.5)$ By, a continuity correction is needed as the response are binary & the counts have a Binomial distⁿ with $n=100$, $p=0.45$.

$$\begin{aligned}\mu &= np \\ &= 100 \times 0.45 \\ &= 45\end{aligned}$$

$$\begin{aligned}\sigma &= \sqrt{npq} \\ &= \sqrt{100 \times 0.45 \times 0.55} \\ &= 4.9749\end{aligned}$$


$$P(X \geq 57) \cong P(X > 56.5)$$

$$= P\left(\frac{X-45}{4.9749} > \frac{56.5-45}{4.9749}\right)$$

$$= P(Z > 2.31)$$

$$= P(Z < -2.31)$$

$$= 0.0104$$



10) Let X be the length of the pea pods for its olive-green pea crop.

$$X \sim N(11.5, 1.6^2)$$

$$n = 16$$

$$a) \sigma_x = \frac{\sigma}{\sqrt{n}} = \frac{1.6}{\sqrt{16}} = 0.4$$

$$b) P(\bar{X} > 12.3)$$


$$= P\left(\frac{\bar{X} - 11.5}{0.4} > \frac{12.3 - 11.5}{0.4}\right)$$

$$= P(Z > 2)$$

$$= 1 - P(Z < 2)$$


$$= 1 - 0.9772$$

$$= 0.0228$$



$$c) P(\bar{x}_1 < \bar{X} < \bar{x}_2) = 0.68$$

$$P(Z_1 < Z < Z_2) = 0.68$$



$$z = \frac{\bar{x} - \mu_x}{\sigma_x}$$

$$\bar{x}_1 = \mu_x + z_1 \sigma_x$$

$$= 11.5 + (-1)(0.4)$$

$$= 11.1$$

$$\bar{x}_2 = \mu_x + z_2 \sigma_x$$

$$= 11.5 + 1(0.4)$$

$$= 11.9$$

$$\begin{aligned}
 \text{a) Mean} &= np \\
 &= 1500 \times 0.7 \\
 &= 1050
 \end{aligned}$$

$$\begin{aligned}
 \text{standard deviation} &= \sqrt{np(1-p)} \\
 &= \sqrt{1500 \times 0.7 \times 0.3} \\
 &= 17.7482
 \end{aligned}$$

$$\text{b) } np \geq 5, nq \geq 5$$

So, we can use normal approximation
 $X \sim N(1050, 17.7482^2)$

$$\begin{aligned}
 P(X \geq 1000) &= P\left(\frac{X-\mu}{\sigma} \geq \frac{1000-1050}{17.7482}\right) \\
 &\cong P(X > 999.5) \rightarrow
 \end{aligned}$$

$$= P(Z \geq \frac{-2.8453}{-2.8171})$$

$$= 1 - P(Z \leq -2.85)$$

$$= 1 - 0.0022$$

$$= 0.9978$$



$$\text{c) } P(X > 1200) = P\left(\frac{X-\mu}{\sigma} > \frac{1200-1050}{17.7482}\right)$$

$$\cong P(X > 1200.5)$$

$$= P(Z > \frac{8.4797}{8.4516})$$

$$= 0$$

d)

$$N = 1700$$

$$\text{mean} = 1700 \times 0.7 \\ = 1190$$

$$S.D = \sqrt{1700 \times 0.7 \times 0.3} \\ = 18.8944$$

$$\begin{aligned} P(X > 1200) &= P\left(\frac{X - \mu}{\sigma} > \frac{1200 - 1190}{18.8944}\right) \\ &\equiv P(X > 1200.5) \rightarrow P(Z > 0.5557) \\ &= P(Z > 0.5557) \\ &= 1 - P(Z < 0.5557) \\ &= 1 - 0.7122 \\ &= 0.2878 \\ &= 0.2877 \end{aligned}$$



12
★

$$f_X(x) = \lambda e^{-\lambda x} \quad x > 0$$

$$\lambda = \frac{1}{5} = 0.2/\text{year}$$

$$\frac{14}{12} = 1.167 \text{ years}$$

$$P(T < \frac{14}{12}) = \int_0^{\frac{14}{12}} 0.2 e^{-0.2x} dx$$

$$= -e^{-0.2x} \Big|_0^{\frac{14}{12}}$$

$$= -e^{-0.2 \times \frac{14}{12}} + 1$$

$$= -0.7919 + 1 = 0.2081$$

13

Let x be the lifetime of a particular fan.

$$x \sim \exp(\lambda = 0.0003)$$

$$\text{p.d.f. } f(x) = \lambda e^{-\lambda x} \quad x > 0$$

$$P(x > 10000) = \int_{10,000}^{\infty} \lambda e^{-\lambda x} dx$$

$$= -e^{-\lambda x} \Big|_{10,000}^{\infty}$$

$$= 0 + e^{-0.0003 \times 10,000}$$

$$= e^{-3}$$

$$= 0.04978$$

$$= 0.0498$$

$$\lambda = 0.00035$$

$$P(x > 10,000) = \int_{10,000}^{\infty} \lambda e^{-\lambda x} dx$$

$$= -e^{-\lambda x} \Big|_{10,000}^{\infty}$$

$$= 0 + e^{-0.00035 \times 10,000}$$

$$= e^{-3.5}$$

$$= 0.030197$$

$$= 0.0302$$

It will fewer fans compare with earlya one.

14) Let x be the time interval between successive barges.

$$\text{mean} = 8$$

$$\frac{1}{\lambda} = 8$$

$$\lambda = \frac{1}{8} = 0.125$$

$$x \sim \exp\left(\frac{1}{8}\right)$$

$$f_x(x) = \lambda e^{-\lambda x} \quad x > 0$$

$$a) P(X < 5) = \int_0^5 \lambda e^{-\lambda x} dx$$

$$= -e^{-\lambda x} \Big|_0^5$$

$$= -e^{-\frac{1}{8} \cdot 5} + 1$$

and

$$= 0.4647$$

$$b) P(T > t) = \int_t^{\infty} 0.125 e^{-0.125t} dt$$

$$= -e^{-0.125t} \Big|_t^{\infty}$$

$$= e^{-0.125t}$$

$$e^{-0.125t} = 0.95$$

$$e^{-\frac{t}{8}} = 0.95$$

$$-\frac{t}{8} = \ln 0.95$$

$$t = -8 \ln 0.95$$

$$= 0.4103$$