

BANGLADESH JOURNAL OF NEUROSURGERY

□ VOLUME 5 □ NUMBER 2 □ JANUARY 2016

CONTENTS

Editorial	29
Original Articles	
Primary bone tumour of the spine: Surgical treatment and Histopathological analysis <i>Md. Shafiul Alam, Md. Shafiqul Kabir Khan, Monsur Ahmed, Muhammad Shamsul Arefin, Kazi Hafiz Uddin, Mohammad Hossain, Md. Rezaul Amin</i>	31
Outcome of Conservative Management of Traumatic Acute Subdural Haematoma in Relation to Underlying Brain Injury <i>Sanat Kumar saha, Mohammad Abu Sayed, Joynul Islam, Shafiul Alam, ATM Asadullah, Misbah Uddin Ahamed, Shamsul Islam Khan</i>	37
Signs and Symptoms of Early VP Shunt Infection in a Tertiary Hospital <i>Kaisar Haroon, Tania Taher, Mokhlasur Rahman, Kanak Kanti Barua</i>	41
Assessment of Serum Homocysteine Level in Epileptic Patients on Carbamazepine Treatment <i>Md. Abu Hanif, M A Hannan, M Iqbal Arslan, SK Mahbub Alam, Mohammad Hossain, Md. Aynul Hoque, Niloy Ranjan Roy</i>	44
Case Reports	
Tubercular Spinal Epidural Abscess Involving the Lumbo-Sacral Region: A Case Report with Review of Literature <i>Akhalaque Hossain Khan, Ranjit Kumar Chaurasiya, Narendra Shalike, Mohammed Aminul Hasnat, Bipin Chaurasia, Abhishek Chaturbedi, Asifur Rahman, Rezaul Amin</i>	48
Giant Ganglioneuroma of Thoracic Spine: A Case Report <i>Das S, Alam K, , Rashid MM, Islam MM, Mahbub H, Khan MSI</i>	51



**OFFICIAL ORGAN OF
BANGLADESH SOCIETY OF NEUROSURGEONS**

BANGLADESH JOURNAL OF NEUROSURGERY

Volume 5, Number 2, January 2016

OFFICIAL ORGAN OF BANGLADESH SOCIETY OF NEUROSURGEONS

Shahbag, Dhaka -1000

Email: bdneurosurgeons@yahoo.com

Website: <http://www.bsnsbd.org>

EDITORIAL BOARD

Chief Editor	: Prof. Kanak Kanti Barua
Executive Editor	: Dr. Asif Moazzam Barkatullah
Managing Editor	: Dr. Md. Raziul Haque
Assistant Editor	: Dr. Md. Shafiul Alam Dr. Md. Rezaul Amin Dr. Md. Rafiqul Islam
Members	: Prof. Abul Khair Prof. Sk. Sader Hossain Prof. ATM Mosharef Hossain Prof. Md. Zillur Rahman Dr. Md. Kamal Uddin Dr. Akhlaque Hossain Khan Dr. Asit Ch Sarker Dr. Mohammad Hossain Dr. Sukrity Das Dr. Haradhan Debnath Dr. Asifur Rahman

INTERNATIONAL EDITORIAL ADVISORS

Prof. Madjid Samii	Prof. P.S. Ramani
Prof. Kenji Ohata	Prof. Maurice Choux
Prof. Atul Goel	Prof. R.P. Sen Gupta
Prof. Yoko Kato	Prof. Timothy Lee

Editorial Policy

Bangladesh Journal of Neurosurgery (BJNS) is a double blind peer reviewed journal. It started publishing from July 2011. It is published biannually in January & July each year by Bangladesh Society of Neurosurgeons (BSNS). It publishes original article based on laboratory work, field work, clinical trial and various other studies by scientific means related to disciplines of Neurosurgery conducted in different institutes in Bangladesh and other countries. It also accepts review article, meta analysis, case report, short communication, letter to editor.

All rights reserved. No part of any publication may be reproduced, stored in retrieval system or transmitted in any form without the written permission of publisher except for nonprofit educational purpose.

The editors reserve the customary right to style and if necessary may shorten the material accepted for publication and to determine the priority and time of publication. Editors assume that work is based on honest observation. It is not task of the editors to investigate scientific fraud paper. The editorial board will not take the responsibility for any misleading inaccurate information or any statement or opinion.

ADDRESS OF CORRESPONDENCE

Prof. Kanak Kanti Barua

Chief Editor BJNS

Room 505, Department of Neurosurgery

Bangabandhu Sheikh Mujib Medical University

Shahbag, Dhaka

Phone: +88-01711530424

Email: bdneurosurgeons@yahoo.com

INFORMATION FOR THE CONTRIBUTORS

Manuscripts prepared following the “Uniform Requirements for manuscripts submitted to biomedical Journals” is acceptable to this journal for publishing. Authors are advised to visit the Web site www.bsnsbd.org for this purpose.

Editorial scope:

- The Bangladesh Journal of Neurosurgery (BJNS) is intended to promote prompt publication of concise, scientific article based on the study in all fields of medical and health sciences.
- Submitted manuscripts should not be previously published or being considered for publication elsewhere.
- All submitted articles will undergo double blind peer review as per recommendations by subject specific experts selected by editors.
- Reviewed manuscripts will be sent to the corresponding author for appropriate response if it is indicated.
- Acceptance is based on significance, originality and clarity and fulfillment of the criteria of the publication policy of this journal.
- The Editor-in-Chief will make all final decisions regarding acceptance.
- Selection of the reviewed and accepted manuscripts intended for publication in a particular issue will be decided by Editorial Board.
- Rejected manuscript will be returned if accompanied by stamped or self-addressed envelop.
- Upon acceptance for publication the copy right of the paper automatically transfers to the Bangladesh Society of Neurosurgeons (BSNS) and will not be published elsewhere either in part or whole without written permission of the copyright holder.
- Review article should be written by a subject expert.

Ethical aspects:

- Manuscripts based on the study should have been conducted according to the ethical standards laid down in the 1964 Declaration of Helsinki revised 2000.
- Manuscript must contain a statement in the method section that all human subjects involved studies have been approved by appropriate ethical committee after careful examination of the ethical aspects.

- Permission of the patients or their families to reproduce photographs of the patients where identity is not disguised
- Author should obtain written permission to reproduce any table, illustration from any other source.

Manuscript submission::

The authors are requested to strictly follow the guide lines below for submission of manuscript to for publication. The following documents with manuscripts are to be submitted for publication.

- A covering letter addressed to the Chief Editor of the journal (Sample given at the end).
- A Title page ((Sample given at the end).
- Abstract and key words in the first page followed by the text, prepared in the format of IMRAD.
- Authors are requested to submit their articles in the web bsnsbd.com as instructed.
- On special cases, submission through E-mail with file attachment of all documents is welcome.

Covering letter:

- All authors must sign after seeing the manuscript with the statement that they are the only authors.
- The corresponding author should mention the contribution of each author to the work.
- It should contain a declaration that this manuscript has not been submitted elsewhere or not under consideration in any journal.
- It should clearly indicate the publication type (Original/Review/Case report/letter etc.)
- Should also mention the expected benefit of the medical science from publishing of this article.

Authors are requested to submit new and revised manuscript and correspondence to:

Prof. Kanak Kanti Barua

Chief Editor BJNS

Room 505, Department of Neurosurgery
Bangabandhu Sheikh Mujib Medical University
Shahbag, Dhaka

Phone: +88-01711530424

Email: bdneurosurgeons@yahoo.com

Manuscript Organization:**Typing**

- Double spaced throughout with Justified and 2.5 cm margins
- Font type is Times New Roman with size 12.
- Manuscript should have uniform style, correct journal format, carefully proofread for grammar, spelling and punctuation

Manuscript format:

In general, original article should be divided into following sections: Title page, Abstract, Text, Acknowledgement, References, Tables with titles and foot notes, alternatively graph with title and Illustrations with legends. Each of the sections is to start on a separate page. Pages should be numbered consecutively beginning from the abstract.

Title page:

- Title of the article.
- Names of all authors with their designation and institutional affiliations
- Name of the department and institute where the study was undertaken
- Name of the corresponding author with contact address, telephone number, Email address
- Disclosure of conflict of interest if any
- Disclosure of sources of funding or sponsor
- A short running head
- Number of tables given
- Number of figures given
- Word count of abstract

Abstract:

- Title of the article
- Structured with headings (Background, Objectives, Methods with statistical analysis, Result & Conclusion).
- Authors name should not be given
- Preferably within about 300 words
- Avoid abbreviations in the title and abstract except standard abbreviation
- A non structured abstract is suggested for review article and case report.

Text:

Text should be arranged into following sections: Introduction, Methods, Results, Discussions, Conclusion, Acknowledgement & References

Introduction:

- Statement of the problem with a short discussion of its importance and significance
- Review of the literature related to the problem with pertinent reference
- Objectives/ hypothesis/ benefits expected stated in 1-2 paragraph

Methods:

- Study type, place and time
- Description of study variables
- Description of study subjects and grouping
- Selection criteria
- Approval of the study involving human subjects by ethical review committee and description of the ethical aspects in such study
- Description of procedure, methods, apparatus, drugs or chemicals as applicable
- Description of statistical procedure with enough detail to enable a knowledgeable reader with access to the original data to verify the reported results

Results:

- Present result in logical sequence in text, table and illustration with most important finding first
- Describe without comment
- Restrict number of table and figure needed to support assessment of paper
- Do not duplicate data in table and figure

Table:

- Simple self explanatory with brief title, not duplicate in text
- Each table should be numbered in Romans and printed in separate page
- Do not use internal horizontal and vertical rules
- Uses of many tables are not encouraged

Illustration:

- All illustrations must be numbered consecutively in Arabic numerals as they appear in the text

- Submit print photograph of each illustration along with its electronic file
- Figure number, title of manuscript, name of the corresponding author and arrow indicating top should be written on a sticky label affixed on the back of each photograph
- Scanned picture, graph, chart with title and figure number should be printed on separate page and its original data presentation file should be inserted in the CD along with text

Legend:

- Must be typed in a separate sheet of paper
- Photomicrograph should indicate the magnification, internal scale and the method of staining
- All drugs should be mentioned in their generic form. The commercial name may be used in parenthesis

Acknowledgement:

- Individuals, institution, sponsor, organization or bodies can be acknowledged in the article for their contribution or financial or any form of assistance to the work

References:

- For reference, use author number style (Vancouver) which is based on an ANSI standard adapted by the National Library of Medicine(NLM)
- References should be numbered consecutively in the order in which they are first mentioned in the text
- Identify references in the text, tables and legends by Arabic numerals in superscript
- All citations to electronic references should be presented in numbered references following the text
- The titles of the journals should be abbreviated according to the style used in Index Medicus
- Write names of 6 authors followed by et al. if authors number is more than six.
- The reference list is not usually checked by the editorial staff or reviewer. It is the total responsibility of author to provide accurate information.

Standard journal article:

Example:

Khalil M, Chowdhury MAI, Rahman H, Mannan S, Sultana SZ, Rahman MM, et al Splenic Mass and its

relation to age, sex and height of the individual in Bangladeshi People. J Bangladesh Soc Physiol 2008; 3(1): 71-78.

Journal article with organization as author:Journal article with organization as author:

American diabetes Association.diabetes update. Nursing, 2003 Nov; Suppl: 19-20.

Journal article with multiple organization as author:Journal article with multiple organization as author:

American Dietetic association; Dietitians of Canada. Position of Dietetic association and Dietitians of Canada: nutrition and women's health. J Am Diet Assoc 2004 Jun; 104(6): 984-1001.

Journal article with Governmental body as author:Journal article with Governmental body as author:

National Institute on Drug Abuse (US); Caribbean Epidemiology Centre; Pan American Health Organization; World Health Organization. Building a collaborative research agenda: drug abuse and HIV/AIDS in the Caribbean 2002-2004. West Indian Med J. 2004 Nov; 53 Suppl 4: 1-78.

Standard book with initials for authors:Standard book with initials for authors:

Eyre HJ, Lange DP, Morris LB. Informed decisions: the complete book of cancer diagnosis, treatment and recovery. 2nd ed. Atlanta: American Cancer Society; 2002. 768 p.

Contributed chapter of a book:Contributed chapter of a book:

Rojko JL, Hardy WD. Feline leukemia virus and other retroviruses. In:Sherding RG, editor. The cat: diseases and clinical management. New York: Churchill Livingstone; 1989. p. 229-332.

Conference Proceedings:Conference Proceedings:

Pacak K, Aguilera G, Sabban, E, Kvetnansky R, editors. Stress: current neuroendocrine and genetic approaches. 8th symposium on Catecholamines and Other Neurotransmitters in Stress: 2003 Jun 28-July 3; Smolenice Castle(place of conference), Slovakia. New York(place of publication): New York Academy of Sciences (publisher); 2004 Jun. 590 p.

Scientific and Technical Reports:Scientific and Technical Reports:

Page E, Harney JM. Health hazard evaluation report. Cincinnati(OH)(Place of publication: National Institute for Occupational Safety and Health (US)(Publisher); 2001 Feb.24 p(Total number of pages). Report No.: HETA2000-0139-2824

Dissertation & Thesis:

Entire Reference

Kempner JL. Aching heads, making medicine: gender and legitimacy in headache (title) [dissertation]. [Philadelphia]: University of Pennsylvania;2004.271p.

Alam M. Study of Heart Rate Variability in Adolescent Athletes [M Phil thesis]. [Dhaka]: Bangabandhu Sheikh Mujib Medical University; 2008. 178p.

Part of Dissertation & Thesis:

Mackowski MP. Human factors: aerospace medicine and the origins of manned space flight in the United States[dissertation].[Tempe(AZ)]:Arizona State University;2002 May. Part 2,Space medicine;p.188-377.

Alam M. Heart Rate Variability in Adolescent Athletes[M Phil thesis]. [Dhaka (Bangladesh)]: Bangabandhu Medical University;2008 July. Appendix (Name of the part) 4(Number of the part),Classification of Physical Activity Intensity(Title of the part). p.7(Location of the part).

Standard journal article on the Internet:

Kaul S, Diamond GA. Good enough: a primer on the analysis and interpretation of noninferiority trials. Ann Intern Med [Internet].2006Jul4 [cited 2007 Jan 4];145(1):662-9.

Available from :<http://www.annals.org/cgi/reprint/145/1/62.pdf>

Journal article on the Internet with organization (s) as author:

National Osteoporosis Foundation of South Africa. Use of generic alendronate in treatment of osteroporosis. S Afr Med J [Internet].2006 Aug[cited 2007 Jan 9];9(8):696-7. Available from: <http://blues.sabinet.co.za/WebZ/Authorize?>

Journal article on the Internet with governmental body as author:Journal article on the Internet with governmental body as author:

Centers for Disease Control and Prevention(US), National center for HIV/AIDS, Hepatitis, STD, and TB Prevention and control of tuberculosis in correctional and detention facilities: recommendations from CDC. Endorsed by the Advisory Council for the elimination of tuberculosis, the national Commission of Correctional Health Care and the American correctional Association. MMWR R Rep[Internet].2006Jul7[cited2007Jan 9];55(RR-9):1-44.Available from:<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5509a1.htm>

Journal article on the Internet with no author:

Prevention strategies for asthma-secondary prevention. CMAJ [Internet].2005 Sept[cited 2007Jan 5];173 (6Suppl):S25-7.Available from:http://www.cmaj.ca/cgi/content/full/173/6_suppl/S25

Journal article on the Internet without standard volume, issue or article number:

Jacobs JL, Lee MT, Lindberg M, Kamin C. Problem based learning, multimedia paucity of behavioral issue learning. Med Educ. Online[Internet].2005[cited2005]: [5p]. Available from:<http://www.med-ed-online.org/pdf/10000006.pdf>

For Subscription:

Annual subscription for

Institution	: Taka 300.00
Individuals	: Taka 200.00
Senior faculty	: Taka 200.00
Junior faculty	: Taka 100.00
Overseas patrons	: US\$ 20.00

Send subscription to:

Prof. Kanak Kanti Barua

Chief Editor BJNS

Room 505, Department of Neurosurgery

Bangabandhu Sheikh Mujib Medical University

Shahbag, Dhaka

Phone: +88-01711530424

Email: bdneurosurgeons@yahoo.com

DISCLOSURE OF CONTRIBUTION OF AUTHORS

Corresponding author:

As principal investigator, Dr.....

had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept & design:

Acquisition of data:

Analysis and interpretation of data:

Drafting of the manuscript:

Critical revision of the manuscript for important intellectual content:

Statistical analysis:

Obtained funding:

Administrative, technical, or material support:

Supervision:

TITLE PAGE OF ARTICLE

The title of the article:

.....

1. Authors' names and institutional affiliations:1,2,
.....3,4,5

1.Medical College, Dhaka

2. Professor & Chairman of the Department of
Bangabandhu Sheikh Mujib Medical University (BSMMU), Bangladesh

3. Associate Professor, Department of
Bangabandhu Sheikh Mujib Medical University (BSMMU), Bangladesh.

4. Assistant Professor, Department of
Bangabandhu Sheikh Mujib Medical University (BSMMU), Bangladesh

5. Assistant Professor, Department of
Bangabandhu Sheikh Mujib Medical University (BSMMU), Bangladesh

2. The name of the department(s) and institution(s) to which the work should be attributed:
Department of
Bangabandhu Sheikh Mujib Medical University (BSMMU), Bangladesh

3. Disclaimers any: Yes, subject to non profit purpose

4. Corresponding author name:

Mailing address:

.....

Telephone:

E-mail address:

5. The name and address of the author to whom requests for reprints should be addressed:

Mailing address:

.....

7. Source(s) of support in the form of grants, equipment, drugs, or all of these- [Self finance]

8. A short running head:

9. Word counts of abstract :

10. The number of tables :

11. The number of figures:

FORWARDING LETTER FOR SUBMISSION TO BJNS

Date.....

To
Chief Editor,
Prof.

.....
Bangladesh Journal of Neurosurgery (BJNS)
Sub: Submission of manuscript

Dear Sir,

We are submitting our manuscript entitled,by,1,
.....2,3,4,
.....5. for publication in your journal. This article **has not been published**

or
submitted for publication elsewhere.

We believe that this article may be of value to medical professionals engaged in Neurosurgery.

We therefore, hope that you would be kind enough to consider our manuscript for publication in your journal
as original / Review article / Case Report.

Thanks and best regards

More than 17,000 people in the United States and 18,000 in India are diagnosed each year with a brain tumor. Unfortunately we do not have any accurate data for Bangladesh.

Some tumors are benign (noncancerous). Noncancerous tumors can usually be removed and are not likely to recur. Other tumors are malignant (cancerous). These tumors interfere with vital functions and are life threatening. Cancerous brain tumors usually grow rapidly, crowding and invading tissue.

Primary brain tumors (gliomas) start in the brain and affect the central nervous system (CNS). They can be noncancerous or cancerous. Secondary brain tumors, which are 10 times more common, are cancers that originated elsewhere in the body and have metastasized (spread) to the brain. Secondary tumors are about 3 times more common than primary tumors of the brain.

People with the following conditions or characteristics may be at risk for developing a brain tumor:

- Radiation exposure
- Exposure to pesticides, herbicides, fertilizer
- Certain occupations, such as lead, petroleum, plastic, rubber, and textile workers, as well as aircraft and vehicle operators
- Exposure to electromagnetic fields
- Certain viruses, especially Epstein-Barr virus
- Certain genetic disorders
- People who have had transplants and individuals with AIDS
- A small percentage of brain tumors may be hereditary (approximately 5% of gliomas)
- People over age 45

Studies have not shown an association between brain cancer and cell phone use.

Now it will be the right time to ask all the neurosurgeons of Bangladesh to take an initiation in opening and maintaining a combined central brain tumor registry.

Primary Bone Tumour of the Spine: Surgical Treatment and Histopathological Analysis

Md. Shafiul Alam¹, Md. Shafiqul Kabir Khan², Monsur Ahmed³, Muhammad Shamsul Arefin⁴, Kazi Hafiz Uddin⁵, Mohammad Hossain⁶, Md. Rezaul Amin⁶

Abstract

Background: Primary bone tumour of the spine are rare but not uncommon. These include both benign and malignant lesions. Surgical resection and stabilization in some cases are the treatment of choice. Adjuvant radio and/or chemotherapy are needed in malignant cases.

Methods: We retrospectively reviewed all the primary bone tumour of the spine that were surgically treated in the department of Neurosurgery of Dhaka Medical College Hospital and National Institute of Neurosciences and Hospital Dhaka, Bangladesh between 2010 and 2015.

Results: Eighteen cases were included that presented at a median age of 36 years (range 14 to 49 years). Pain was the most common presenting symptom. Three patients had significant neurological deficits at time of presentation and in most cases there was an improvement after surgery. A variety of surgical strategies was employed with the use of adjuvant radio- or chemotherapy in six cases. Six benign and twelve malignant tumours were treated.

Conclusions: Primary bone tumours of the spine are often associated with a significant delay in diagnosis. Surgical strategy should be individualized for each case. Acceptable results can be achieved with early surgery and adequate chemo-radiotherapy.

Key words: primary bone tumour, spine, surgical treatment, histopathology.

Bang. J Neurosurgery 2016; 5(2): 31- 36

Introduction:

Primary bone tumour of the spine is a heterogynous group of conditions that includes both benign and malignant tumours, with only 6% of these tumours considered malignant.¹ The spine is a frequent site of metastasis, but primary bone tumours of the spine are rare and make up only 4.2% of spinal tumours.² Eleven per cent (11%) of primary musculoskeletal tumours occur in the spine with certain tumours having an affinity for the spine. Although rare, these tumours offer a considerable diagnostic dilemma and therapeutic challenge. Timely diagnosis is crucial, as immediate attention could affect overall prognosis.³

The most common primary tumour of the spine is metastatic deposits. A number of both benign and

malignant tumours may arise primarily from the spine. Among benign tumour osteoid osteoma, osteoblastoma, osteochondroma, giant cell tumour (GCT), aneurysmal bone cyst (ABC), eosinophilic granuloma (EG), haemangioma are common. Among malignant lesion malignant lymphoma, sarcoma, ewing sarcoma, osteosarcoma, chondrosarcoma, malignant fibrous histiocytoma (MFH), fibrosarcoma, chordoma, plasmacytoma/multiple myeloma and peripheral primitive neuroectodermal tumour (pPNET) are common.

Primary bone tumours are more common in the thoracic and lumbosacral regions than in the cervical spine.⁴ These tumours occur according to a typical anatomical distribution within the vertebra. In general, malignant tumours occur more frequently in the anterior elements and benign tumours in the posterior elements.³⁻⁵

The aims of treatment of these axial neoplasms include complete resection if feasible, preservation of neurological function and stabilisation of the spine if needed.⁴ The aim of our study was to review the primary bone tumours of the spine treated at our unit regarding the presentation, the surgical strategy employed and the outcome.

Materials and Methods:

This is a retrospective study. All the primary bone tumours of the spine that received surgical treatment

1. Associate Professor, Department of Gamma Knife, National Institute of Neurosciences & Hospital.

2. Associate Professor, Department of Neurosurgery, Dhaka Medical College Hospital.

3. Resident Surgeon, Department of Neurosurgery, National Institute of Neurosciences & Hospital.

4. Junior Consultant, , Department of Neuroanaesthesiology, , National Institute of Neurosciences & Hospital.

5. Registrar, Department of Clinical Neurosurgery, National Institute of Neurosciences & Hospital.

6. Associate Professor, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka

Address of Correspondence: Associate Professor, Department of Gamma Knife, National Institute of Neurosciences & Hospital, E mail: dr_chapal@hotmail.com, Tel: 01711567358.

at the department of Neurosurgery of Dhaka Medical College Hospital and National Institute of Neurosciences and Hospital Dhaka, Bangladesh from 2010 to 2015 were included in this study. We excluded all secondary deposition in the spine. We treated 18 cases during this period that were included according to these criteria. A case note and imaging review was done where patient demographics, details of presentation, clinical information and the management strategy were analyzed. All complications and mortalities were recorded.

Results:

Between 2010 and 2015, a total 18 patients were diagnosed as primary bone tumour of the spine. The median age at presentation was 36 years (range 14 to 49 years) with eleven(61.12%) male and seven(38.88%) female patients affected. The median follow-up was 18 months (range 3 to 60 months).

Table-I
Distribution of patient by age.

Age group	Frequency	Percentage
0 to 10 years	0	0
11 to 20 years	3	16.67
21 to 30 years	7	38.89
31 to 40 years	4	22.22
41 to 50 years	4	22.22
Total	18	100

Table-II
Distribution of patient by sex.

Sex	Frequency	Percentage
Male	11	61.12
Female	7	38.88
Total	18	100

Table-III
Distribution of patient by site of involvement.

Site of involvement	Frequency	Percentage
Cervical	1	5.55
Dorsal	7	38.89
Lumber	6	33.34
Sacrum	4	22.22
Total	18	100

The most utilized diagnostic tool was MRI and CT scan of the spine. It helped to diagnose the case as well as planning of surgery. X ray is no longer a standard diagnostic tool of spine tumour, it was used in all cases as a complimentary diagnostic tool. It also helped to find out the extra spinal involvement like rib cases in two cases. Pre-operative CT guided FNAC was done in seven cases. Post-operative histopathological study was done in all cases.

Primary bone tumour of the spine most commonly involved the dorsal (38.89%) and the Lumber (33.34%). It affect the sacral (22.22%) and the cervical spine (5.55%) less frequently. (Table-3)

Table-IV
Distribution of patient by histopathological findings.

Type	Histopathology	Frequency	percentage
Benign	-Osteoid osteoma	1	5.55
	-Haemangioma	1	5.55
	-Giant cell tumour	1	5.55
	-Aneurysmal bone cyst	2	11.11
	-Eosinophilic granuloma	1	5.55
Malignant	-Myxoid Chondrosarcoma	1	5.55
	-ewing's sarcoma	2	11.11
	-Chondrosarcoma	2	22.22
	-Chordoma	3	33.33
	-Multiple myeloma	2	22.22
	-pNET	2	22.22
Total		18	100

Pain was the most common presenting symptom (15 from 18 cases) and four cases presented with pain alone. Pain was mostly axial pain in the affected region (11 of 18 cases), although three patients presented with chest pain and one with flank pain. Other presenting symptoms included neurological symptoms and odynophagia. One patient presented with a recurrence of a malignant aggressive tumour (Ewing sarcoma of sacral spine) following incomplete resection at another unit. Ten patients had no neurological deficit at time of presentation. Two patients presented with radicular symptoms and another three had significant neurological deficits at time of presentation. One case presented with significant neurological deficit with a diagnosis of benign aggressive lesions (Aneurysmal bone cyst). Two of these cases had significant improvement after surgery. We had no cases of neurological deterioration following surgery. We treated six benign and twelve malignant tumours. The histopathological diagnoses are summarised in Table 4. In all cases the surgery was performed by the senior author. A

variety of surgical strategies was utilised. An intralesional resection or debulking of the tumour was performed in six cases and a marginal resection was performed in nine cases to correlate with diagnosis. In six patients a fusion was performed by transpedicular screws and rods. Adjuvant therapy was used where indicated. Two of the recurrent cases

were in malignant tumours and one in benign aggressive tumour. In the one recurrent malignant tumour a repeat attempted marginal resection was performed. The other two recurrent cases were not re-operated as resection was not possible with acceptable morbidity. We had three deaths in the series on further follow up.




Fig.-1: CT Scan of the lumbosacral spine showing Sacral Chordoma.




Fig.-2: MRI of the dorsal spine showing Ewing Sarcoma of D-9 vertebrae with involvement of ribs.




Fig.-3: MRI showing extensive Chondrosarcoma of the dorsal spine.




Fig.-4: MRI of the dorsolumbar spine showing Aneurysmal Bone Cyst of the L-2 vertebrae.




Fig.-5: Pre and post-Operative X-ray of L-2 Aneurysmal Bone Cyst.

Discussion:

The most common presenting complaint of patients with primary bone tumour of the spine is pain, with roughly 60% complaining of axial pain and 25% of radicular symptoms.³ The presence of a neurological deficit carries a worse prognosis.¹¹ Pain at rest or night pain is red flag symptoms that warrant further investigation. Spinal deformity is obvious when present, but occurs in less than 10% of patients. The most common cause of a painful scoliosis in adolescents is an osteoid osteoma.⁵ A delay in diagnosis is common and our finding is confirmed in other series.⁴ This is due to the presenting symptoms often being vague, pain often being the only symptom, and the initial radiographs often only showing subtle or no abnormalities leading to the symptoms not being properly investigated. Most patients report a slow, gradual onset of pain in the involved area. In benign tumours of the spine an average of 26 weeks of symptoms before presentation are reported.⁴ This was confirmed in our series where the average delay in presentation was 7 months.

A variety of classifications systems is used for primary bone tumours of the spine. The Enneking staging system¹² was initially described for primary bone tumours of the appendicular skeleton, but is applied to tumours of the spine as well. It prognosticates as well as guides surgical resection margins and has been validated in the literature. The Weinstein-Boriani-Biagini surgical staging system¹³ is an anatomical classification based on the Enneking system that describes the number of vertebra as well as the number of sectors within a vertebra involved. Recommendations are then made for the most appropriate approach to resection of the lesion. We used the Enneking staging system. According to this system we treated six benign and twelve malignant tumours. Of the benign group five could be classified as active and seven as aggressive. Appropriate treatment may be observational (Enneking grade 0 latent lesions) or surgical (most other lesions), depending on the level of pain, instability, neurologic compromise and the natural history of the lesion.⁵ The aims of surgery are complete resection of the lesion where feasible with preservation of neurological function. We used a variety of surgical strategies customised to the patient and the tumour. In peripheral malignant sarcomas the most important factor affecting survival is complete resection of the tumour with a wide

margin.^{14,15} There are however no reports prescribing surgical margins in spinal tumours. Traditional intralesional resection in a piecemeal fashion is thought to increase the likelihood of local recurrence. True wide resection would cause unacceptable morbidity as it would involve resecting the segment of spinal cord. Total en bloc spondylectomy in a single stage from an all posterior approach was described by Tomita.¹⁴ Here wide margins are achieved except at the pedicles and occasionally the spinal canal. This approach decreases local recurrence rates.¹⁶⁻¹⁸ In benign aggressive lesions surgical eradication also provides the best long-term cure.^{6,18} We had three cases of local recurrence in our series. Two were in malignant tumours where a marginal resection was performed. The other recurrent case was a thoracic giant cell tumour encapsulating the aorta and forcing the surgeon to perform an intralesional resection.

Recent advances in chemotherapy have led to improved survival in malignant primary bone tumours, including in the spine.^{7,19} Adjuvant radiotherapy is indicated in some malignant and benign aggressive lesions.^{6,9} The use of local radiotherapy does increase the risk of major local complications, including sepsis.^{8,10} We used radio- and chemotherapy in a number of cases as guided by our local oncology unit. The main limitations of this study are the retrospective character and the relatively small numbers.

Conclusion:

Most primary bone tumour of the spine in our population occurred in adults and was malignant. Surgical treatment resulted in pain reduction and neurological improvement, allowing a rapid adjuvant therapy in good clinical condition. An experienced multidisciplinary team of tertiary care center should perform diagnosis and treatment of these extremely rare tumors. In order to optimize management and elaborate guidelines, we encourage national or international registries, so that correct epidemiology, better classification and understanding, improved diagnostics, superior treatment and an overall better outcome may be achieved. We conclude that primary bone tumours of the spine are often associated with a significant delay in diagnosis. Surgical strategy and adjuvant therapy should be individualized for each case. Acceptable results with minimal complications can be achieved with this approach.

References:

1. Boriani S, Biagini R, De Iure F, et al. Primary bone tumors of the spine: a survey of the evaluation and treatment at the Istituto Ortopedico Rizzoli. *Orthopedics*. 1995;18(10):993–1000.
2. Davis JH. Anatomical classification and surgical considerations: Primary spinal tumours. An overview. *South African Med J*. 2011;10(3):26–30.
3. Sansur CA, Pouratian N, Dumont AS, Schiff D, Shaffrey CI, Shaffrey ME. Part II: spinal-cord neoplasms—primary tumours of the bony spine and adjacent soft tissues. *Lancet Oncol*. 2007;8(2):137–47.
4. Beer SJ, Menezes AH. Primary tumors of the spine in children: Natural history, management and long-term follow-up. *Spine (Phila Pa 1976)*. 1997;22(6):649–58.
5. Thakur N a, Daniels AH, Schiller J, et al. Benign tumors of the spine. *J Am Acad Orthop Surg*. 2012;20(11):715–24.
6. Harrop JS, Schmidt MH, Boriani S, Shaffrey CI. Aggressive 'benign' primary spine neoplasms. Osteoblastoma, aneurysmal bone cyst and giant cell tumour. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S39–S47.
7. Fisher CG, Andersson GBJ, Weinstein JN. Spine focus issue. Summary of management recommendations in spine oncology. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S2–S6.
8. Fisher CG, Keynan O, Ondra S, Gokaslan Z. Introduction to focus issue in spine oncology: the synthesis of evidence and expert opinion for best practice recommendation. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S21–S25.
9. Choi D, Crockard A, Bunker C, et al. Review of metastatic spine tumour classification and indications for surgery: the consensus statement of the Global Spine Tumour Study Group. *Eur Spine J*. 2010;19(2):215–22.
10. Boriani S, Bandiera S, Donthineni R, et al. Morbidity of en bloc resections in the spine. *Eur Spine J*. 2010;19(2): 231–41.
11. Kelley SP, Ashford RU, Rao AS, Dickson RA. Primary bone tumours of the spine: a 42-year survey from the Leeds Regional Bone Tumour Registry. *Eur Spine J*. 2007;16(3):405–409.
12. Enneking WF, Spanier SS, Goodman MA. A system for the surgical staging of musculoskeletal sarcoma. *Clin Orthop Relat Res*. 1980;153:106–20.
13. Boriani S, Weinstein JN, Biagini R. Primary bone tumors of the spine: Terminology and surgical staging. *Spine (Phila Pa 1976)*. 1997;22(9):1036–44.
14. Tomita K, Kawahara N, Baba H, Tsuchiya H, Fujita T, Toribatake Y. Total en bloc spondylectomy: a new surgical technique for primary malignant tumors. *Spine (Phila Pa 1976)*. 1997;22(3):324–33.
5. Gore L, Greffe BS, Rothenberg SS, Erickson M, Schreiber DP, Handler MH. Long-term survival after intralesional resection and multi-modal therapy of thoracic spine osteosarcoma. *Med Pediatr Oncol*. 2003; (6):400–402.
16. Boriani S, Saravanja D, Yamada Y, Varga PP, Biagini R, Fisher CG. Challenges of local recurrence and cure in low grade malignant tumors of the spine. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S48–S57.
17. Sciubba DM, Okuno SH, Dekutoski MB, Gokaslan ZL. Ewing and osteogenic sarcoma: evidence for multidisciplinary management. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S58–S68.
18. Yamazaki T, McLoughlin GS, Patel S, Rhines LD, Fournay DR. Feasibility and safety of en bloc resection for primary spine tumors: a systematic review by the Spine Oncology Study Group. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S31–S38.
19. Kim HJ, McLawhorn AS, Goldstein MJ, Boland PJ. Malignant osseous tumors of the pediatric spine. *J Am Acad Orthop Surg*. 2012;20:646–56.
20. Bremnes RM, Hauge HN, Sagsvee R. Radiotherapy in the Treatment of Symptomatic Vertebral Hemangiomas: Technical Case Report. *Neurosurgery*. 1996;39(5):1054–58.

Outcome of Conservative Management of Traumatic Acute Subdural Haematoma in Relation to Underlying Brain Injury

Sanat Kumar Saha¹, Mohammad Abu Sayed², Joynul Islam³, Shafiul Alam⁴, ATM Asadullah⁵, Misbah Uddin Ahamed⁶, Shamsul Islam Khan⁷

Abstract

Background: Acute subdural haematoma is haematoma within the dura & arachnoid mater presenting within 72 hours of injury. It is caused by high speed impact that accelerates the brain relative to the fixed dural structure tearing the bridging veins that traverse between the cortical surface & venous sinuses. Small Acute Subdural haematoma less than 5 mm thick on axial CT images, without sufficient mass effect to cause shifting of midline less than 5 mm can be managed conservatively.

Objective: The objective of this study is to assess the morbidity & mortality following conservative management of traumatic acute subdural haematoma.

Methodology: A total 40 patients of Traumatic acute subdural haematoma were selected from Neurosurgery Department, Dhaka Medical College Hospital, Dhaka for this study. All the cases were diagnosed with History, Clinical examination & radiologically by non-contrast CT scan of the Brain. The epidemiological data were recorded and categorized in groups based on some medical and physical features. Then all the categorized data were analyzed using Computer based software SPSS program.

Result: 40 patients were managed conservatively. Among them 30 were not associated with underlying brain injury and 10 were associated with underlying brain injury. In our series mortality of associated underlying brain injury group is 50% but mortality of without associated underlying brain injury is 6.6% and morbidity of associated underlying brain injury group was 40% but disability of without associated brain injury group was 13.33%.

Conclusion: Outcome of conservative management of Traumatic Acute Subdural Haematoma is directly related with underlying brain injury. That is, outcome is worse when there is associated underlying brain injury.

Key Words: Acute Subdural Haematoma, conservative treatment, Morbidity, Mortality.

Bang. J Neurosurgery 2016; 5(2): 37-40

Introduction:

Head injury is the most common cause of death in the world. Acute subdural haematoma is much more common which approximately 30% of all severe head

injuries¹. In our country there is no such epidemiological statistics but report of the Bangladesh Bureau of Statistics, 2001 identifies RTA with head injury as one of the major causes of death⁶. According to the records of two main neurosurgical centers of Bangladesh DMCH & BSMMU 12000-15000 head injury patients get admission every year & the number is increasing day by day. Acute Subdural Haematoma caused by high speed impact that accelerates the brain relative to the fixed dural structure tearing the bridging veins that traverse between the cortical surface & venous sinuses. It can occur due to brain laceration where it is associated with intracerebral contusion & haemorrhage. Acute Subdural Haematoma can result from injury to surface of brain with bleeding from cortical vessels into subdural space. In a general series mortality is around 60% in acute subdural haematoma but can be lowered by very rapid surgical intervention & aggressive medical management. Seelig et al, observed a marked decrease in mortality & morbidity rates in those patients with acute subdural haematoma undergoing craniotomy with evacuation

1. Assistant Professor, Department of Neurosurgery, Sir Salimullah Medical College & Mitford Hospital, Dhaka, Bangladesh.
2. Associate Professor, Department of Neurosurgery, Sir Salimullah Medical College & Mitford Hospital, Dhaka, Bangladesh.
3. Associate Professor, Department of Clinical Neurosurgery, National Institute of Neurosciences & Hospital, Dhaka,
4. Associate Professor, Department of Gamma knife, National Institute of Neurosciences & Hospital, Dhaka, Bangladesh.
5. Associate Professor, Department of Clinical Neurosurgery, National Institute of Neurosciences & Hospital, Dhaka, Bangladesh.
6. Assistant Professor, Department of Clinical Neurosurgery, National Institute of Neurosciences & Hospital, Dhaka, Bangladesh.
7. Assistant Registrar, Department of Clinical Neurosurgery, National Institute of Neurosciences & Hospital, Dhaka, Bangladesh.

Address of Correspondence: Dr. Sanat Kumar Saha, Assistant Professor, Dept. of Neurosurgery, Sir Salimullah Medical College & Mitford Hospital, Dhaka, Bangladesh. Mobile: 01712113393, Email: sanatsaha2005@yahoo.com

of haematoma within 4 hours of injury; 30% of the patient died & 65% had functional recovery². However when surgery was delayed for more than 4 hours post injury, the mortality rate increase to 85% & only 7% of the patients had functional recovery^{11,13}. Haselberger et al, documented that when the interval between the injury & operation exceeded 2 hours, the mortality rise from 47 to 80%³. According to Henry H. Schmidek, MD, FACS, 2000, acute subdural haematoma form against a broad spectrum of primary brain injury of diffuse axonal type⁴. According to Tom Scaletta et al January 2005, all patients of acute subdural haematoma with GCS score less than 12 should be managed in ICU (Intensive Care Unit) with endotracheal intubation⁵. But in our series it was not possible to manage all patients GCS less than 12 in ICU due to limited ICU facilities. We have managed in ICU, all patients GCS less than 9, with severe respiratory distress & falling blood O₂ level with endotracheal intubations. Conservative management includes anticonvulsant, diuretics, maintain fluid & electrolyte balance, antibiotics and other supportive treatment were not taken in ICU in our present study^{7,8,9}. This study tried to find out a plan for management of these patients to reduce mortality & morbidity within an acceptable limit.

Methodology

This is a prospective study. A total 40 patients of Traumatic Acute Subdural Haematoma were selected randomly from Neurosurgery Department, Dhaka Medical College Hospital, Dhaka from July 2005 to June 2007. All the cases were diagnosed with History, Clinical examination & Radiologically by non contrast CT scan of the Brain. The epidemiological data were recorded. Functional recovery associated morbidity & mortality were assessed & recorded in every case as per Glasgow outcome scale (GOS). The collected data was edited, compiled & statistical analysis was done. The data were presented in tables & graphs. In our series 40 patients were managed conservatively. 35 of them was radiologically indicated for conservative management & rest 5 was managed conservatively either due to refusal of surgery or patient was not physically fit for surgical management. On the basis of type of conservatively management of patients some Criteria were followed such as in CT scan of Brain shows maximum thickness of acute subdural haematoma is less than 5 mm, little or no

midline shifting (Fig. 1), refusal of surgical management by the attendants, patient is not physically fit for surgery.




Fig-1: CT Scan of the brain axial view showing acute subdural haematoma (Right) less than 5 mm without midline shifting.

Result:

In our study highest number of patient were in the most active period of life, 70% patients that is 28 patients out of 40 patients were found between 21 to 40 years of age. Below 20 years of age 5 cases (12.5%) were found & 7 cases (17.5%) were found above 40 years of age (Table I).

Table-I
Age distribution in study group (n = 40)

Age in year	No of cases	%
<20	5	12.5
21-30	13	32.5
31-40	15	37.5
>40	7	17.5
Total	40	100

Out of 40 patients in our study 32 cases (80%) were male & only 20% (8 cases) were female. The number of occurrence of brain injury was approximately four times higher in male than in female (Table-2).

Table-II
Sex distribution in study group (n=40)

Sex	No of cases	%
Male	32	80
Female	8	20
Total	40	100

Highest numbers of head injury were found due to road traffic accident which was 50% (20 cases). Next to road traffic accident 2nd highest causes of head injury was fall from height which was 25% (10 cases) & 7 cases were due to assault (17.5%).

Table-III
Incidence of underlying brain injury according to mode of injury

Mode of injury	ASDH without Brain Injury		ASDH with Brain Injury	
	No.	%	No.	%
RTA	16	53.33	4	40
Fall from height	7	23.33	3	30
Assault	5	16.66	2	20
Others	2	6.66	1	10
Total	30		10	

CT scan of Brain done in all 40 (100%) cases, as CT scan in the investigation of choice for its high

sensitivity & specificity. Plain X-ray skull (B/V) done in 32 cases (80%). Random Blood Sugar, S. Electrolytes & Blood gases analysis was done in some selective cases, according to need but not included in this study.

Out of 40, 10 patient of associated underlying brain injury, highest incidence of brain injury was underlying brain contusion which was 60% cases (6 patients). 3 patients had associated intracerebral clot which was about 30% & associated infarction was found in 1 patient (10%) (Table-4).

Table-IV
Type of underlying brain injury in study group (n = 10)

Type of injury	No of patients	%
Brain contusion	6	60
Intracerebral clot	3	30
Infarction	1	10
Total	10	100

So, in conservatively managed group of patient, prognosis is directly related with underlying brain injury. That is, prognosis is worse when there is associated underlying brain injury.

Table-V
Follow up of the conservatively managed patients as Glasgow Outcome Scale (n = 40)

Total no of patients		Death		Vegetative		Severe disability		Moderate disability		Good recovery	
		No.	%	No	%	No	%	No	%	No	%
ASDH Without brain injury	30	2	6.6	1	3.3	0	0	3	10	24	80
ASDH With brain injury	10	5	50	1	10	1	10	2	20	1	10
Total	40	7		2		1		5		25	

Table-VI
Outcome of conservative treatment (n = 40)

For analysis of data vegetative, severe disability & moderate disability is counted as single group as disability.

ASDH	Death	Disability	Good recovery	Number
Without brain injury	2	4	24	30
With brain injury	5	4	1	10
Total	7	8	25	40

Chi-Square Tests:

	χ^2	df	p value
Pearson Chi-square	22.14	2	p<.001

Discussion:

Acute subdural haematoma is a neurosurgical emergency worrying the surgeon with its malignant behavior. High mortality & morbidity of patients following acute subdural haematoma has led to search for better treatment modalities¹⁰. In advanced world despite of rapid transportation, emergency medical services, and improved radio imaging acute subdural haematoma continue to contribute significantly poor outcome is severely head injured patients.

Outcome of the conservative treatment was evaluated on the basis of Glasgow Outcome Scale (GOS)¹². According to Rengachary S.S 1994, the overall mortality rate of patients with a treated ASDH is roughly 50%, but in our series overall mortality is 17.5% (7 out of 40). In our series mortality of associated underlying brain injury group is 50% but mortality of without associated underlying brain injury is 6.6%. So mortality of associated underlying brain injury is much more higher. In our series overall disability was 20%. Disability of associated underlying brain injury group was 40% but disability of without associated brain injury group was 13.33%, So, disability of associated brain injury is also higher. On the other hand overall good recovery was 62.5%, in our series. Good recovery of without associated brain injury was 80% but with underlying brain injury 10%, so good recovery of associated underlying brain injury is worse than without underlying brain injury.

Morbidity & mortality following acute subdural haematoma is still much more higher. Mortality can be lowered by rapid surgical intervention & intensive medical management. According to Kotwica 1993, an acute subdural haematoma commonly is associated with extensive primary brain injury¹¹. This diffuse parenchymal injury correlate strongly with the outcome of the patient. According to Henry H. Schmidek 2000, Rates of mortality & morbidity after an acute subdural haematoma are the highest of all traumatic mass lesions⁴. The poor outcome results largely from associated parenchymal injuries & subsequent intracranial hypertension. Approximately 50% of patients have associated lesions, but in our

series associated underlying brain injury was 25% (10, out of 40). So, according to different authors underlying brain injury is more common in acute subdural haematoma & morbidity, mortality is higher in underlying brain injury.

Conclusion:

Outcome of Traumatic acute subdural haematoma depends on underlying brain injury. Our study revealed that both mortality and morbidity are higher in traumatic acute sub dural haematoma with underlying brain injury than without underlying brain injury after conservative management.

Reference:

1. Ramamurthy B, Tandon P.N. Extradural haematomas: Textbook of Neurosurgery: 2nd ed; Churchill Livingstone; 1996; vol (1).
2. Seelig JM, Becker DP, Miller JD, Miller JD, et al. Traumatic acute subdural haematoma. Major mortality reduction in comatose patients treated within four hours. N Engl J Med 1981;304:1511-1512.
3. Haselsberger K, Pucher R, Auer LM, Prognosis after acute subdural or epidural haemorrhage. Acta Neurochir (Wien) 1988; 90: 111-116
4. Henry H. Schmidek, MD, FACS, Operative Neurosurgical Techniques Fourth Edition Volume 1-2 Cilt, 2000
5. Scaletta, Tom. "Subdural Hematoma," *eMedicine Medical Library*, January, 2005.
6. Ahmed R, Hossain MA: Craniocerebral Trauma; Epidemiology & Aetiology. Bangladesh Journal of neuroscience 1989, 5:66-75.
7. Becker DP, Miller JD, Young HF, et al. Diagnosis and treatment of head injury in adults. In Youmans LR (ed): Neurological Surgery. Philadelphia; Saunders, 1982, pp 1938-2083.
8. Greenberg M.S: Hand of neurosurgery 4th ed; 1997, Inc vol. 11, pp 729-31.
9. Lindsay KW : Neurology & Neurosurgery Illustrated, 3rd ed, 1997, p 215.
10. Rengachary SS: Principles of Neurosurgery 1994, 4th ed, p-1.2, 1.22, 16.5, 19.4, 19.5.
11. Seelig JM, Becker DP, Miller JD, Miller JD, et al. Traumatic acute subdural haematoma. Major mortality reduction in comatose patients treated within four hours. N Engl J Med 1981;304:1511-1512.
12. Servadei F: Prognostic factors in severely head injured adult patients with acute subdural haematomas. Acta Neurochir (Wien) 1997; 139(4): 279-85.
13. Wilberger JE Jr, Harris M, Diamond DL: Acute subdural hematoma: morbidity and mortality related to timing of operative intervention. J Trauma 1990 Jun; 30(6): 733-6.

Signs and Symptoms of Early VP Shunt Infection in a Tertiary Hospital

Kaisar Haroon¹, Tania Taher², Mokhlasur Rahman³, Kanak Kanti Barua⁴

Abstract:

Background: VP shunt infection is a major concern for the treating surgeon. Shunt infection causes shunt malfunction and is source of much morbidity for the patient.

Introduction: Hydrocephalus was diagnosed in patients clinically by history, symptoms and signs. This was confirmed by CT and/or MRI scans and was treated with ventriculoperitoneal shunt system. In this study we have analyzed signs and symptoms of early shunt infection. That will help in early detection of shunt infection and faster management.

Methods: This is an observational study. The study was done from September 2008 up to January 2010 in BSMMU. Thirty patients were operated on. They were followed up to 15 days after surgery. They were evaluated with presence of non-remitting fever in the absence of other causes, indication of infection along the shunt tract or incision site and signs of meningeal irritation or peritonitis.

Result: Seven patients of the thirty patients treated with ventriculoperitoneal shunt surgery had VP shunt infection. That is 23.33% of patients. This is more in the young and elderly age group. It was also more at the lower socioeconomic age group. Most patients had fever, headache and neck rigidity.

Key Words: Hydrocephalus, ventriculomegaly, shunt infection, headache, papilledema, meningitis, ventriculitis, ventriculoperitoneal shunt.

Bang. J Neurosurgery 2016; 5(2): 41-43

Introduction:

Hydrocephalus is defined as an abnormal enlargement of the ventricles due to an excessive accumulation of CSF from a disturbance of its flow, absorption or uncommonly, secretion¹. Ventriculoperitoneal shunt catheter placement is a relatively common neurosurgical procedure, performed for the treatment of hydrocephalus as well as for associated conditions in which the natural flow of cerebrospinal fluid is obstructed². According to McGirt et al. the signs of shunt infection are meningitis, ventriculitis, CSF leak from wound, skin infection—anywhere along the tract, peritonitis, seizure disorder³.

Trojanowski had commented that between 5 and 15% of the devices become infected, of which more than a half within the first month after surgery⁴. Ahmed et al. have studied complications on fifty patients. Of these 6 (12%) patients had shunt infection⁵. Drake and Iantosca reported that the incidence of shunt

infection is approximately 5%- 10%⁶. Bokhary and Kamal incidence rate is nearly 10%⁷. Kinasha et al. in their study in Tanzania reported the shunt infection rate about 24.6%⁸. McGirt et al. had concluded from their observation that shunt infection occurred in 11% of cases (92 of 820)³. Leach and Kerr had reported that shunt infection should be within 1% and 7% of shunt insertions⁹.

Materials and Method:

This is an observational study. This was conducted from 01-07-2008 to 17-05-2010. All patients were admitted in the BSMMU Hospital for ventriculoperitoneal shunt surgery. Patients who met the inclusion criteria from all age groups were included in the study. Number of patients were thirty ($n=30$).

Shunt infection was defined if at least one of the following criteria was present:

1. Presence of non-remitting fever in the absence of other causes
 2. Indication of infection along the shunt tract or incision site
 3. Signs of meningeal irritation or peritonitis.
- Shunt system used: In all patients we used Chhabra medium pressure ventriculoperitoneal shunt system (Surgiview Inc.)

1. Assistant Professor, Department of Neurosurgery, NINSH
2. Assistant professor, Department of Paediatrics, MH Shamorita Medical College Hospital
3. Assistant Professor, Department of Neurosurgery, NINSH
4. Professor and Head, Department of Neurosurgery, BSMMU
Address of Correspondence: Dr. Kaisar Haroon, Assistant Professor, Department of Neurosurgery, NINS. Email: kaisar298@gmail.com

- Inclusion criteria:

All patients with ventriculomegaly in imaging and signs of hydrocephalus undergoing ventriculoperitoneal shunt surgery were selected.

- Exclusion criteria:

- o Patients with preoperative fever or any other focus of infection
- o Patients with other medical problems e.g., diabetes mellitus, valvular heart disease, renal impairment and ruptured meningocele.
- o Patients who had previous CSF diversion surgery (e.g. ETV, EVD)

Results:

After proper collection and processing of samples, the data is analyzed. The results and observations from the analysis were presented in tables and figures. Data were expressed in numbers or percentage as appropriate.

Table-I
Distribution of the patients by age

Age in years	Frequency		Total
	Shunt infection	No shunt infection	
0-2	2	6	8
2-12	1	5	6
12-30	1	3	4
30-50	1	4	5
>50	2	5	7
Total	7	23	30

From this table we can see that shut infection is more in the below 12 years and more than 50 years.

Table-II
Distribution of the patients by sex

Variables	Male	Female	Total
Shunt infection	4	3	7
No shunt infection	16	7	23
Total	20	10	30

In this table we can see that 4 male patients and 3 female patients had shunt infection.

Table-III
Distribution of the patients by post-operative morbidity

Post operative complications	Total	Percentage
Persisting Fever	07	23.33%
Surgical site infection	05	16.66%
Shunt tract infection	05	16.66%
Skin necrosis	1	3.33%
Headache	6	20%
Vomiting	6	20%
Neck rigidity	1	3.33%
Shunt block	1	3.33%
Peritonitis	2	6.66%

From the above table it is seen that seven patients (23.33%) developed persisting fever. SSI occurred in 5 patients (16.66%). Four patients (13.33%) had shunt tract infection and only one patient had skin necrosis. Headache and vomiting was present in 6 patients (20%), neck rigidity was present in one patient and shunt block occurred in one patient. Two patients (6.66) developed signs of peritonitis.

Table-IV
Distribution of the patients by post-operative vomiting, headache and non-remitting fever

Age in years	Vomiting	Headache	Non-remitting fever
0-2	3	2	3
2-12	1	1	1
12-30	0	0	0
30-50	1	2	2
>50	1	1	1
Total	6	6	7

From the above table it is seen that total 6 patients had vomiting in the post operative period. 3 patients were under the age of 2 years. In this table 6 patients also headache. Most patients were above 12 years of age (66.67%). It also shows that total 7 patients had non-remitting fever.

Table-V
Distribution of the patients by surgical site infection (n=6)

Age in years	SSI	Neck rigidity	Peritonitis
0-2	3	0	1
2-12	2	0	1
12-30	0	1	0
30-50	1	0	0
>50	0	0	0
Total	6	1	2

From the above table we can see that six patients had surgical site infection, one patient had neck rigidity and two patients had signs of peritonitis.




Fig.-1: Distribution of the patients by shunt infection in all patients (n=30).

From this above chart it is seen that shunt infection occurred in seven patients out of thirty patients. This was about 23.33%.

Discussion:

Choux et al. in 1992 had defined post-operative shunt infection as an infection confirmed within six months of operation and was diagnosed if there was inflammation along the length of shunt, wound discharge or wound dehiscence. Signs of meningitis, ventriculitis or non-focal systemic indication of infection were also investigated for shunt infection¹⁰. In this study, 5 patients (16.66%) had these signs. Signs of meningitis was present in one patient. Five patients had headache and vomiting. Kontny et al. had mentioned that the main sign and symptoms were fever, shunt malfunction and meningeal irritation, and with VP-shunts only abdominal pain¹¹. Seven patients had fever in this study following shunt insertion. Pople in 2002 had suggested the following as most important clinical features of shunt infection: general malaise, pyrexia, headaches, vomiting, neck stiffness, abdominal tenderness or distension, recurrent lower end shunt obstruction, occasionally pain and erythema around the shunt¹². In our study seven patients (23.33%) had pyrexia, five patients (16.66%) had shunt malfunction and peritonitis in two patients

(6.66%). The distribution of the shunt infection is congruent with the patients of the other studies.

Conclusion:

Ventriculoperitoneal shunt surgery is a common procedure. Shunt infection can reduce the success of this surgery. If we can diagnose the shunt infection early, then we can reduce morbidity as well as mortality of these patients. Therefore, we have to diagnose shunt infection very quickly and accurately and take necessary measures for benefit of the patient.

References:

1. Kaye A. Raised intracranial pressure and hydrocephalus. In: Kaye A, ed. *Essential Neurosurgery*. 3rd ed. Massachusetts, USA: Blackwell Publishing Ltd 2005:31-36.
2. Claus E, Duncan C. Shunt infection. In: Winn H, ed. *Youman's Neurological surgery*. Philadelphia: Saunders 2004:3419-24.
3. McGirt MJ, Zaas A, Fuchs HE, et al. Risk factors for pediatric ventriculoperitoneal shunt infection and predictors of infectious pathogens. *Clinical infectious diseases* 2003;36(7):858-62.
4. Trojanowski T. Hydrocephalus in adults. In: Sindou M, ed. *Practical Handbook of Neurosurgery*, vol-1, pp Germany: Springer, 2009:429.
5. Ahmed A, Sandlas G, Kothari P, et al. Outcome analysis of shunt surgery in hydrocephalus. *Journal of Indian Association of Pediatric Surgeons* 2009;14(3):98.
6. Drake J, Iantosca M. Cerebrospinal fluid shunting and management of paediatric hydrocephalus. In: Schimidek H, Roberts D, eds. *Schimidek and Sweet Operative Neurosurgical Techniques: indications, methods and results*. Philadelphia: Saunders Elsevier 2006:488-99.
7. Bokhary Aly M, Kamal H. Ventriculo-peritoneal shunt infections in infants and children. *Libyan Journal of Medicine* 2008;3(1):20-22.
8. Kinasha A, Kahamba J, Semali I. Complications of ventriculoperitoneal shunts in children in Dar es Salaam. *East and Central African Journal of Surgery* 2005;10(2): 55-59.
9. Leach J, Kerr R. Elective Neurosurgery. In: Williams, Bulstrode, O'Connell, eds. *Bailey and Love's Short Practice of Surgery*. 25th ed. London: Hodder Arnold 2008:625-28.
10. Choux M, Genitori L, Lang D, et al. Shunt implantation: reducing the incidence of shunt infection. *Journal of neurosurgery* 1992;77(6):875-80.
11. Kontny U, Höfling B, Gutjahr P, et al. CSF shunt infections in children. *Infection* 1993;21(2):89-92.
12. Pople IK. Hydrocephalus and shunts: what the neurologist should know. *Journal of Neurology, Neurosurgery & Psychiatry* 2002;73(suppl 1):i17-i22.

Assessment of Serum Homocysteine Level in Epileptic Patients on Carbamazepine Treatment

Md. Abu Hanif¹, M A Hannan², M Iqbal Arslan³, SK Mahbub Alam⁴, Mohammad Hossain⁵, Md. Aynul Hoque⁶, Niloy Ranjan Roy⁷

Abstract:

Background: This comparative cross sectional study was carried out to evaluate the level of serum homocysteine with carbamazepine treatment in epileptic patients.

Materials and Methods: This study was conducted in the Department of Neurology, BSMMU, Dhaka, from March 2013 to December 2015 for the duration of two years and nine months period. Patients, who were diagnosed as epileptic and receiving Carbamazepine at least for six months and no other anti epileptic drugs (AEDs), were considered as case group and Patients other than epilepsy attending Neurology outpatient department (NOPD) were considered as control group.

Results: A total of 104 patients were recruited as study population. Of them 52 patients grouped into case and the rest 52 patients in control. The mean (SD) value of serum homocysteine in epileptic patients using carbamazepine as anti-epileptic drugs found 16.33 (5.34) $\mu\text{mol/L}$ which was highly statistically significant difference than control group measuring 8.77 (3.75) $\mu\text{mol/L}$. Considering the normal level 5- 15 $\mu\text{mol/L}$, in epileptic patients with carbamazepine therapy, increased level was found in 53.8% patients. On the other hand, in control group, 13.5% had increased value of serum homocysteine.

Conclusion: Carbamazepine therapy which is usually given to epileptic patients may interfere with metabolic pathways of homocysteine (Hcy) as well as may lead to an alteration of its serum levels.

Bang. J Neurosurgery 2016; 5(2): 44-47

Introduction:

Many developing countries such as Bangladesh, India are densely populated. Like tuberculosis, malaria epilepsy is also a burden for our country. Anti- epileptic drugs are commonly used to treat epilepsy.

Commonly used first-generation antiepileptic drugs- carbamazepine, sodium valproate, phenytoin and phenobarbitone. Second-generation antiepileptic drugs are- lamotrigine, gabapentin, topiramate etc. Oxcarbazepine and levetiracetam- available but are highly expensive

However consumption of carbamazepine causes significant elevation of serum homocysteine level¹.

Elevated serum homocysteine (Hcy) is responsible for ischaemic heart disease, deep vein thrombosis, pulmonary embolism and stroke². Near about 10-40% of epileptic patients develop hyperhomocysteinaemia³. High plasma homocysteine is an independent risk factor for cardiovascular disease and thrombosis⁴. Anti-epileptic drugs raise serum homocysteine (Hcy) by reducing blood folate levels⁵.

Atherosclerotic vascular diseases are associated with well-known risk factors such as systemic arterial hypertension, diabetes mellitus, smoking and obesity, but in the last decade other emerging risk markers have been identified, one of them is being Hcy. Atherosclerosis is a chronic inflammatory disease of the blood vessel walls, in which deposits of lipids, cholesterol, calcium and other substances build up in the endothelial layer of the arteries. Many studies have identified moderate elevation of serum Hcy as an independent risk factor for atherosclerotic vascular disease⁶.

Lowering of serum folate and elevation of Hcy concentrations in blood are associated with poor cognition in general population. High levels of Hcy result in neurotoxic and vasotoxic effects in dementia and Alzheimer's disease, suggesting that Hcy is a direct marker of early cognitive deficit⁷.

1. OSD, Medical officer, DGHS, Attachment- NIDCH, Dhaka,
2. Professor & Chairman, Neurology Bangabandhu Sheikh Mujib Medical University(BSMMU),
3. Professor & Chairman, Department of Biochemistry, Dean, Basic Medical Science Faculty and Syndicate Member(BSMMU),
4. Associate Professor, Neurology, BSMMU, Dhaka
5. Associate Professor, Neurosurgery, BSMMU, Dhaka
6. OSD, DGHS,
7. Assistant Registrar, Shaheed Tajuddin Ahmed Medical College.

Address of Correspondence: Dr. Md. Abu Hanif, OSD, Medical officer, DGHS, Attachment- NIDCH, Dhaka,

The pathogenic role of high serum Hcy is in vascular dementia is yet no clear. The levels of serum Hcy, vitamin B12 and plasma folate were studied in Alzheimer's disease and vascular dementia, and it was found that Hcy was increased, while vitamin B12 and folic acid were decreased in both of these conditions. This suggests that supplementation of vitamin B12 and folate may be beneficial in vascular dementia and Alzheimer's disease ⁸.

Materials and Methods:

This study was conducted in the Department of Neurology, BSMMU, Dhaka, from March 2013 to December 2015 for the duration of two years and nine months period. Cases were selected by simple random sampling. Age ranges of the patients were 14 to 60 years. Because baseline serum Hcy level have similarity in these age group. Moreover we have collected cases from epilepsy clinic of adult neurology. Epileptic patients with other systemic diseases such as: Chronic kidney disease, Hypothyroidism, chronic liver disease, Leukemia, Inflammatory bowel disease, psoriasis and homocysteinuria were excluded from this study⁹. Patient with Diabètes Mellitus, Smoking, Pregnancy, age less than 14 years and greater than 60 years, taking Methotrexate, Isoniazide or other AEDs, folic acid supplement, receiving vit-B6, Vit- B12 or on folate antagonists were also excluded from this study

Data were collected by a pre-designed proforma. Detailed history and clinical examination were carried out and structured questionnaire were used to collect the necessary information. Similar numbers of controls were also selected. Relevant base line investigations were done which included Complete blood count with peripheral blood film where applicable, Urine R/M/E, Serum creatinine, ALT, Thyroid function tests in suspected hypothyroidism.

Serum Hcy assay was done from venous blood of both cases and control in the Department of Biochemistry, BSMMU on the day of sample collection. The ARCHITECT Homocysteine assay, a chemiluminescent micro particle immunoassay (CMIA) technology was used to measure homocysteine value. Statistical analysis of data was carried out with appropriate techniques and systems.

Results:

A total of 104 patients were recruited as study population. Of them 52 patients grouped into case and the rest 52 patients in control.

Table-I

Distribution of patients (Case group) according to epilepsy type (n= 52)

Type of Epilepsy	Frequency (n)	Percentage
Simple partial	17	32.7
Complex partial	5	9.6
Primary generalized	26	50.0
Secondary generalized	4	7.7

This table I shows distribution of Case group patients according to epilepsy type. Here, half of the patients, 50% had primary generalized epilepsy which was followed by simple partial type 32.7%. Complex partial was accounted 9.6% whereas secondary generalized 7.7%.

Table-II

Distribution of case group patients according to duration of diagnosis and treatment (n=52).

Duration (Years)	Mean	SD
Duration of diagnosis	4.29	4.51
Duration of treatment	4.05	3.59

This table II shows distribution of case group patients according to duration of diagnosis and treatment. The mean (SD) value of duration of diagnosis was 4.29 (4.51) years while duration of treatment was 4.05 (3.59) years.

Table-III

Distribution of case group patients according to Carbamazepine dose level (n=52)

Carbamazepine dose (per day)	Frequency (n)	Percentage
200 mg	2	3.8
400 mg	18	34.6
800 mg	30	57.7
1000 mg	2	3.8

This table III shows distribution of case group patients according to Carbamazepine dose level. Out of all patients, 57.7% patients were taken 800 mg carbamazepine daily. The second highest portion of patients 34.6% was taken 400 mg of carbamazepine daily.

Table-IV

Distribution of patients according to serum level of homocysteine (N=104)

Type of Patients	Serum Homocysteine (micro mol/L)		p-value*
	Mean	SD	
Case	16.33	5.34	<0.0001 ^s
Control	8.77	3.75	

S=significant

* P-value derived from independent sample t test

Table IV: shows distribution of patients according to serum homocysteine level. The mean (SD) value of serum homocysteine in epileptic patients using carbamazepine as anti-epileptic drugs found highly increased than control group [16.33 (5.34) vs. 8.77 (3.75)]

Table-V

Distribution of patients according to range of serum Homocysteine level (N=104).

Type of Patients	Serum Homocysteine (¼mol/L)	
	Minimum	Maximum
Case	8.45	30.28
Control	5.30	19.00

The table II shows distribution of patients by range of serum homocysteine level. Here, in case group, the range was 8.45 to 30.28 ¼mol/L and that of 5.30 to 19.00 ¼mol/L in control group

Discussion:

Carbamazepine (CBZ) is a potent antiepileptic drug. It is commonly used in partial onset and secondary generalized seizures. It has also good efficacy in the treatment of primary generalized tonic-clonic seizures. Some studies showed in epileptic patients receiving CBZ, there is significant elevation of serum homocysteine levels¹⁰. No previous study has done in Bangladeshi subjects. It is one of the foremost studies of epilepsy patients taking carbamazepine therapy to consider their serum homocysteine level in Bangladesh context. We compared our study findings with result of some other published articles elsewhere in the world.

According to gender distribution, no statistically significant difference was observed between two groups. Out of all patients, 67.3% were male and 32.7%

female in carbamazepine treated patients. In control group patients, 63.5% was male and 36.5% female.

In analysis of epilepsy type, half of the patients, 50% had primary generalized epilepsy which was followed by simple partial type 32.7%. Complex partial was accounted 9.6% whereas secondary generalized 7.7%.

The mean (SD) value of duration of diagnosis was 4.29 (4.51) years while duration of treatment was 4.05 (3.59) years. Out of all patients, 57.7% patients were taken 800 mg carbamazepine daily. The second highest portions of patients 34.6% were taken 400 mg of carbamazepine daily.

The present study showed that epileptic patients receiving carbamazepine as AEDs have increased serum levels of homocysteine. The mean value of serum homocysteine in epileptic patients taking carbamazepine as anti-epileptic drugs found 16.33 (5.34) ¼mol/L which was highly statistically significant difference than control group measuring 8.77 (3.75) ¼mol/L. However, in case group the highest number of 13 patients had their value in 15 – 20 ¼mol/L and highest value 30 ¼mol/L had only in 1 patient. At the same time, in control group highest number of 21 patients had their value in 5 – 10 ¼mol/L and highest value near about 20 ¼mol/L had in 2 patients. In case group, the range of serum homocysteine was found 8.45 to 30.28 ¼mol/L and that of 5.30 to 19.00 ¼mol/L in control group.

Our results are in consistent with the previous reports on homocysteine concentrations in patients with epilepsy taking AED specially CBZ. Paknahad et al, also recorded mean of serum Hcy concentration in epileptic patients receiving CBZ was significantly higher compared to that in the controls [13.66 (0.95) vs. 12.97 (0.46) ¼mol/L, $p = 0.04$]⁴. The similar result was obtained if patients with only CBZ monotherapy were included, Minitzer et al, found 11.1 (4.2) ¼mol/L¹¹. Epilepsy patients switching from the enzyme inducing AEDs, phenytoin or CBZ to the non inducing AEDs drugs, LTG or levetiracetam results in significant declines in serum homocysteine level.

Conclusion:

In conclusion, our data demonstrate that serum homocysteine level significantly increased in carbamazepine treated epileptic patients. Therefore,

it should be monitored the level of homocysteine in epileptic patients and managed accordingly.

Reference:

1. Gorjipour F, Asadi Y, Osguei NK, Effatkhan M, Samadikuchaksaraei A. Serum levels of Homocysteine, Folate and Vitamin-B12 in Epileptic patients under Carbamazepine and Sodium valproate treatment ; A Systematic Review and Meta-Analysis. *Iranian Red Crescent Medical Journal*. 2013; 15(3): 249-53.
2. Wald DS, Law M, Morris JK. Homocysteine & cardiovascular disease: Evidence on Causality from a meta-analysis. *BMJ*. 2002; 325:1202.
3. Apeland T, Mansoor MA, Pentieva K, McNulty H, Seljeftot I, Strandjord RE. The effect of B-vitamins on hyperhomocysteinemia in patients on antiepileptic drugs. *Epilepsy Res*. 2002; 51: 237-247.
4. Paknahad Z, Ahmad C, Akbar HZ, Elham S. Effects of Common Anti-epileptic Drugs on the Serum Levels of Homocysteine and Folic Acid. *Int J Prev Med*. 2012;3(Suppl1): 86–90.
5. Kishi T, Fujita N, Eguchi T, Uedak K. Mechanisms for reduction of serum folate by antiepileptic drugs during prolonged therapy. *J Neurol Sci*. 1997; 145: 109.
6. Weiss N, Keller C, Hoffmann U, Loscalzo J. Endothelial dysfunction and atherothrombosis in mild hyperhomocysteinemia. *Vas Med*. 2002; 7: 227-23
7. Ravaglia G, Forti P, Maioli F, Martelli M, Servadi L, Brunetti N, et al. Homocysteine and folate as risk factors for dementia and Alzheimer's disease. *Am J Clin Nutr*. 2005; 82: 636-643.
8. Koseoglu E, Karaman Y. Relations between homocysteine, folate and vitamine B-12 in vascular dementia and Alzheimer's disease. *Clin Biochem*. 2007; 40: 859-863.
9. Harrison's. Internal Medicine' TR Hyperhomocysteinemia. 18th edition. New-York: Mc Graw Hill, 2012; pp 1953
10. Kurul S, Unalp A, Yis U. Homocysteine levels in epileptic children receiving antiepileptic drugs. *J Child Neurol*. 2007; 22: 1389-92.
11. Mintzer S, Skidmore CT, Abidin CS, Morales MC, Chervoneva I, Capuzzi DM et al. Effects of Antiepileptic Drugs on Lipids, Homocysteine, and C-Reactive Protein. *Ann Neurol*. 2009; 65: 448–456.

Case Reports

Tubercular Spinal Epidural Abscess Involving the Lumbo-Sacral Region: A Case Report with Review of Literature

Akhlaque Hossain Khan¹, Ranjit Kumar Chaurasiya², Narendra Shalike³, Mohammed Aminul Hasnat⁴, Bipin Chaurasia⁵, Abhishek Chaturbedi⁶, Asifur Rahman⁷, Rezaul Amin⁸

Abstract:

Tubercular spinal epidural abscess (SEA) without bony involvement is a rare condition but may be devastating and fatal.

This is a report of a rare case of a tubercular spinal epidural abscess in a 60 years old man who presented with paraparesis and Cauda-Equina Syndrome with Fournier's gangrene, but without vertebral bony involvement. MRI of the lumbosacral spine revealed huge collection of thick pus extradurally and posteriorly extending from L2 vertebra to sacral region. The abscess was treated surgically by posterior surgical decompression that involved laminectomy of L2 to L5 vertebra and pus was completely drained. Histopathology demonstrated Mycobacterium Tuberculosis bacteria in the drained pus. Anti-tubercular drugs were started postoperatively. The patient showed progressive improvement in his neurological deficits and started walking without support at 6 months of follow-up after hospital discharge.

Keywords: Epidural; abscess; paralysis; lumbar-sacral; laminectomy; ATT; Ambulation

Bang. J Neurosurgery 2016; 5(2): 48-50

Introduction:

Spinal epidural abscess represents a severe pyogenic infection of the epidural space which may compress neural elements and require urgent surgical intervention to avoid permanent neurological deficits.[1] It is a rare variety reported with an incidence of 0.2 -2 cases per 10000 admissions at

hospital.[2] The incidence of spinal epidural abscess is increasing in developing country which usually occurs secondary to tubercular spondylitis but may rarely develop by haematogenous spread by any foci. Tubercular spondylitis and its resulting complications are common in developing nations.[3] Early diagnosis with advanced imaging techniques, proper surgical intervention and effective coverage of broad spectrum antibiotics have led to gradual decrease in mortality rates but still data show high mortality rates of 5-33% in some series .[4] The purpose of this study was to analyse the clinical manifestations, prognostic indicators, source of infection and final outcomes in patients with SEA.[4]We present a rare case of spinal epidural abscess of tubercular origin involving the lumbar and sacral spine without osseous involvement.

Case Report:

A 60 years old man, farmer by occupation, developed pain, tingling and numbness in his both lower limbs for 2 months which was insidious in onset. After one and half months later, he developed dribbling of urine and slowly over one week, he developed incontinence of urine. During the course of his illness, he had history of intermittent type of fever and weight loss, for which he visited a general practitioner. He was

1. Associate Professor, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.
2. Resident, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.
3. Resident, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.
4. Resident, Department of Anaesthesia, National Institute of Neurosciences & Hospital, Dhaka, Bangladesh.
5. Resident, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.
6. Resident, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.
7. Assistant Professor, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.
8. Assistant Professor, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.

Address of Correspondence: Dr. Akhlaque Hossain Khan, Associate Professor, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.

diagnosed as a case of UTI and was advised some drugs. But the patient condition got worsened over time and got admitted at nearby hospital. A self-retaining urinary catheter with some medicines were advised and discharged after two weeks. After 2 weeks, patient developed Fournier's gangrene with bed sore in left sacral region. Surgical wound debridement and S.P.C was done under spinal block in September 2014 and was put on anti-tubercular regimen but patient later developed drug induced hepatitis. The drug was stopped and referred to higher centre and the patient was admitted in our hospital. MRI of the lumbosacral spine with whole spine screening film was done which showed there was significant compression over thecal sac extending from L2 to sacral region by huge collection of pus extradurally and posteriorly (Figure 1). Chest x-ray showed some patchy infiltrates around the mediastinal-hilar region. Neutrophil was 88%, ESR

was 85 in 1st hour and was diagnosed as a case of epidural abscess from L2 to sacral region with lower limbs paraparesis and incontinence of urine.

Posterior surgical decompression was done by laminectomy of L2 to L5 vertebra. A thick collection of pus was found extradurally and posteriorly which was sticky, yellowish in colour and was evacuated and thorough irrigation done with normal saline mixed with antibiotic Gentamycin. Dural sac was made free and pus was sent for Gram stain, culture and sensitivity, and Acid Fast Bacillus (AFB). Wound was closed over a suction drain.

Pus for AFB was found to be positive. Again, anti-tubercular therapy (ATT) was started following the report and was discharged under supervision. After one week, patient developed ulceration in the genito-perineal region which was treated conservatively. Postoperatively no significant improvement was noted and passive physiotherapy was advised. Gradually,




Fig.-1. Preoperative MRI of Lumbosacral spine with different views showing significant compression over thecal sac extending from L2 to sacral region by huge collection of pus extradurally and posteriorly.

patient regained improvement in power of his lower limbs and at 6 months of follow up, patient was found to have muscle power of 4 out of 5 in both his lower limbs and also control over his bladder function.

Discussion:

Most spinal infections in developed countries are due to pyogenic infections where as non pyogenic organisms are responsible for most spinal infections in developing countries and also in the immunocompromised patients in the developed nations.^{1,3} Mycobacterium Tuberculosis is the most important organisms in developing nations.¹ One of the common cause of death in the world is tuberculosis (TB) which is assumed that two billion people are infected with TB, who can develop TB disease in their course of illness. TB mostly affects children and young adults. The risk factors regarding the development of the disease depends on the patient compliance and the geographical variations because of the different incidence of TB in each country. [5,9] Infection of the spine can pose a wide range of problems for both patient and the surgeon. Tuberculosis demonstrates a varieties of clinical and radiological findings and has a known propensity for dissemination from its primary site; therefore it can mimic a number of disorders. The microbiological diagnosis of spinal infection is very important because despite the surgical treatment, the medical management with ATT drugs is essential postoperatively for resolution of infection. An estimated 20% - 40% involved spinal tuberculosis patients have another focus of infection and due to lack of proper antimicrobial treatment, poor compliance of the patients and proper duration lead to flare of the infection. Despite proper management, the resistance to spinal tuberculosis is increasing.[6] Innumerable patients and lack of risk factors in most of the patient results in diagnosis delay during the process of the disease¹¹.

Cauda equina syndrome is uncommon; accounting for an incidence of 1–5% of spinal pathology in the literature. Acute trauma is a rare cause of this syndrome. The most common presenting features are bladder symptoms, saddle anaesthesia and loss of motor function. Early surgical decompression has been suggested to obtain a satisfactory recovery^{7, 8, 10}. Patient with spinal epidural abscess (SEA) may have unknown focus of infection with pain and tenderness of the spine clue to proper investigation lead to early diagnosis of the disease¹².

Conclusion:

Spinal epidural abscess without bony and pulmonary involvement is uncommon but potentially devastating condition that challenge many clinicians regarding

its proper diagnosis in timely fashion for appropriate management. MRI of the spine is a noninvasive and sophisticated tool for the early diagnosis of SEA due to spinal TB. Early diagnosis and adequate surgical decompression of the SEA compressing the thecal sac and nerve roots along with proper postoperative anti tubercular therapy leads to cure of the patient having good chances of recovery of neurological functions.

Conflict of interest

Authors declare no conflict of interest.

References:

1. Reihnsaus E, Waldbaur H, Seeling W. Spinal epidural abscess: a meta-analysis of 915 patients. *Neurosurgical review*. 2000 Dec 25;23(4):175-204.
2. Agrawal V, Patgoankar PR, Nagariya SP. Tuberculosis of the spine. *J Craniovertebr Junction Spine*. 2010; 1(2): 74–85
3. Arora S, Kumar R. Tubercular spinal epidural abscess involving the dorsal-lumbar-sacral region without osseous involvement. *The Journal of Infection in Developing Countries*. 2011 Jul 5;5(07):544-9.
4. Chen HC, Tzaan WC, Lui TN, spinal epidural abscesses: A retrospective analysis of clinical manifestations, sources of infection and outcomes. *Chang Gung medical journal*. 2004 May; 27(5): 351-8.
5. Polley P, Dunn R. Noncontiguous spinal tuberculosis: incidence and management. *European Spine Journal*. 2009 Aug 1;18(8):1096-101.
6. Merino P, Candel FJ, Gestoso I, Baos E, Picazo J. Microbiological diagnosis of spinal tuberculosis. *International Orthopaedics*. 2012;36(2):233-8.
7. Eyre Brook AL. A study of late results from disk operations present employment and residual complaints. *British Journal of Surgery*. 1952 Jan 1;39(156):289-96.
8. Kennedy JG, Soffe KE, McGrath A, Stephens MM, Walsh MG, McManus F Predictors of outcome in cauda equina syndrome. *Eur Spine J* 1999; 8: 317-322.
9. Garg RK, Somvanshi DS. Spinal tuberculosis: a review. *J Spinal Cord Med*. 2011;34(5):440-54
10. Lakhey R, Kafle D, Pokharel R. Tubercular epidural abscess presenting as acute cauda equina syndrome. *Journal of Institute of Medicine*. 2016;39(1).
11. Allison Bond and Farrin A. Manian, "Spinal Epidural Abscess: A Review with Special Emphasis on Earlier Diagnosis," *BioMed Research International*, vol. 2016, Article ID 1614328, 6 pages, 2016. doi:10.1155/2016/1614328
12. Mackenzie A, Laing R, Smith C, Kaar G, Smith F. Spinal epidural abscess: the importance of early diagnosis and treatment. *Journal of Neurology, Neurosurgery & Psychiatry*. 1998;65(2):209-12.

Giant Ganglioneuroma of Thoracic Spine: A Case Report

Das S¹, Alam K², Rashid MM³, Islam MM⁴, Mahbub H⁵, Khan MSI⁶

Abstract:

Ganglioneuromas most commonly arise from sympathetic ganglia. These neoplasms may be located wherever ganglion cells are normally found from the skull base to the pelvis, including the adrenal medulla. We describe a 15-year-old girl with giant Ganglioneuromas in the thoracic spine, who underwent successful resection (T4-5 level) of the tumor. Histopathological examination confirmed the diagnosis. Ganglioneuromas should be considered in the differential diagnosis of any paraspinal mass. A high index of suspicion and correlation of clinico-radiological findings is necessary in differentiating a large benign tumor from a malignant growth. Complete surgical excision is the treatment of choice; however tumor size and location need to be considered for the surgical approach (one-step or multiple surgeries). Close follow-up after surgery is mandatory.

Bang. J Neurosurgery 2016; 5(2): 51-54

Introduction:

Ganglioneuromas are rare benign tumors that originate from a neural crest or sympathetic ganglion¹. They most commonly appear in the posterior mediastinum and abdomen². The patients exhibit no obvious symptoms upon nervous system examination. The ganglioneuromas are often found in females, while the male/female ratio is approximately 2/3³. The incidence of ganglioneuroma is not well documented, but it is estimated to characterize 0.1 to 0.5 % of total central nervous system (CNS) tumors⁴. Paravertebral ganglioneuroma and scoliosis is rarer and has only been sporadically reported. We report a giant paraspinal Ganglioneuroma extending into extradural space and thoracic cavity.

Case Report:

A 15-year-old girl presented with a one month history of progressive onset lower extremity weakness. The patient had no other significant complaints. On examination, she was unable to stand or walk without support.

A neurological examination revealed 3/5 strength in all muscle groups in the patient's lower extremities. Sensory was markedly decreased below the nipple line. Deep tendon reflexes were hyperreflexic throughout the lower extremities along with markedly increased tone, right greater than left. Sustained clonus was noted at both ankles.

Magnetic resonance imaging showed a huge well defined mixed signal intensity oval lesion in posterior mediastinum predominantly on the right side. The lesion predominantly hypointense in T1WI and predominantly hyperintense in T2WI. Foci of signal void areas within the lesion represents calcification T1WI and T2WI increased signal intensity areas represents fatty component. The lesion causes collapse of right lung sparing basal segments of right lower lobe. Mediastinal structure are pushed of left. Epidural extension of the lesion is noted into the spinal canal causing widening of the right neural foramen resulting compression over the spinal cord and corresponding nerve root compression at D4-D5 level. Spinal cord pushed to left.


Right Posterolateral thoracotomy was done. A huge tumor occupying right paravertebral space pushing the right lung medially and inferiorly. It has a broad base, Posterior end of right 4th rib thin out and became triangular shaped. Tumor invaded in between the space.

1. Dr. Sukriti Das, Associate Professor, Department of Neurosurgery, Dhaka Medical College & Hospital, Dhaka.
2. Dr. Kamrul Alam, Professor and Head of the Department of thoracic surgery, Dhaka Medical College & Hospital, Dhaka.
3. Dr. Md. Mamunur Rashid, Phase-B, Resident, Department of Neurosurgery, Dhaka Medical College & Hospital, Dhaka
4. Dr. Md. Manirul Islam, Medical Officer, Department of Neurosurgery, Dhaka Medical College & Hospital, Dhaka.
5. Dr. Hasan Mahbub, Phase-B, Resident, Department of Neurosurgery, Dhaka Medical College & Hospital, Dhaka.
6. Dr. Mohammed. Shamsul Islam Khan, Medical Officer, Department of Neurosurgery, Dhaka Medical College & Hospital, Dhaka.



Address of Correspondence: Dr. Sukriti Das, Associate Professor, Department of Neurosurgery, Dhaka Medical College & Hospital, Dhaka, Cell: +8801711676848, e-mail: sukriti66@yahoo.com,



Pre-op chest x-ray




Pre-op CT scan of chest




A. Non-contrast
Pre-op MRI of dorsal spine- sagittal (T1)




B. Contrast




Sagittal section
Pre-op MRI of dorsal spine- (T2)




Coronal section




Pre-op MRI of dorsal spine Axial section



Post-op MRI of dorsal spine- sagittal view



Post-op MRI of dorsal spine- axial view



Post-op MRI of dorsal spine- coronal view

Discussion:

About 10% of Ganglioneuromas may involve the spinal canal⁵. Paraspinal Ganglioneuromas can extend into the spinal canal, forming dumb bell shaped tumor. However, in rare cases intradural extension has been reported. Most Ganglioneuromas are incidentally detected, and the symptoms, if any, are usually due to the mass effect. Rarely, the tumor may secrete vasoactive intestinal polypeptide, resulting in diarrhea. As this slow growing tumor extends through the neural foramen into the spinal cord, some patients may present with neurological deficits or scoliosis⁶. Thoracic intradural extramedullary Ganglioneuromas are very rare. In our patient, the characteristic feature was the remarkably large tumor size infiltrating into the thoracic cavity along with an intradural component. Microscopically, these tumors contain large ganglion cells and show areas with smaller lymphocyte-like cells within a matrix of fibrous stroma and schwann cells. The distinction from malignant tumor is based on the absence of necrosis or presence of any immature ganglion cells⁷.

It is usually safe and feasible to perform complete excision of Ganglioneuromas. However, in case of multiple and/or large-sized tumors, multi-stage dissection should be considered. In the present case, there were dense adhesions of the tumor with the nerve roots at the foraminal portions, which were left undisturbed during dissection. Ganglioneuromas generally has a favorable prognosis given its low metastatic potential⁸.

Conclusion:

This report describes a rare case of multiple Ganglioneuromas of the thoracic spine with intradural and thoracic extension. Ganglioneuromas should be considered in the differential diagnosis of a paraspinal mass. Although complete surgical resection is the best treatment option, stage-wise surgical resection should be considered in large-sized and/or multiple tumors, with close follow up.

References:

1. Geoerger B, Hero B, Harms D, Grebe J, Scheidhauer K, Berthold F. Metabolic activity and clinical features of primary ganglioneuromas. *Cancer*. 2001; 91(10):1905–13.

2. Lai PL, Lui TN, Jung SM, Chen WJ. Spinal ganglioneuroma mimicking adolescent idiopathic scoliosis. *Pediatr Neurosurg*. 2005;41(4):216–9.
3. Stout AP. Ganglioneuroma of the sympathetic nervous system. *Surg Gynecol Obstet*. 1947;84:101–10.
4. Choi YH, Kim IO, Cheon JE, Kim WS, Yeon KM, Wang KC, et al. Gangliocytoma of the spinal cord: a case report. *Pediatr Radiol*. 2001;31(5): 377–80.
5. Pang BC, Tchoyoson Lim CC, Tan KK : Giant spinal ganglioneuroma. *J Clin Neurosci* 12 : 967-972, 2005
6. Kara T, Oztunali C : Radiologic findings of thoracic scoliosis due to giant ganglioneuroma. *Clin Imaging* 37 : 767-768, 2013
7. Ugarriza LF, Cabezudo JM, Ramirez JM, Lorenzana LM, Porras LF : Bilateral and symmetric C1-C2 dumbbell ganglioneuromas producing severe spinal cord compression. *Surg Neurol* 55 : 228-231, 2001
8. Son DW, Song GS, Kim YH, Lee SW : Ventrally located cervical dumbbell ganglioneuroma producing spinal cord compression. *Korean J Spine* 10 : 246-248, 2013