

Name: _____

Advanced Calculus 2

Instructor Piotr Hajłasz

Final Exam

Due on May 1, 2020

Problem	Possible points	Score
1	10	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
9	10	
10	10	
11	10	
12	10	
Total	120	

You need 100 points so 2 problems is a bonus.

Problem 1. Suppose $f : \mathbb{R}_+^2 \rightarrow \mathbb{R}$ is a continuous function defined on

$$\mathbb{R}_+^2 = \{(x, y) : x \in \mathbb{R}, y > 0\}.$$

Assume also that the limits

$$g(u, v) = \lim_{t \rightarrow 0} \frac{f((u+t) \cos v, (u+t) \sin v) - f(u \cos v, u \sin v)}{t},$$

and

$$h(u, v) = \lim_{t \rightarrow 0} \frac{f(u \cos(v+t), u \sin(v+t)) - f(u \cos v, u \sin v)}{t},$$

exist and define continuous functions g, h on the domain

$$D = \{(u, v) : u > 0, 0 < v < \pi\}.$$

Prove that the function f is differentiable on \mathbb{R}_+^2 .

Proof. WRITE YOUR SOLUTION HERE. □

Problem 2. Let $f \in C^1(\mathbb{R})$ be a continuously differentiable function such that $|f'(x)| \leq 1/2$ for all $x \in \mathbb{R}$. Define $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$g(x, y) = (x + f(y), y + f(x)).$$

Prove that

- (1) g is a diffeomorphism,
- (2) $g(\mathbb{R}^2) = \mathbb{R}^2$,
- (3) the area $|g([0, 1]^2)|$ of the image of the unit square belongs to the interval $[3/4, 5/4]$.

Hint: Among other tools use the contraction principle.

Proof. WRITE YOUR SOLUTION HERE. □

Problem 3. Prove that the tangent planes to the surface S defined by $x^2 + y^2 - z^2 = 1$ at the points $(x, y, 0) \in S$ are parallel to the z -axis.

Proof. WRITE YOUR SOLUTION HERE. □

Problem 4. Prove that if $f : [0, 1] \rightarrow [0, 1]$ is a continuous function then its graph as a subset of \mathbb{R}^2 has measure zero.

Proof. WRITE YOUR SOLUTION HERE. □

Problem 5. Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a continuous function. Let $K \subset \mathbb{R}^3$ be a compact set such that $|f(x) - f(y)| \leq 2012|x - y|^2$ for all $x, y \in K$. Prove that the set $f(K) \subset \mathbb{R}^2$ has measure zero as a subset of \mathbb{R}^2 .

Proof. WRITE YOUR SOLUTION HERE. □

Problem 6. Assume that $f : [0, 1] \rightarrow [0, 1]$ is a continuous function such that the set $\{x \in [0, 1] : f(x) = 1\}$ has measure zero. Prove directly (without using any results like monotone or dominated convergence theorem) that

$$\lim_{n \rightarrow \infty} \int_0^1 f(x)^n dx = 0.$$

Proof. WRITE YOUR SOLUTION HERE. □

Problem 7. Let $\gamma : \mathbb{R} \rightarrow \mathbb{R}^n$ and $\mathbf{v}_i : \mathbb{R} \rightarrow \mathbb{R}^n, i = 1, 2, \dots, n-1$, be C^∞ smooth functions such that for any $t \in \mathbb{R}$ the vectors

$$\gamma'(t), \mathbf{v}_1(t), \dots, \mathbf{v}_{n-1}(t)$$

form an orthonormal basis of \mathbb{R}^n (here we differentiate γ but **do not** differentiate $\mathbf{v}_i, i = 1, 2, \dots, n-1$).

Consider the mapping $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by

$$\Phi(x_1, \dots, x_n) = \gamma(x_n) + \sum_{i=1}^{n-1} x_i \mathbf{v}_i(x_n).$$

- (a) Find the derivative $D\Phi(x_1, \dots, x_n)$;
- (b) Prove that Φ is a diffeomorphism in a neighborhood of and point of the form $(0, \dots, 0, x_n)$;
- (c) Find the limit

$$\lim_{r \rightarrow 0} \frac{|\Phi(B^n(0, r))|}{|B^n(0, r)|},$$

where $B^n(0, r)$ denotes the ball of radius r centered at the origin and $|A|$ stands for the volume of the set A .

Proof. WRITE YOUR SOLUTION HERE. □

Problem 8. Prove that if $K \in C^1(\mathbb{R}^2 \setminus \{(0, 0)\})$ satisfies the estimate

$$|\nabla K(x)| \leq \frac{1}{|x|^3} \quad \text{for all } x \neq (0, 0)$$

then there is a constant $C > 0$ such that

$$\iint_{\{x \in \mathbb{R}^2 : |x| > 2|y|\}} |K(x - y) - K(x)| dx \leq C$$

for all $y \in \mathbb{R}^2$.

Hint: Use the mean value theorem to estimate $|K(x - y) - K(x)|$ and then integrate in polar coordinates.

Proof. WRITE YOUR SOLUTION HERE. □

Problem 9. Use Green's theorem to prove the following result: If the vertices of a polygon, in counterclockwise order, are $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$, then the area of the polygon is

$$A = \frac{1}{2} \sum_{i=1}^n (x_i y_{i+1} - x_{i+1} y_i),$$

where we use notation $x_{n+1} = x_1, y_{n+1} = y_1$.

Proof. WRITE YOUR SOLUTION HERE. □

Problem 10. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with C^1 boundary and let $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \Phi(x, y) = (u(x, y), v(x, y))$, be a C^2 diffeomorphism. Prove that

$$\int_{\partial\Omega} uv_x dx + uv_y dy = \pm |\Phi(\Omega)|,$$

where $|\Phi(\Omega)|$ denotes the area of $\Phi(\Omega)$ and $\partial\Omega$ has positive orientation. Show on examples that both cases $+|\Phi(\Omega)|$ and $-|\Phi(\Omega)|$ are possible.

Proof. WRITE YOUR SOLUTION HERE. □

Problem 11. Let f be a polynomial of total degree at most three in $(x, y, z) \in \mathbb{R}^3$. Prove that:

$$\int_{x^2+y^2+z^2 \leq 1} f(x, y, z) dx dy dz = \frac{4\pi f((0, 0, 0))}{3} + \frac{2\pi (\Delta f)((0, 0, 0))}{15}.$$

Here $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ is the Laplacian operator on \mathbb{R}^3 .

Proof. WRITE YOUR SOLUTION HERE. □

Problem 12. Suppose that $K \in C^\infty(\mathbb{R}^2 \setminus \{0\})$ and

$$K(x) = \frac{K(x/\|x\|)}{\|x\|} \quad \text{for all } x \in \mathbb{R}^2 \setminus \{0\}.$$

- (a) Prove that $\nabla K(tx) = t^{-2}\nabla K(x)$ for $x \neq 0$ and $t > 0$.
- (b) Use the divergence theorem to prove that (on both sides we integrate vector valued functions)

$$\int_{\{1 \leq \|x\| \leq 2019\}} \nabla K(x) dx = \int_{\partial\{1 \leq \|x\| \leq 2019\}} K(x) \vec{n} d\sigma(x).$$

- (c) Prove that

$$\int_{\{1 \leq \|x\| \leq 2019\}} \nabla K(x) dx = 0.$$

Hint. Show first that $K(tx) = t^{-1}K(x)$ for $x \neq 0$, $t > 0$. In (a) differentiate $K(tx)$. Part (a) is not needed for parts (b) and (c).

Proof. WRITE YOUR SOLUTION HERE. □