

Title: Variation in leaf transcriptome responses to elevated ozone corresponds with physiological sensitivity to ozone across maize inbred lines

Adalena V. Nanni<sup>1,4</sup>, Alison M. Morse<sup>1,4</sup>, Jeremy R. B. Newman<sup>3,4</sup>, Nicole E. Choquette<sup>5</sup>, Jessica M. Wedow<sup>5</sup>, Zihao Liu<sup>1,4</sup>, Andrew D. B. Leakey<sup>5</sup>, Ana Conesa<sup>2,7</sup>, Elizabeth A. Ainsworth<sup>5,6</sup>, Lauren M McIntyre<sup>\*1,4</sup>

<sup>1</sup>Department of Molecular Genetics and Microbiology, University of Florida, Gainesville Florida 32611

<sup>2</sup>Department of Cell and Microbial Sciences, University of Florida, Gainesville Florida 32611

<sup>3</sup>Department of Pathology, University of Florida, Gainesville Florida 32611

<sup>4</sup>Genetics Institute, University of Florida, Gainesville Florida 32611

<sup>5</sup>Carl R. Woese Institute for Genomic Biology, Department of Plant Biology, and Department of Crop Sciences, University of Illinois at Urbana-Champaign

<sup>6</sup>USDA ARS Global Change and Photosynthesis Research Unit, Urbana, IL

<sup>7</sup>Institute for Integrative Systems Biology, Spanish National Research Council, Paterna, Spain.

\* Lauren M. McIntyre corresponding author

**Email:** [mcintyre@ufl.edu](mailto:mcintyre@ufl.edu)

**ORCID:**

0000-0003-4286-6489 Alison M Morse; 0000-0002-0077-3359 – Lauren M McIntyre; 0000-0001-8486-2801- Jeremy Newman; 0000-0001-5410-9026 Jessica Wedow; 0000-0002-3199-8999. Elizabeth Ainsworth; 0000-0001-6251-024X Andrew Leakey; 0000-0001-9597-311X Ana Conesa

none of the authors have any competing interests

## Abstract

We examine the impact of sustained elevated ozone concentration on the leaf transcriptome of 5 diverse maize inbred genotypes, which vary in physiological sensitivity to ozone (B73, Mo17, 6 Hp301, C123, NC338), using long reads to assemble transcripts and short reads to quantify 7 expression of these transcripts. More than 99% of the long reads, 99% of the assembled 8 transcripts, and 97% of the short reads map to both B73 and Mo17 reference genomes. 9 Approximately 95% of the genes with assembled transcripts belong to known B73-Mo17 10 syntenic loci and 94% of genes with assembled transcripts are present in all temperate lines in 11 the NAM pan-genome. While there is limited evidence for alternative splicing in response to 12 ozone stress, there is a difference in the magnitude of differential expression among the 5 13 genotypes. The transcriptional response to sustained ozone stress in the ozone resistant B73 14 genotype (151 genes) was modest, while more than 3,300 genes were significantly differentially 15 expressed in the more sensitive NC338 genotype. There is the potential for tandem duplication in 16 30% of genes with assembled transcripts, but there is no obvious association between potential 17 tandem duplication and differential expression. Genes with a common response across the 5 18 genotypes (83 genes) were associated with photosynthesis, in particular photosystem I. The 19 functional annotation of genes not differentially expressed in B73 but responsive in the other 4 20 genotypes (789) identifies reactive oxygen species. This suggests that B73 has a different 21 response to long term ozone exposure than the other 4 genotypes. The relative magnitude of the 22 genotypic response to ozone, and the enrichment analyses are consistent regardless of whether 23 aligning short reads to: long read assembled transcripts; the B73 reference; the Mo17 reference. 24 We find that prolonged ozone exposure directly impacts the photosynthetic machinery of the 25 leaf.

## Keywords

corn, stress response, climate change, Iso-seq, RNA-seq

## 57 Background

58 Tropospheric ozone is a prevalent, phytotoxic air pollutant, which significantly reduces global  
59 crop production (BURNEY and RAMANATHAN 2014; MCGRATH *et al.* 2015; MILLS *et al.* 2018).  
60 In the U.S., ozone pollution over the past 30 years was estimated to decrease maize yields by  
61 10% (MCGRATH *et al.* 2015); a loss equivalent to the impact of aridity stress, temperature stress  
62 or nutrient stress (MILLS *et al.* 2018). Exposure of plants to elevated ozone concentration causes  
63 reactive oxygen species (ROS) production in cells and knowledge of the responses this elicits is  
64 expected to aid understanding of oxidative stress associated with other abiotic and biotic factors  
65 (CONKLIN and BARTH 2004; GADJEV *et al.* 2006). Loss of photosynthetic capacity associated  
66 with accelerated senescence is a key response of maize to elevated ozone and differs among  
67 genotypes (YENDREK *et al.* 2017a). Ozone stress increased the heritability of photosynthetic  
68 traits and strengthened genetic correlations among traits in maize with substantial differences  
69 among hybrids observed (CHOQUETTE *et al.* 2019).

70 In a range of species, presence/absence variation (PAV) of genes related to abiotic stress  
71 response have been observed (YAO *et al.* 2015; GOLICZ *et al.* 2016; JIN *et al.* 2016;  
72 MONTENEGRO *et al.* 2017; WANG *et al.* 2018b; ZHAO *et al.* 2018; GAO *et al.* 2019; HOOPES *et al.*  
73 2019; LIU *et al.* 2020). In Amborella, PAV of genes have been linked primarily to responses to  
74 abiotic stress such as salt, water deprivation, and heat (HU *et al.* 2021). Gene families that have  
75 observed PAV associated with abiotic stress response include late embryonic abundant (LEA)  
76 genes (GOYAL *et al.* 2005; JIN *et al.* 2016), ROS signaling genes (MONTENEGRO *et al.* 2017), and  
77 genes related to the stay-green trait (QIAN *et al.* 2016). Alternative splicing (AS) has been  
78 reported to be associated with environmental stress response (MARRS and WALBOT 1997;  
79 EGAWA *et al.* 2006; LI *et al.* 2013; DING *et al.* 2014; CALIXTO *et al.* 2018). Divergence of AS,  
80 due to sequence divergence in splice sites or motifs, has been associated with stress response  
81 genes in Arabidopsis (WANG *et al.* 2019).

82 This study investigated the transcriptional response of maize to ozone stress in 5 diverse  
83 genotypes (B73, Mo17, Hp301, C123, NC338). Differences in a transcriptional response to  
84 ozone may be a consequence of PAV, differential splicing, and differential expression (DE).  
85 Large portions of grass genomes are syntenic (HULBERT *et al.* 1990; AHN and TANKSLEY 1993;  
86 MOORE *et al.* 1995; GALE and DEVOS 1998; FEUILLET and KELLER 2002). A recent *de novo*

87 assembly of the maize Mo17 genotype (SUN *et al.* 2018) and corresponding in-depth comparison  
88 to reference B73 genotype found only 122 genes present exclusively in one of the two genomes,  
89 and another pairwise comparison of B73 to a European inbred found ~400 genes were not shared  
90 (DARRACQ *et al.* 2018). Despite macrocollinearity, conservation of genes and gene order, there is  
91 evidence for genic rearrangements (reviewed in BENNETZEN 2005). Within maize, there is  
92 support for a pan-genome with documented divergence among genomes (SCHNABLE *et al.* 2009;  
93 LAI *et al.* 2010; HIRSCH *et al.* 2014), a megabase deletion (HAN *et al.* 2019), pervasive  
94 presence/absence variation (PAV) (SPRINGER *et al.* 2009; SWANSON-WAGNER *et al.* 2010), and  
95 exon shuffling (DOONER and WEIL 2007). In addition, a study of 503 inbred lines suggested that  
96 a “substantial portion of variation may lie outside the single reference genome for the species”  
97 (HIRSCH *et al.* 2014). A recent comprehensive evaluation of 26 maize genotypes (parents of the  
98 Nested Association Mapping Panel, NAM (YU *et al.* 2008)) finds evidence of a pan-genome  
99 with more than 100,000 genes and only approximately 30% shared among all 26 genotypes. B73  
100 is reported to contain between 63% and 74% of all genes in the pan-genome (HUFFORD *et al.*  
101 2021). At the heart of these discrepancies are varied views on how to account for nearly identical  
102 paralogs, tandem duplicates, and genes that have been duplicated through transposition.  
103 Whatever the nomenclature, it is important to consider that there are potential differences in  
104 dosage of highly similar transcripts and proteins between genotypes.

105 Differential expression in response to abiotic stress occurs in many plant species (reviewed in  
106 FUJITA *et al.* 2006), including maize (reviewed in SALIKA and RIFFAT 2021). Changes to  
107 expression following abiotic stresses related to climate change (e.g., heat, drought, elevated  
108 greenhouse gases, etc.) have also been observed (reviewed in AHUJA *et al.* 2010). Differential  
109 expression must be due to variation in cis-regulatory elements either directly affecting expression  
110 or indirectly leading to downstream effects (reviewed in WITTKOPP 2007). Cis-regulation  
111 induced by abiotic stress can include factors such as transposable element (e.g., ITO *et al.* 2013;  
112 MAKAREVITCH *et al.* 2015), epigenetic responses (e.g., VERHOEVEN *et al.* 2018;  
113 ENTRAMBASAGUAS *et al.* 2021; reviewed in KIM 2021) or other cis-regulatory factors (e.g.,  
114 MARUYAMA *et al.* 2004; ZOU *et al.* 2011; RICCI *et al.* 2019).

115 Maize genotypes exist that show important phenotypic differences, including response to stress  
116 and ozone susceptibility (HULBERT *et al.* 1990; YENDREK *et al.* 2017a; AINSWORTH *et al.* 2020;

CHOQUETTE *et al.* 2020; WEDOW *et al.* 2021a). We seek to answer what the basis of the observed differences in ozone response among maize cultivars is, such as differences in the gene content (PAV), differences in splicing (AS), or differences in transcriptional regulation (DE)? We sequenced the leaf transcriptome of five diverse maize genotypes, B73, Mo17, C123, NC338 and Hp301, in ambient and elevated ozone conditions. While C123 and Mo17 are somewhat closely related, the other genotypes represent a diverse array of temperate genotypes including Hp301, a popcorn genotype (See Supplementary Figure 1 in LIU *et al.* 2003). These five genotypes are selected based on variation in physiological response to ozone in field experiments (CHOQUETTE *et al.* 2019) where B73 was the least sensitive of the 5 genotypes. We focus on the following questions: Is the expressed transcriptome part of the shared genome? Is there evidence for alternative splicing in response to stress? Is the transcriptional response to ozone similar among all 5 temperate genotypes? We used long read sequencing and an unbiased approach to assemble transcripts for each genotype and treatment (n=10) independently. These assembled transcripts were rigorously examined for diversity and then the union of the assembled transcriptomes was used as a transcriptional reference for quantitation of short read data from an experiment with twelve independent replicate plants for each genotype/condition (n=120).

## Results

*Do the leaf transcriptomes of different maize lines represent the shared genome, or, on the contrary, reveal genotype specific genes?*

The majority of clusters are high quality with few technical artifacts. We use IsoSeq3, which does not require a reference genome (<https://github.com/PacificBiosciences/IsoSeq3>), to obtain high-quality polished clusters for each genotype/condition. Clustering is an intermediate step in the transcript assembly process (see methods). Clusters from each genotype/condition are mapped to the three available reference genomes (Figure 1, Supplementary Table 1, line 32). Despite genomic divergence (SUN *et al.* 2018), the cluster mapping rates for clusters from the Hp301, C123 and NC338 samples are similar to the B73 and Mo17 samples (Figure 1B) with 92.8-99.5% mapping at 95% identity or higher and at least 95% of the read length when mapped to the B73 and Mo17 references (Figure 1B, Supplementary Table 1, line 35).

The relatively few unmapped clusters are likely technical artifacts. There are **12** unmapped clusters in the B73 set of high-quality polished clusters mapped to the B73 genome: 5 in the ambient condition and 7 in the ozone condition. Only 2 of these 12 are greater than 1 kb in length. These are further examined by a BLAST (ALTSCHUL *et al.* 1990) against the B73 reference genome. Three of the 12 B73 clusters hit to the same small (61bp) region of the B73 genome while the remaining 9 sequences did not return a hit. These results indicate that these 12 unmapped clusters from B73 likely reflect technical artifacts. Similarly, for the 3 Mo17 samples, 50 high quality polished clusters are unmapped to the Mo17 Cau reference (14 in the ambient ozone library and 36 in the 2 elevated ozone libraries) (Supplementary Table 1, line 32). Similar to B73, BLAST results of partial hits from Mo17 clusters to the Mo17 genome contain mismatches and gaps. Taken together, this indicates that less than 1% of the polished clusters for these 4 samples are likely technical artifacts. BLAST results for unmapped clusters from Hp301, C123 and NC338 have similar results ([https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)).\_


Potential copy number variation. We note that there are larger numbers of ‘ignored’ clusters (clusters that map but have low identity or low coverage) for Mo17, C123, NC338, and Hp301 mapping to B73 and for B73, C123, NC338, and Hp301 mapping to Mo17 compared to B73 mapping to B73 or Mo17 Mapping to Mo17(Supplementary Table 1 line 37). Some of these clusters are likely incomplete transcripts. Clusters less than 1000 nt (20%-40% of those ignored) suggest potential incomplete transcript sequences. Most of the low identity and low coverage clusters with at least 1,000 nt have BLAST hits to maize with mismatches and gaps (83-90%) (Supplementary Table 1 line 39, [https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)) potentially suggesting genetic variation. The B73 clusters mapping with low identity to Mo17, but not B73, were examined to provide potential insight into the excess of clusters of low coverage in the Hp301, C123 and NC338 genomes. These clusters overall map well to B73 and have a similar length distribution as the clusters that map to both B73 and Mo17. We note that several of the best hits are to gene families (Supplementary Table 1 line 40) so we cannot preclude the possibility that there is ambiguity in assignment of clusters to genes due to variants among the genotypes that have diverged or the possibility that B73 and Mo17 have unique duplications for some of these

176 genes (tandem or dispersed). For completeness all cluster sequences are provided  
177 ([https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)).

178 Assembled transcripts are in core or near-core genes when compared to the NAM pan-genome.  
179 We assemble transcripts from mapped clusters using Cupcake ToFU  
180 ([https://github.com/Magdoll/cDNA\\_Cupcake](https://github.com/Magdoll/cDNA_Cupcake)). We match the junctions in sequential order  
181 (junction chain) of the resulting assembled transcripts to the existing reference annotations for  
182 B73 or Mo17 using SQANTI QC (TARDAGUILA *et al.* 2018). The B73 and Mo17 Yan references  
183 contain junction chains that match the majority of the assembled transcripts (Supplementary  
184 Figure 2B, Supplementary Table 1, lines 49 and 50). The short reads from all genotypes also  
185 map to both B73 and Mo17 genomes with only 1% to 3% unmapped (Supplementary Table 2).

186 A unique list of B73 v4 reference genes (n=15,055) representing the genes associated with the  
187 assembled transcripts across all samples (Supplementary Table 1, line 46) is compared to genes  
188 in the NAM pan-genome (HUFFORD *et al.* 2021). In this list there are 13,369 genes associated  
189 with a NAM pan-gene with 90.6% annotated as core (present in all parents of the NAM). An  
190 additional ~6.1% are near-core (present in all but 1 of the 26 NAM founders) and ~94.4% are  
191 present in all temperate NAM genotypes (HUFFORD *et al.* 2021). Of the 13,369 reference genes  
192 with associated assembled transcripts, 29 are annotated in the NAM population as private to B73  
193 (present in only NAM B73). Of these 29 genes, 12 are expressed only in the B73 samples  
194 (Supplementary Table 4) and the other 17 genes are detected in our experiments of temperate  
195 lines not in the NAM collection.

196 Assembled transcriptome is 90%-94% B73-Mo17 syntenic. The high mapping rates to both  
197 reference genomes led us to hypothesize that the genes expressed in the leaf are part of the  
198 syntenic genome in B73-Mo17. To determine whether the assembled transcripts are from  
199 syntenic loci in B73 and Mo17, we evaluate B73-Mo17 synteny information using SynMap  
200 (HAUG-BALTZELL *et al.* 2017) and SynFind within CoGe (LYONS and FREELING 2008).  
201 Approximately 90 percent of the assembled transcripts from B73 and Mo17 map to B73-Mo17  
202 syntenic loci. Using the syntenic list of B73-Mo17 loci from Sun *et al.* (SUN *et al.* 2018) shows a  
203 similar result (~94% mapping of assembled transcripts to syntenic loci).



**Figure 1. A) CCS long reads and B) Clusters map equally well** to the B73 v4 or Mo17 Yan reference genomes for all 5 genotypes. The darker color (dark blue or dark red) indicates percent reads (A) or clusters (B) with alignment identity greater than or equal to 95% to the indicated reference sequence and alignment length greater than or equal to 95% of the read or cluster length. Percent reads or clusters with alignment identity less than 95% and/or alignment length less than 95% of the read or cluster length is indicated by a light blue or light red color. The percent mapped CCS long reads ranges from 99.4% to 99.9% and percent mapped clusters range from 99.5% to 99.99% (Supplementary Table 1). Mapping to annotated reference transcripts is also similar among all 5 genotypes (Supplementary Figure 3).

Assembled transcripts that map to intergenic regions ('novel' in SQANTI) or to more than one genic locus ('fusion' in SQANTI) are expressed in multiple genotypes. To determine whether



218 these loci are detected in the temperate genotypes in this study, we look at both short and long  
219 reads. Out of the 1,575 novel, fusion, or antisense loci, 928 are detected in all genotypes and  
220 1,335 are detected in multiple genotypes (Supplementary Figure 4A, B). There are 3 novel loci  
221 expressed exclusively in B73. BLAST results of the transcripts within these loci indicate partial  
222 hits to B73, Mo17, and other maize genomes including putative transposase and retrotransposon  
223 genes ([https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)).


224 There is potential ambiguity in gene identity among putative tandem duplicates and putative  
225 identical paralogs. Hoopes et al. (HOOPES *et al.* 2019) identified paralogous groups of genes. For  
226 the 12,604 annotated genes in the assembled leaf transcriptome, there are 1,032 paralogous  
227 groups containing 2,390 genes where each paralogous group contains a minimum of 2 genes.  
228 This demonstrates that transcripts are assembled separately for paralogs with sequence diversity.  
229 Approximately ~1% of maize genes have nearly identical paralogs (EMRICH *et al.* 2007),  
230 meaning they exhibit 99% similarity. In this case, long reads originating from nearly identical  
231 paralogs would not be separable. This is a problem with RNA-seq experiments in maize and is a  
232 general limitation in such studies. To further examine this issue, we use the annotation of  
233 putative tandem duplicates among the parents of the NAM population (HUFFORD *et al.* 2021).  
234 We find that 30% of the 12,604 annotated genes in this study have a potential tandem duplication  
235 in at least one of the parents of the NAM. This is an enrichment compared to the 16% of all  
236 genes reported as potential tandem duplicates in the NAM pan-genome (HUFFORD *et al.* 2021).  
237 The high number of potential duplicates and the ambiguity among paralogs is a limitation of this  
238 study and all studies of expression in this species (Supplementary File 5).

239 There is limited evidence for transcriptional presence/absence variation in this study. There is a  
240 distinction between evidence for expression and being able to conduct an analysis for differential  
241 expression (Figure 2). Differences in transcriptional presence/absence variation among maize  
242 inbred lines has been observed (BROHAMMER *et al.* 2018; HOOPES *et al.* 2019) and it has been  
243 hypothesized that presence/absence of transcription contributes to stress responses in maize  
244 (HOOPES *et al.* 2019). For genes with assembled transcripts (n=12,604), we have evidence of  
245 expression in all five genotypes for all but 81 genes (Figure 2A). Of these 81 annotated genes, 40  
246 were previously identified as genes with potential PAV (BROHAMMER *et al.* 2018; HOOPES *et al.*

247 2019). It is possible that transcriptional PAV is present in these genes. These genes can be found  
 248 in Supplementary File 5 (where *flag\_detect\_num\_geno\_gt0\_lt5* = 1 and *flag\_pav* = 1).

249 The ability to analyze the 12,604 annotated genes for expression is more variable across  
 250 genotypes (Figure 2B). The majority of the genes (8951, 71%) are analyzable (TPM greater than  
 251 5 in at least 50% of replicates) in all 5 genotypes. Only 549 are analyzable in only one genotype.  
 252 Expression is lower for the genes with fewer analyzable genotypes (Figure 3). Of the 549 genes  
 253 analyzable in only one genotype, 473 are detected in all 5 genotypes. This finding is likely due  
 254 to sampling variation (EMRICH *et al.* 2007).


255 Our long and short read data map to B73 and Mo17 references with ~94% of the genes  
 256 previously identified as B73-Mo17 syntenic. There is some limited evidence for genes expressed  
 257 in the leaf in C123, Hp301, or NC338 in either ambient or elevated ozone that are exclusive to  
 258 those genotypes.



259

260 **Figure 2. Genes with assembled transcripts have evidence of expression and are analyzable.**  
 261 (A) Evidence of expression, or detection, is defined as the presence of a single CCS long read or  
 262 short read mapping to exonic gene regions. A gene is considered to have evidence for expression

in a genotype if it is detected in either ambient or ozone conditions. The upset plot rows are the total number of genes with evidence of expression in each genotype. The columns indicate the unique combinations. The sum of the columns is the number of genes with at least 1 assembled transcript (12,604). There are only 81 genes without evidence of expression in all 5 genotypes. (B) Analyzable expression is defined as at least one transcript with TPM greater than 5 in at least 50% of RNA-seq replicates. The upset plot has the same format as in panel A. The majority of genes (77%) are analyzable in all five genotypes.



**Figure 3. Expression as a function of the number of genotypes analyzable for differential expression.** Mean gene expression (transcripts per million, TPM) for each genotype and treatment. B73 is blue, C123 is red, Hp301 is gray, Mo17 is green and NC338 is orange. Genes are analyzable if at least one transcript has a TPM greater than 5 in 50% of short-read replicates in either ambient or elevated ozone condition for that genotype. The number of genes for each group analyzable in 1 to 5 genotypes is indicated below each set of representative boxplots, totaling 11,813 genes. Most genes are analyzable in all 5 genotypes (8,951). For genes analyzable in fewer than 5 genotypes, expression values are relatively low. This indicates that it is unlikely that presence/absence variation is at play, and instead more likely that presence/absence is due to sampling variation.


*Is there evidence for alternative splicing in response to stress?*

The majority of genes in the assembled leaf transcriptome have a single transcript. The union of the individually assembled transcripts from all genotypes and conditions are combined into a single unique set of assembled transcripts using chaining ([https://github.com/Magdoll/cDNA\\_Cupcake](https://github.com/Magdoll/cDNA_Cupcake)). The assembled transcripts are annotated using the B73 v4 (JIAO *et al.* 2017) reference annotation with SQANTI (TARDAGUILA *et al.* 2018)

(Supplementary Table 1, lines 62-71). After chaining there are 1,531 putative novel loci (187 of which are novel antisense loci of annotated genes). To remove redundancies, transcripts with the same splice junctions are consolidated and unspliced fragments in annotated genes with multi-exon reference genes are removed. Since transcripts with identical splice junction chains cannot be easily distinguished with RNA-seq, we selected the longest representative of each unique splicing pattern in the curated transcriptome. The curated assembled transcriptome has 31,388 assembled transcripts and 12,604 genes. More than half (6,895 genes, ~54%) of the annotated genes have a single associated assembled transcript, and ~17% have more than 3 associated transcripts (Supplementary Figure 5B). The average number of transcripts per annotated gene is ~2.5 and the median is 1. The 9,451 genes in Wang et al. (WANG *et al.* 2018a) (~550,000 starting CCS reads in the leaf tissue dataset) represent 21% of all B73 v4 reference genes (Supplementary Figure 6A). Of these genes, 9% have 3 or more transcripts and 77% have one transcript. Starting with roughly 6 times the number of CCS reads as Wang et al. (WANG *et al.* 2018a), we find approximately twice as many assembled transcripts (31,388 vs. 13,580). Other tissues also showed a median of 1 transcript per gene (Supplementary Figure 6B-F).

Alternative splicing is not a major contributor to the maize leaf response to ozone stress. We estimate gene expression in alternative transcripts by mapping Illumina short read data from 120 individual plants (n=12 per experimental condition; n=24 per genotype) to the assembled leaf transcriptome using RSEM (LI and DEWEY 2011). We then evaluate differential isoform usage in the transcriptional response to ozone stress using tappAS (DE LA FUENTE *et al.* 2019). We assayed all genes with multiple transcripts in a particular genotype for differential isoform usage between ambient and elevated ozone conditions (Supplementary Figure 7). Despite reports of splicing as a response to stress in plants (IIDA *et al.* 2004; FILICHKIN *et al.* 2010; THATCHER *et al.* 2016; reviewed in LALOUM *et al.* 2018), only 59 annotated genes are identified as having major isoform switching in NC338 (< 2% of multi-transcript genes detected in NC338), and fewer than 35 genes (~1% or less of all multi-transcript genes) in each of the other 4 genotypes, with 1 gene (Zm00001d043613) having evidence for major isoform switching in B73 (Supplementary Table 5). Eighteen annotated genes are identified as having differential isoform usage with annotations linked to plant stress responses.

319 We note that there are quite a few annotated genes with differential isoform usage where the  
320 predicted coding regions of the transcripts are non-overlapping, leading to potentially distinct  
321 protein sequences. An example is shown in Supplementary Figure 8. This suggests that ozone  
322 stress response may differentially regulate the proteins produced by these genes, or that there are  
323 two similar genes in proximity, annotated as a single gene, that respond differently to ozone  
324 stress.



**Figure 4. Differentially expressed genes.** (A) Count of differential expression by genotype for all genes analyzed. Genes significantly differentially expressed in the labeled genotype left bar. The top histogram totals to the number of genes analyzed for differential expression (11,401) and the categories are mutually exclusive. For example, genes differentially expressed in NC338 only (998, orange). Relatively few genes are differentially expressed in all genotypes (83, red circle). More are differentially expressed in the 4 genotypes M017, C123, Hp301 and NC338. (B) For each genotype, a scatterplot is shown of the  $\log_2(\text{TPM})$  ambient on the X axis and the  $\log_2(\text{TPM})$  ozone on the Y axis. Differentially expressed (DE) genes in the indicated genotype are in pink, and in exactly one other genotype are blue, in two other genotypes are yellow, in 3 other genotypes are purple and in all 5 genotypes (gray diamonds). The largest fold changes are consistently apparent in those genes with the gray diamonds. Transcripts DE in only one genotype have smaller fold changes than those DE in more than one genotype (as seen by the orange points nearer the diagonal). (C) Genes significant for differential expression in at least one genotype (4,432). Rows are genes and columns are genotypes. Categories are ordered as in Panel A (matching color key is the right bar, numbers indicate the numbers of genes in each of the groups). Within each category in the right bar the rows are sorted by the B73 response, in order to provide a consistent reference point. The first (orange) group is significant for NC338 and the second group visualizes the 789 that are significant for the 4 genotypes M017, C123, Hp301 and NC338.

*Is the transcriptional response to ozone similar among all 5 temperate genotypes?*

Genotypes differ in the magnitude of the response to ozone stress. B73 shows a limited response to elevated ozone, with 151 differentially expressed genes, while NC338 shows evidence for more than 3,300 differentially expressed genes (Figure 4A). The number of differentially expressed genes is progressively greater in the other genotypes: Hp301 (1807 genes), C123 (2085 genes) and Mo17 (2382 genes). The number of genes differentially expressed in a genotype-specific manner followed the same pattern: B73 (8 genes), Hp301 (157 genes), C123 (223 genes), Mo17 (333 genes) and NC338 (998 genes) (Figure 4A). Across the four more ozone sensitive genotypes (non-B73) there are 789 differentially expressed genes. Eighty-three genes are identified as differentially expressed in all five genotypes (Supplementary File 4, flag\_analyze\_DE\_all5), thus representing a common transcriptional response to elevated ozone concentration.


For genes significant in only one genotype, the estimated fold change is relatively modest, with the response to ozone in B73 substantially smaller in all comparisons (Figure 4B, 3C). In each genotype, enriched gene sets (SUBRAMANIAN *et al.* 2005) for the Gene Ontology biological

process category included photosynthesis (e. g., photosystem I, protein-chromophore linkages) (Supplementary Table 6, *flag\_analyze\_DE\_genotype*).

Diverse genotypes show a common transcriptional response to ozone stress. Are differential responses to ozone stress associated with gene expression differences? The largest fold changes are found in genes significant in all genotypes (Figure 4B, Supplementary Figure 9,10), and the proportion of up- and down-regulated genes is balanced for all genotypes, including B73 and NC338 (Figure 4B, Supplementary Figure 11). The regulation of expression in response to ozone is in the same direction for 82 of the 83 transcripts differentially expressed in all 5 genotypes (Figure 4C, red). Zm00001d027525 (basic endochitinase B, EnsemblPlants) is the only transcript that is down-regulated in B73 under ozone conditions but up-regulated in the other 4 genotypes. Genes differentially expressed in all 5 genotypes are enriched for the following GO categories: biological process (heat shock response, photosynthesis (light harvesting in photosystem I, light reaction, response to red light, and protein-chromophore linkage)); cellular component category (photosystem I, photosystem II and plastoglobule); and molecular function category (chlorophyll binding and pigment binding) (Supplementary Table 5, *flag\_analyze\_DE\_all5*).

In addition to photosynthesis, additional GO enrichment categories are found when genes significantly differentially expressed in the four more sensitive genotypes are analyzed (Figure 4A, n=789). These include GO biological processes related to metabolism (aromatic amino acid family biosynthetic process, carotenoid biosynthetic process, coenzyme biosynthetic process, glycine catabolic process, hydrogen peroxide catabolic process, oxylipin biosynthetic process, regulation of lipid metabolic process, unsaturated fatty acid biosynthetic process), additional photosynthesis terms (photosynthetic electron transport in photosystem I, photosystem I assembly, reductive pentose-phosphate cycle, triose phosphate transmembrane transport, response to red light, response to light stimulus, response to low light intensity stimulus), protein-chromophore linkage, iron-sulfur cluster assembly, and detection of biotic stimulus (Supplementary Table 6, *flag\_analyze\_DE\_all\_noB73*). The direction of regulation of the genes significant in these 4 more ozone sensitive genotypes is in the same direction for 788 of the 789 genes (~46% up-regulated, ~54% down-regulated) with Zm00001d037513 (germ-like protein1, EnsemblPlants) up-regulated in C123 and down-regulated in the other 3 genotypes (Figure 4C).

Chlorophyll content estimate from a spectrophotometer follows the same pattern as gene expression. The transcriptional response to ozone stress is largely attributable to genes involved in photosynthetic and respiratory functions in the leaf. These general effects are common across genotypes, but the magnitude of response varies. We hypothesize that the magnitude of the transcriptional response to ozone is correlated with changes in leaf chlorophyll content. To evaluate this hypothesis, we re-analyze the ambient and ozone data in Yendrek et al. (YENDREK *et al.* 2017b) for the maize genotypes in this study. In agreement with our transcriptional analysis, we find that chlorophyll levels are significantly lower under elevated ozone conditions for NC338 ( $p < 0.00001$ ), Mo17 ( $p < 0.01$ ), and C123 ( $p = 0.003$ ), while there is no significant difference in chlorophyll content in the ambient and elevated ozone for B73 (Figure 5).



399

**Figure 5:** The effect of elevated ozone on the PLSR chlorophyll score for temperate genotypes B73, C123, Hp301, Mo17 and NC338. The chlorophyll content was estimated by high throughput spectrophotometry in Yendrek et al. (2017b). Shown here are effect sizes estimated from least squares means of the PLSR chlorophyll in elevated ozone relative to ambient ozone conditions. B73 has the highest levels of PLSR chlorophyll and is these levels are not significantly affected by ozone.



Results are robust to different short read mapping strategies. In short read data derived from gene families, there can be ambiguity as to the origin of these reads. In the literature, there is a general consensus that, overall, there is a reduction in bias when genes and transcripts unlikely to be expressed are eliminated from consideration in mapping (MATSUOKA *et al.* 2002; MCMULLEN *et al.* 2009; HIRSCH and SPRINGER 2017; NEWMAN *et al.* 2018; TARDAGUILA *et al.* 2018). However, it is possible that while there is overall bias reduction, for some genes true signal may be missed with this strategy. We conducted 3 additional differential expression analyses where we map short reads to the genome instead of the estimated transcriptome in order to evaluate whether this strategy impacted our findings (Supplementary Methods). We mapped all short reads to the B73 genome, and separately to the Mo17 Cau genome and for all analyzable genes performed differential expression analyses and a GO enrichment. In order to compare results, we restricted the GO enrichment to the set of genes with one-to-one B73-Mo17 syntenic gene pairs. We compared numbers of reads mapping for the one-to-one syntenic gene pairs and they are similar (Supplementary Figure 3). All inferences about differential expression are consistent with the results described above. In all cases, B73 has low levels of differential expression in response to ozone exposure and NC338 has orders of magnitude higher expression (See [https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)). Many of the GO terms were identical among all 4 analyses, and variation among GO terms did not identify new potential mechanisms of action but rather pointed to the same overall processes. All results from all 4 analyses are provided, and the summary of the GO enrichment across all 4 analyses is also provided (See [https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)).

*In summary, our results indicate that the magnitude of the expression response to elevated ozone mirrors physiological differences among maize genotypes.*

## **Discussion**

Despite the diversity of the maize genotypes and the extensive literature on structural variation among maize genotypes, the long and short read data collected from leaf tissue in these experiments of temperate genotypes largely reflect a set of shared transcripts. Clustering of CCS long reads was carried out independently and without a reference, in order to avoid potential reference bias. A close examination of the CCS long reads, the clusters for each genotype, and the assembled transcriptome reveal that the overwhelming majority (~94%) of genes expressed

in the leaf are part of the annotated B73-Mo17 syntenic genome. The biological inferences of the differential expression analyses are robust to alternative mapping strategies.

The pattern of the expression response to elevated ozone is very similar in loci differentially expressed in all 5 genotypes (82/83). Loci differentially expressed in NC338, Hp301, Mo17 and C123 also display the same pattern in over/under expression among these 4 genotypes (788/789), even as the magnitude of expression in response to elevated ozone is dramatically different, with NC338 having the strongest response. We see no evidence for structural variation (or tandem duplication) in the transcriptional response to ozone but cannot rule out variation in *cis* regulation of these genes, which may well be influenced by adjacent variable genomic content (STUPAR and SPRINGER 2006; DOONER and WEIL 2007; HAWKINS *et al.* 2014; MAKAREVITCH *et al.* 2015; HIRSCH and SPRINGER 2017; STEIGE *et al.* 2017; ROESSLER *et al.* 2018).

There are 83 differentially expressed genes in response to elevated ozone in all 5 maize genotypes. Decreased expression of genes involved in light harvesting dominated the conserved transcriptional response. These include chlorophyll biosynthesis genes, chlorophyll a/b binding proteins, and subunits of photosynthetic proteins. Down-regulation of photosynthesis is a common transcriptional response to ozone (XU *et al.* 2015), and chlorophyll biosynthetic genes are also reduced in an ozone sensitive Medicago accession (PUCKETTE *et al.* 2008). In an ozone sensitive soybean genotype, decreased photosystem II gene expression is observed after 24 hrs of ozone exposure (WALDECK *et al.* 2017). The sensitivity of photosynthesis to ozone is clear across both C<sub>3</sub> and C<sub>4</sub> crops, and remains a critical target for improving ozone tolerance (EMBERSON *et al.* 2018). Perhaps to combat damage to photosynthetic proteins, increased expression of heat shock proteins is another common response of the 5 genotypes to elevated ozone.

The genotypes assayed in this study differed in photosynthetic capacity under elevated ozone in the field (YENDREK *et al.* 2017a) and they differ in transcriptional responses to ozone stress. Allelic variation in transcriptional response to stress has been documented (GUO *et al.* 2004). Over 20 times more genes are differentially expressed in NC338 in ambient and elevated ozone compared B73, and the transcriptional response of B73 is muted relative to the other genotypes. In a companion field experiment, there is a greater decline in chlorophyll content associated with accelerated senescence in inbreds C123 and NC338 as compared to B73 (YENDREK *et al.*

2017b). The metabolite profile of B73 was not responsive to elevated ozone in the field (WEDOW *et al.* 2021b). Additionally, hybrids containing parents Hp301 and NC338 showed greater reductions in photosynthesis under elevated ozone in the field (CHOQUETTE *et al.* 2019). Ozone tolerance has been associated with a dampened transcriptional response in genotypes of other species including *Arabidopsis* (XU *et al.* 2015), soybean (BURTON *et al.* 2016; WALDECK *et al.* 2017) and *Medicago* (PUCKETTE *et al.* 2008). A dampened transcriptional response to other abiotic stresses including drought (FRACASSO *et al.* 2016), cold stress (DA MAIA *et al.* 2017) and salt stress (YANG *et al.* 2017b) was observed in tolerant genotypes.

These results are consistent with oxidative stress tolerance being associated with a weaker response to a reactive oxygen species signal. Notably, B73 has equivalent, or greater, stomatal conductance than the more sensitive maize genotypes studied here (YENDREK *et al.* 2017b). This means that the flux of ozone into the leaf mesophyll would tend to be greatest in B73. That an equal or greater ozone dose elicited a weaker transcriptional response, suggests that genetic variation in downstream processes is key to ozone tolerance; i.e., the processes that either sense ROS in the apoplast, transduce signals, produce the secondary burst of ROS in the cytoplasm, or respond to the secondary ROS burst (AINSWORTH 2017). For example, GO enrichment analysis (Supplementary Table 6), indicated differential expression of genes related to *cellular response to hydrogen peroxide*, *detection of biotic stimulus*, *hydrogen peroxide catabolic process*, and *regulation of response to biotic stimulus* in the ozone sensitive genotypes, but not B73. At the same time, there is differential gene expression related to *response to salicylic acid* in B73 that is not observed in the more ozone sensitive genotypes. The strength of ozone impacts on *Arabidopsis* varies with the strength of ethylene-signaling components of the pathway when modulated genetically or chemically (RAO *et al.* 2002). The transcriptional data presented here suggest a similar strategy might be relevant in maize and could assist in the identification of potential targets for manipulation.

## Conclusions:

The long and short read data collected from leaf tissue in these experiments of temperate genotypes (B73, Mo17, Nc338, C123, Hp301) largely reflect a set of shared transcripts. While the overall response to ozone is largely consistent in direction, the extent of the response is dramatically different among the genotypes, ranging from 150 genes significantly differentially

expressed (B73) to more than 3,000 genes significantly differentially expressed (NC338). B73 has a dampened transcriptional response, suggesting potential fundamental differences compared to the other 4 genotypes.

We found that all genotypes show regulatory responses in genes related to photosynthesis. Our findings show that ozone directly impacts the photosynthetic machinery of the leaf, and a potential key element to reduction in sensitivity to ozone is a controlled response to the presence of ozone.

## **Methods:**

Experimental design and sample collection: Twenty-four plants from each genotype were grown in 5.7 L pots in LC1 mix (Sun Gro Horticulture Distribution Inc., Bellevue, WA, USA) in six growth chambers (Growth Chamber, Chagrin Falls, OH, USA) set to maintain a 15 hr day at 25 °C and photosynthetic photon flux (PPF) of  $\sim 350 \mu\text{mol m}^{-2} \text{s}^{-1}$ , 9 hr night at 21 °C, and a relative humidity of 60%. We grew 120 plants in the six growth chambers, with three chambers maintained at low ozone conditions ( $< 10$  ppb) and three at elevated ozone conditions (100 ppb). Ozone is generated using a variable output UV-C light bulb ballast (HVAC 560 ozone generator, Crystal Air, Langley, Canada) and plants are fumigated with 100 ppb ozone for 6 hrs per day from emergence to sampling. Four plants of each of the five genotypes, B73, C124, Hp301, NC338, Mo17 were randomly assigned to each chamber for a total of twenty-four plants per genotype. Plants were fertilized (Osmocote Blend 19-5-8) at the start of the experiment. Twenty-six days after planting, leaf tissue from the 5<sup>th</sup> leaf is sampled directly into liquid N<sub>2</sub> for a total of 120 independent leaf samples (5 genotypes x 4 plants per genotype x 2 treatment conditions x 3 growth chambers per treatment) for a total of 12 replicates per genotype and ozone treatment. Leaf samples were sent to University of Florida where leaf punches were collected in a cold room on dry ice. A separate sample from a single plant from each genotype and ozone treatment was collected for long read sequencing.

Long-read library preparation and sequencing: Leaf tissue ( $\sim 175$  mg) from a single individual plant of each genotype/treatment combination (n=10) was collected and snap frozen in liquid nitrogen. Total RNA was isolated independently for each of the 10 samples (5 genotypes x 2 treatments) followed by full-length cDNA synthesis using the SMARTer PCR cDNA Synthesis kit and single molecule SMRTbell library construction followed by PCR optimization and ELF

size selection. One library (Mo17 elevated) was selected to calibrate the run, and then all ten libraries sequenced on the PacBio SEQUEL platform using 1 LR SMRT cell per sample and SMRT Link 6.0.0 chemistry to generate raw subreads for each sample by the University of Florida Interdisciplinary Center for Biotechnology Research (ICBR). Data (subread.bam files) have been deposited at the SRA (BioProject accession PRJNA604929).

RNA-seq library preparation and sequencing: Approximately 20mg of flash frozen leaf tissue from each sample was placed into individual mini tubes in a 96-tube plate format (Axygen MTS-11-C-R). Sample freeze-drying, mRNA purification, cDNA synthesis and library preparation were carried out by Rapid Genomics (Gainesville, FL) to generate an individual dual indexed library per sample. Individual libraries were quantified and pooled to generate equimolar samples for sequencing. The final pooled library was sequenced on 3 lanes of an Illumina HiSeq 3000 at Rapid Genomics and 1 lane of Illumina Novaseq at Novogene (Sacramento, CA). Data have been deposited at the SRA (BioProject accession PRJNA604929).

Mapping short-reads to available reference genomes: Adapter trimmed unique short reads are mapped to B73 (Ensembl version 41, [ftp://ftp.ensemblgenomes.org/pub/plants/release-41/fasta/zea\\_mays/](ftp://ftp.ensemblgenomes.org/pub/plants/release-41/fasta/zea_mays/), (JIAO *et al.* 2017)), Mo17 Yan (<https://ftp.maizegdb.org/MaizeGDB/FTP/Mo17-YAN/>, (YANG *et al.* 2017a)) and Mo17 Cau (<https://ftp.maizegdb.org/MaizeGDB/FTP/Mo17-CAU/>, (SUN *et al.* 2018)) using BWA-mem with default parameters and ‘-M’ to mark short split reads as secondary alignment (v0.7.17, (LI 2013)). Unique primary alignments with an alignment score of at least 30 (default of BWA-mem) are selected for further analysis. The maximum percentage of unmapped reads was 3.8% therefore more than 96% of the reads mapped to all three genomes (Supplementary Table 2). On average, 98% of reads across all samples mapped to the B73 v4 reference genome. The sample with the lowest mapping proportion of reads of all samples was a NC338 elevated ozone sample ([https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022))

Long-read pre-processing: The IsoSeq3 pipeline (v3.0.0, <https://github.com/PacificBiosciences/IsoSeq3>) was used to process subreads into circular consensus reads (CCS (v3.1.0)). CCS for each sample had, on average, approximately 3 subreads, indicating successful multiple passes during sequencing and enabling error correction. Primers were removed from CCS using lima (v1.7.1). IsoSeq3 cluster was used to trim polyA

556 tails, remove false concatemers, and cluster full-length reads by sequence similarity. Consensus  
557 ('polished') sequences for each cluster were generated using IsoSeq3 polish. The expected error  
558 rate after this step was less than < 2% (WEIRATHER *et al.* 2017). Up to this point no genome  
559 references were used.

560 Mapping CCS: CCS, after primer trimming, were mapped to the three available reference  
561 genomes using minimap2 (v2.12, (LI 2018)) and parameters recommended for PacBio IsoSeq  
562 processing (-ax splice -uf --secondary=no -C5,  
563 [https://github.com/Magdoll/cDNA\\_Cupcake/wiki/Best-practice-for-aligning-Iso-Seq-to-](https://github.com/Magdoll/cDNA_Cupcake/wiki/Best-practice-for-aligning-Iso-Seq-to-reference-genome:-minimap2,-deSALT,-GMAP,-STAR,-BLAT/24228b7e9c329268b0fdc85226097129d9c44a82)  
564 [reference-genome:-minimap2,-deSALT,-GMAP,-STAR,-BLAT/](https://github.com/Magdoll/cDNA_Cupcake/wiki/Best-practice-for-aligning-Iso-Seq-to-reference-genome:-minimap2,-deSALT,-GMAP,-STAR,-BLAT/24228b7e9c329268b0fdc85226097129d9c44a82)  
565 [24228b7e9c329268b0fdc85226097129d9c44a82](https://github.com/Magdoll/cDNA_Cupcake/wiki/Best-practice-for-aligning-Iso-Seq-to-reference-genome:-minimap2,-deSALT,-GMAP,-STAR,-BLAT/24228b7e9c329268b0fdc85226097129d9c44a82)) (Supplementary Figure 12). Mapping rates  
566 exceeded 99% (Supplementary Table 1, lines 23 and 32). Unmapped reads were compared to  
567 the NCBI nucleotide non-redundant database using BLAST (ALTSCHUL *et al.* 1990). Greater  
568 than 47% (3,613 out of 7,637) of unmapped reads had no BLAST hit, the remaining reads  
569 resulted in partial hits, many with numerous mismatches and gaps.

570 Mapping clusters: Although there were similar numbers of CCS reads across samples  
571 (Supplementary Table 1, line 8), the number of clusters was more variable: 24,000 to 44,000  
572 clusters (Supplementary Table 1, line 23). The polished cluster sequences were mapped to the  
573 three available references using minimap2 (v2.12, (LI 2018)) and parameters recommended for  
574 PacBio IsoSeq3 processing (-ax splice -uf --secondary=no -C5). Unmapped clusters were binned  
575 according to length and unmapped clusters larger than 1 kb were evaluated by BLAST  
576 (ALTSCHUL *et al.* 1990).

577 Assembled Transcripts for each sample: For clusters that map to each of the three references  
578 (Supplementary Table 1, line 24), we used the IsoSeq3 supporting script in Cupcake ToFU  
579 (version 20180629, [https://github.com/Magdoll/cDNA\\_Cupcake](https://github.com/Magdoll/cDNA_Cupcake), [collapse\\_isoforms\\_by\\_sam.py](https://github.com/Magdoll/cDNA_Cupcake/blob/master/collapse_isoforms_by_sam.py))  
580 to identify identical isoforms within each library and collapse them into a single unique  
581 assembled transcript (Supplementary Table 1, line 27).

582 SQANTI QC: SQANTI QC was used to match the assembled transcripts to an annotated  
583 reference transcript (TARDAGUILA *et al.* 2018) and to classify the resulting splice junction  
584 pattern. A full-splice match (FSM) is a transcript where all splice junctions in the assembled  
585 transcript match exactly to an annotated reference transcript. Incomplete-splice matches (ISM),

586 have junctions in the same order as a reference transcript, but may be missing some junctions on  
587 the 5' and/or 3' end of the transcript. Transcripts that contained novel combinations of annotated  
588 junctions, novel in catalog (NIC), and those with new junctions within an annotated gene, novel  
589 not in catalog (NNC) were also identified. In addition, novel transcripts within potentially genic  
590 areas were identified (TARDAGUILA *et al.* 2018). SQANTI QC was run on CCS long reads  
591 (Supplementary Table 1, line 17-20) assembled transcripts (Supplementary Table 1, line 44-52)  
592 and the assembled leaf transcriptome (Supplementary Table 1, line 62-71).

593 Evaluation of filtered clusters: Clusters that did not pass Cupcake ToFU filtering process  
594 ('ignored') due to low identity (< 95%) or low coverage (< 99%) alignments to the B73  
595 reference genome were potential candidates for genes with sequence divergence compared to the  
596 B73 reference. Alternatively, they may be paralogous genes in the non-B73 genotypes with  
597 PAV. Ignored clusters greater than 1 kb (of sufficient length to suggest a complete transcript)  
598 were evaluated using BLAST BLAST (ALTSCHUL *et al.* 1990) against the NCBI non-redundant  
599 nucleotide database (Supplementary Table 1, line 37-40). Best hits were defined as the BLAST  
600 hit with the largest bitscore and smallest e-value for each filtered cluster  
601 ([https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)). Best hits were further  
602 assessed for subject names associated with gene families ("family" or "kinase" in name). Ignored  
603 clusters from C123, Hp301, and NC338 were also compared to the assembled transcripts from  
604 the B73 and Mo17 samples using BLAST  
605 ([https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)).

606 Comparison of assembled transcripts to NAM pan-gene classifications. A recent assembly and  
607 annotation of 26 maize lines from the nested association mapping (NAM) population (including  
608 B73 and Hp301) identified pan-gene associations and classified B73 genes as "core" (in all 26  
609 lines), "near-core" (in 24 of the 26 lines), "dispensable" (in 2 to 23 lines), and "private" (only in  
610 a single line) (HUFFORD *et al.* 2021). We used the Ensembl B73 v4 reference annotation which  
611 required a conversion to the NAM associated v5 gene identifiers to compare pan-gene  
612 classifications. A file from maizeGDB (<https://maizegdb.org/>)(WOODHOUSE *et al.* 2021) was  
613 used to link v5 genes to each v4 gene  
614 ([https://download.maizegdb.org/Pan-genes/B73\\_gene\\_xref/B73v4\\_to\\_B73v5.tsv](https://download.maizegdb.org/Pan-genes/B73_gene_xref/B73v4_to_B73v5.tsv)). A unique list  
615 of B73 v4 reference genes representing the genes associated with the assembled transcripts

616 across all samples was compared to the pan-genes in the NAM population. Additionally, the  
617 12,604 annotated genes in the assembled leaf transcriptome were linked to the NAM pan-gene  
618 classifications.

619 Chaining multiple samples to identify redundancy: Cupcake ToFU IsoSeq3 supporting scripts  
620 (*summarize\_sample\_GFF\_junctions.py*, *scrub\_sample\_GFF\_junctions.py* and  
621 *chain\_samples.py*) were used to create a non-redundant union of assembled transcripts. The  
622 clustering and chaining protocol was carried out on unique assembled transcripts mapped to the  
623 B73 or Mo17 Yan references separately (Supplementary Figure 13), yielding two chained sets of  
624 assembled transcripts (Supplementary Table 1). The B73 and Mo17 Yan assembled  
625 transcriptomes had similar numbers of transcripts (79,620 and 76,980 respectively;  
626 Supplementary Table 1, line 60). Transcript quality control was performed with SQANTI QC  
627 and filter functions (TARDAGUILA *et al.* 2018) and included identification of transcripts with  
628 known technological artifacts such reverse transcriptase switching (TARDAGUILA *et al.* 2018).  
629 There was a slightly higher number of annotated genes (Supplementary Table 1 line 65) and  
630 transcripts with annotated junction chains (Supplementary Table 1 line 68-69) when the  
631 assembled transcriptomes were mapped to B73 v4 as compared to Mo17 Yan. There were fewer  
632 novel loci identified when the assembled transcriptomes were mapped to B73 v4 as compared to  
633 Mo17 Yan (Supplementary Table 1 line 69).

634 Synteny: SynMap (HAUG-BALTZELL *et al.* 2017) and SynFind within CoGe (LYONS and  
635 FREELING 2008) were used to identify collinear gene-sets between the B73 v4 reference and the  
636 Mo17 Cau reference (Margaret Woodhouse, maizeGDB). We also verified syntenic genes  
637 according to (SUN *et al.* 2018). Information in Supplementary File 5.

638 Novel genes: Assembled transcripts associated with putative novel loci were quantified using  
639 HTSeq (v 0.11.2, (ANDERS *et al.* 2015)). Novel loci show evidence for expression if at least one  
640 read is detected in at least one replicate of either treatment (Supplementary Figure 4).

641 Annotation of assembled transcripts: FASTA files of the assembled transcripts and the  
642 associated translated sequences were used as queries to obtain putative functional, structural or  
643 signaling motif information from InterProScan (version 5.29-68.0, (QUEVILLON *et al.* 2005)),  
644 TMHMM (version 2.0, (SONNHAMMER *et al.* 1998; KROGH *et al.* 2001)) and SignalP (version  
645 4.1, (NIELSEN *et al.* 1997; NIELSEN 2017)) for tappAS (DE LA FUENTE *et al.* 2019). Translated



transcripts were linked to proteins in public databases via sequence similarity (Supplementary Table 9). Repetitive regions were annotated using RepeatMasker (version 4.0.5, <http://www.repeatmasker.org>) and putative RNA regulatory sequences identified using ScanForMotif (BISWAS and BROWN 2014). Default parameters were used. We collated all the above annotation into a GFF-like file (Supplementary File 1).

Evidence for expression: The list of genes with assembled transcripts were evaluated for expression using HTSeq (version 0.11.2, (ANDERS *et al.* 2015)) with CCS primer trimmed reads and adapter trimmed unique short reads for each sample separately. Evidence of expression for each gene in each sample is defined as at least 1 read (short or long) mapping to the gene (Figure 2A).

Secondary Data: The raw B73 PacBio data (.bas.h5) were obtained from Wang et al. (WANG *et al.* 2018a) and consist of the following tissue types: leaf, silk, pericarp, bract, shoot and seedling (Supplementary Figure 6). All PacBio data were processed as described above. Clustering and polishing of trimmed reads yielded 38,741 high quality transcripts in the leaf.

Quantification of the assembled transcriptome: Paired-end Illumina short-read data was used for quantification of the B73 assembled leaf transcriptome. FASTQ files were trimmed of adapter sequence (cutadapt, version 2.1, (MARTIN 2011)) and merged if R1 and R2 reads overlapped (*bbmerge.py* from BBMap version 38.44, (BUSHNELL *et al.* 2017)). Duplicate reads were removed. Gene and transcript expression values estimated using RSEM with STAR (LI and DEWEY 2011; DOBIN *et al.* 2013) against the B73 v4 genome with the B73 assembled transcriptome GTF file (Supplementary File 2). Merged (single-end) and unmerged (paired-end) reads were aligned separately. TPM (transcripts per kilobase million) values are output for genes and transcripts. Single-end (R1-R2 merged reads) and paired-end counts were each assumed to represent 1 read and TPM were summed. Transcripts are analyzable for differential expression if their TPM values are greater than 5 in 50% of the replicates in either ambient or elevated ozone condition (Figure 2B, Supplementary File 3, flag *\_genotype\_condition*).

Differential expression: For each genotype, analyzable transcripts in either ambient or elevated ozone conditions were evaluated for differential expression and differential splicing using tappAS (version 1.0.6, (DE LA FUENTE *et al.* 2019)) at FDR level (BENJAMINI and HOCHBERG 1995) of 0.05 (Supplementary Figure 14). GO enrichment using Fisher's exact test and an FDR

threshold of 0.05 is performed (Supplementary Table 6). Analysis of chlorophyll data: We re-analyzed the PLSR predicted chlorophyll values from the Illinois FACE (Free Air Concentration Enrichment, <https://www.igb.illinois.edu/soyface/>) site under ambient and elevated ozone conditions (Yendrek et al. 2017b). Each ringpair was divided into 5 sub-blocks and genotypes were planted in each sub-block. The difference in chlorophyll (as measured by the PLSR model) between ozone and ambient conditions was tested in a mixed effects model with a random effect of ring-pair and fixed effects of genotype and ozone treatment. The model also included an effect for potential excess ozone application in one of the sub-blocks due to a prevailing wind.

## **Acknowledgments**

NSF Plant Genome Research Program (PGR-1238030; ADBL, LMM, and EAA) MCA-PGR: Genetic and genomic approaches to understand and improve maize responses to ozone NIH R01GM128193 (LMM), R03CA222444 (AC, LMM) USDA SoyFACE Global Change Research. Project Number: 5012-21000-030-17-S. Brad Barbazuk for helpful discussion, Margaret Woodhouse (with maizeGDB for synteny file), Junping Shi for providing the synteny list from Sun et al. 2018. Deborah Morse and James Resnick for assistance with sample processing. Francisco Pardo-Palacios and Pedro Salguero Garcia for help with maize functional annotation. Dr. Bo Wang, Peter van Buren and Dr. Doreen Ware for graciously sharing maize PacBio raw data. Three anonymous reviewers for excellent criticism.

## **Data Availability**

Raw data (BAM files for the PacBio data and FASTQ files for the RNA-seq data) are deposited under SRA BioProject accession PRJNA604929. All processed data are attached to this manuscript, including the GTF file for the transcriptome (Supplementary File 2), quantified expression data for analysis (Supplementary File 3) and annotations for the transcriptome (Supplementary File 1). All scripts and full detailed documentation for all the analyses are posted on github ([https://github.com/McIntyre-Lab/papers/tree/master/nanni\\_maize\\_2022](https://github.com/McIntyre-Lab/papers/tree/master/nanni_maize_2022)). All data files including all FASTA files used for BLAST analyses; all BLAST results, all results from alternate differential expression analyses, and all enrichment results from these alternative

705 analyses are deposited on zenodo (link- we will create this deposit when the final content is  
706 approved, these deposits are not easily updated).

707

#### 708 **Author contributions**

709 **Adalena V. Nanni** developed all analysis plans, analyzed all short-read data, contributed to  
710 analysis of long read data, annotated the maize transcriptomes, managed all references, designed  
711 and contributed to figures, contributed to the writing of the paper. Worked on the github page.

712 **Alison M. Morse** designed and executed the analysis performed all long-read analyses,  
713 interpreted the findings, designed and contributed to figures, contributed to the writing of the  
714 paper and is responsible for the github page.

715 **Jeremy R. B. Newman** participated in analysis, designed and contributed to figures, contributed  
716 to the writing of the paper.

717 **Nicole E. Choquette** identified genotypes for testing, performed the growth chamber experiment  
718 and collected tissue.

719 **Jessica M. Wedow** identified genotypes for testing, performed the growth chamber experiment  
720 and collected tissue.

721 **Zihao Liu** created the transcriptome visualization plots.

722 **Andrew D. B. Leakey** contributed to the overall design of the umbrella project (PGR-1238030),  
723 interpreted the findings, designed figures, interpreted the findings, contributed to the writing of  
724 the paper.

725 **Ana Conesa** contributed to the analyses of long reads, interpreted the findings, contributed to the  
726 writing of the paper.

727 **Elizabeth A. Ainsworth** contributed to the overall design of the umbrella project (PGR-  
728 1238030), designed this particular experiment, identified genotypes for testing, supervised tissue  
729 collection, participated in analysis, designed figures, interpreted the findings, contributed to the  
730 writing of the paper.

731 **Lauren M. McIntyre** contributed to the overall design of the umbrella project (PGR-1238030),  
 732 designed this particular experiment, designed and supervised the analysis, interpreted the  
 733 findings, contributed to the writing of the paper.

734

735 Ahn, S., and S. D. Tanksley, 1993 Comparative linkage maps of the rice and maize genomes.  
 736 Proceedings of the National Academy of Sciences of the United States of America 90:  
 737 7980-7984.

738 Ahuja, I., R. C. H. de Vos, A. M. Bones and R. D. Hall, 2010 Plant molecular stress responses  
 739 face climate change. Trends in Plant Science 15: 664-674.

740 Ainsworth, E. A., 2017 Understanding and improving global crop response to ozone pollution.  
 741 Plant Journal 90: 886-897.

742 Ainsworth, E. A., P. Lemonnier and J. M. Wedow, 2020 The influence of rising tropospheric  
 743 carbon dioxide and ozone on plant productivity. Plant Biology 22: 5-11.

744 Altschul, S. F., W. Gish, W. Miller, E. W. Myers and D. J. Lipman, 1990 Basic local alignment  
 745 search tool. Journal of Molecular Biology 215: 403-410.

746 Anders, S., P. T. Pyl and W. Huber, 2015 HTSeq-a Python framework to work with high-  
 747 throughput sequencing data. Bioinformatics 31: 166-169.

748 Benjamini, Y., and Y. Hochberg, 1995 Controlling the false discovery rate - a practical and  
 749 powerful approach to multiple testing. Journal of the Royal Statistical Society Series B-  
 750 Statistical Methodology 57: 289-300.

751 Bennetzen, J. L., 2005 Transposable elements, gene creation and genome rearrangement in  
 752 flowering plants. Current Opinion in Genetics & Development 15: 621-627.

753 Biswas, A., and C. M. Brown, 2014 Scan for Motifs: a webserver for the analysis of post-  
 754 transcriptional regulatory elements in the 3' untranslated regions (3' UTRs) of mRNAs.  
 755 BMC Bioinformatics 15: 174.

756 Brohammer, A. B., T. J. Y. Kono, N. M. Springer, S. E. McGaugh and C. N. Hirsch, 2018 The  
 757 limited role of differential fractionation in genome content variation and function in  
 758 maize (*Zea mays* L.) inbred lines. Plant Journal 93: 131-141.

759 Burney, J., and V. Ramanathan, 2014 Recent climate and air pollution impacts on Indian  
 760 agriculture. Proc Natl Acad Sci U S A 111: 16319-16324.

761 Burton, A. L., K. O. Burkey, T. E. Carter, J. Orf and P. B. Cregan, 2016 Phenotypic variation  
 762 and identification of quantitative trait loci for ozone tolerance in a Fiskeby III × Mandarin  
 763 (Ottawa) soybean population. Theor Appl Genet 129: 1113-1125.

764 Bushnell, B., J. Rood and E. Singer, 2017 BBMerge - Accurate paired shotgun read merging via  
765 overlap. Plos One 12.

766 Calixto, C. P. G., W. B. Guo, A. B. James, N. A. Tzioutziou, J. C. Entizne *et al.*, 2018 Rapid and  
767 Dynamic Alternative Splicing Impacts the Arabidopsis Cold Response Transcriptome.  
768 Plant Cell 30: 1424-1444.

769 Choquette, N., F. Ogut, T. Wertin, C. Montes, C. Sorgini *et al.*, 2019 Uncovering hidden genetic  
770 variation in photosynthesis of field-grown maize under ozone pollution. Global Change  
771 Biology 25: 4327-4338.

772 Choquette, N. E., E. A. Ainsworth, W. Bezodis and A. P. Cavanagh, 2020 Ozone tolerant maize  
773 hybrids maintain Rubisco content and activity during long-term exposure in the field.  
774 Plant Cell and Environment 43: 3033-3047.

775 Conklin, P. L., and C. Barth, 2004 Ascorbic acid, a familiar small molecule intertwined in the  
776 response of plants to ozone, pathogens, and the onset of senescence. Plant Cell and  
777 Environment 27: 959-970.

778 da Maia, L. C., P. R. B. Cadore, L. C. Benitez, R. Danielowski, E. J. B. Braga *et al.*, 2017  
779 Transcriptome profiling of rice seedlings under cold stress. Functional Plant Biology 44:  
780 419-429.

781 Darracq, A., C. Vitte, S. Nicolas, J. Duarte, J.-P. Pichon *et al.*, 2018 Sequence analysis of  
782 European maize inbred line F2 provides new insights into molecular and chromosomal  
783 characteristics of presence/absence variants. BMC Genomics 19.

784 de la Fuente, L., Á. Arzalluz-Luque, M. Tardáguila, H. del Risco, C. Martí *et al.*, 2019 tappAS: a  
785 comprehensive computational framework for the analysis of the functional impact of  
786 differential splicing. bioRxiv: 690743.

787 Ding, F., P. Cui, Z. Y. Wang, S. D. Zhang, S. Ali *et al.*, 2014 Genome-wide analysis of  
788 alternative splicing of pre-mRNA under salt stress in Arabidopsis. BMC Genomics 15.

789 Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski *et al.*, 2013 STAR: ultrafast  
790 universal RNA-seq aligner. Bioinformatics 29: 15-21.

791 Dooner, H. K., and C. F. Weil, 2007 Give-and-take: interactions between DNA transposons and  
792 their host plant genomes. Current Opinion in Genetics & Development 17: 486-492.

793 Egawa, C., F. Kobayashi, M. Ishibashi, T. Nakamura, C. Nakamura *et al.*, 2006 Differential  
794 regulation of transcript accumulation and alternative splicing of a DREB2 homolog under  
795 abiotic stress conditions in common wheat. Genes & Genetic Systems 81: 77-91.

796 Emberson, L. D., H. Pleijel, E. A. Ainsworth, M. van den Berg, W. Ren *et al.*, 2018 Ozone  
797 effects on crops and consideration in crop models. European Journal of Agronomy 100:  
798 19-34.

799 Emrich, S. J., L. Li, T. J. Wen, M. D. Yandea-Nelson, Y. Fu *et al.*, 2007 Nearly identical  
800 paralogs: Implications for maize (*Zea mays* L.) genome evolution. *Genetics* 175: 429-  
801 439.

802 Entrambasaguas, L., M. Ruocco, K. J. F. Verhoeven, G. Procaccini and L. Marin-Guirao, 2021  
803 Gene body DNA methylation in seagrasses: inter- and intraspecific differences and  
804 interaction with transcriptome plasticity under heat stress. *Scientific Reports* 11.

805 Feuillet, C., and B. Keller, 2002 Comparative Genomics in the grass family: Molecular  
806 characterization of grass genome structure and evolution. *Annals of Botany* 89: 3-10.

807 Filichkin, S. A., H. D. Priest, S. A. Givan, R. K. Shen, D. W. Bryant *et al.*, 2010 Genome-wide  
808 mapping of alternative splicing in *Arabidopsis thaliana*. *Genome Research* 20: 45-58.

809 Fracasso, A., L. M. Trindade and S. Amaducci, 2016 Drought stress tolerance strategies revealed  
810 by RNA-Seq in two sorghum genotypes with contrasting WUE. *BMC Plant Biology* 16:  
811 115.

812 Fujita, M., Y. Fujita, Y. Noutoshi, F. Takahashi, Y. Narusaka *et al.*, 2006 Crosstalk between  
813 abiotic and biotic stress responses: a current view from the points of convergence in the  
814 stress signaling networks. *Current Opinion in Plant Biology* 9: 436-442.

815 Gadjev, I., S. Vanderauwera, T. S. Gechev, C. Laloi, I. N. Minkov *et al.*, 2006 Transcriptomic  
816 footprints disclose specificity of reactive oxygen species signaling in *Arabidopsis*. *Plant*  
817 *Physiology* 141: 436-445.

818 Gale, M., and K. Devos, 1998 Comparative genetics in the grasses. *Proceedings of the National*  
819 *Academy of Sciences of the United States of America* 95: 1971-1974.

820 Gao, L., I. Gonda, H. H. Sun, Q. Y. Ma, K. Bao *et al.*, 2019 The tomato pan-genome uncovers  
821 new genes and a rare allele regulating fruit flavor. *Nature Genetics* 51: 1044-+.

822 Golicz, A. A., P. E. Bayer, G. C. Barker, P. P. Edger, H. Kim *et al.*, 2016 The pangenome of an  
823 agronomically important crop plant *Brassica oleracea*. *Nature Communications* 7.

824 Goyal, K., L. J. Walton and A. Tunnacliffe, 2005 LEA proteins prevent protein aggregation due  
825 to water stress. *Biochemical Journal* 388: 151-157.

826 Guo, M., M. A. Rupe, C. Zinselmeier, J. Habben, B. A. Bowen *et al.*, 2004 Allelic variation of  
827 gene expression in maize hybrids. *The Plant cell* 16: 1707-1716.

828 Han, X., Y. Qin, F. Yu, X. Ren, Z. Zhang *et al.*, 2019 A megabase-scale deletion is associated  
829 with phenotypic variation of multiple traits in maize. *Genetics* 211: 305-316.

830 Haug-Baltzell, A., S. Stephens, S. Davey, C. Scheidegger and E. Lyons, 2017 SynMap2 and  
831 SynMap3D: web-based whole-genome synteny browsers. *Bioinformatics* 33: 2197-2198.

832 Hawkins, J. S., V. Delgado, L. Feng, M. Carlise, H. K. Dooner *et al.*, 2014 Variation in allelic  
833 expression associated with a recombination hotspot in *Zea mays*. *The Plant Journal* 79:  
834 375-384.

835 Hirsch, C. D., and N. M. Springer, 2017 Transposable element influences on gene expression in  
836 plants. *Biochim Biophys Acta Gene Regul Mech* 1860: 157-165.

837 Hirsch, C. N., J. M. Foerster, J. M. Johnson, R. S. Sekhon, G. Muttoni *et al.*, 2014 Insights into  
838 the maize pan-genome and pan-transcriptome. *Plant Cell* 26: 121-135.

839 Hoopes, G. M., J. P. Hamilton, J. C. Wood, E. Esteban, A. Pasha *et al.*, 2019 An updated gene  
840 atlas for maize reveals organ-specific and stress-induced genes. *Plant Journal* 97: 1154-  
841 1167.

842 Hu, H., A. Scheben, B. Verpaalen, S. Tirnaz, P. E. Bayer *et al.*, 2021 Amborella gene  
843 presence/absence variation is associated with abiotic stress responses that may contribute  
844 to environmental adaptation. *The New phytologist*.

845 Hufford, M. B., A. S. Seetharam, M. R. Woodhouse, K. M. Chougule, S. J. Ou *et al.*, 2021 De  
846 novo assembly, annotation, and comparative analysis of 26 diverse maize genomes.  
847 *Science* 373: 655-+.

848 Hulbert, S. H., T. E. Richter, J. D. Axtell and J. L. Bennetzen, 1990 Genetics-mapping and  
849 characterization of sorghum and related crops by means of maize DNA probes.  
850 *Proceedings of the National Academy of Sciences of the United States of America* 87:  
851 4251-4255.

852 Iida, K., M. Seki, T. Sakurai, M. Satou, K. Akiyama *et al.*, 2004 Genome-wide analysis of  
853 alternative pre-mRNA splicing in *Arabidopsis thaliana* based on full-length cDNA  
854 sequences. *Nucleic Acids Research* 32: 5096-5103.

855 Ito, H., T. Yoshida, S. Tsukahara and A. Kawabe, 2013 Evolution of the ONSEN retrotransposon  
856 family activated upon heat stress in Brassicaceae. *Gene* 518: 256-261.

857 Jiao, Y., P. Peluso, J. Shi, T. Liang, M. Stitzer *et al.*, 2017 Improved maize reference genome  
858 with single-molecule technologies. *Nature* 546: 524-+.

859 Jin, M. L., H. J. Liu, C. He, J. J. Fu, Y. J. Xiao *et al.*, 2016 Maize pan-transcriptome provides  
860 novel insights into genome complexity and quantitative trait variation. *Scientific Reports*  
861 6.

862 Kim, J. H., 2021 Multifaceted Chromatin Structure and Transcription Changes in Plant Stress  
863 Response. *International Journal of Molecular Sciences* 22.

864 Krogh, A., B. Larsson, G. von Heijne and E. L. Sonnhammer, 2001 Predicting transmembrane  
865 protein topology with a hidden Markov model: application to complete genomes. *J Mol*  
866 *Biol* 305: 567-580.

867 Lai, J., R. Li, X. Xu, W. Jin, M. Xu *et al.*, 2010 Genome-wide patterns of genetic variation  
868 among elite maize inbred lines. *Nature Genetics* 42: 1027-+.

869 Laloum, T., G. Martin and P. Duque, 2018 Alternative Splicing Control of Abiotic Stress  
870 Responses. *Trends in Plant Science* 23: 140-150.

871 Li, B., and C. N. Dewey, 2011 RSEM: accurate transcript quantification from RNA-Seq data  
872 with or without a reference genome. *BMC Bioinformatics* 12: 323.

873 Li, H., 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM,  
874 pp. arXiv:1303.3997.

875 Li, H., 2018 Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34: 3094-  
876 3100.

877 Li, W. F., W. D. Lin, P. Ray, P. Lan and W. Schmidt, 2013 Genome-Wide Detection of  
878 Condition-Sensitive Alternative Splicing in Arabidopsis Roots. *Plant Physiology* 162:  
879 1750-1763.

880 Liu, K., M. Goodman, S. Muse, J. S. Smith, E. Buckler *et al.*, 2003 Genetic structure and  
881 diversity among maize inbred lines as inferred from DNA microsatellites. *Genetics* 165:  
882 2117-2128.

883 Liu, Y. C., H. L. Du, P. C. Li, Y. T. Shen, H. Peng *et al.*, 2020 Pan-Genome of Wild and  
884 Cultivated Soybeans. *Cell* 182: 162-+.

885 Lyons, E., and M. Freeling, 2008 How to usefully compare homologous plant genes and  
886 chromosomes as DNA sequences. *Plant Journal* 53: 661-673.

887 Makarevitch, I., A. J. Waters, P. T. West, M. Stitzer, C. N. Hirsch *et al.*, 2015 Transposable  
888 Elements Contribute to Activation of Maize Genes in Response to Abiotic Stress. *Plos*  
889 *Genetics* 11.

890 Marrs, K. A., and V. Walbot, 1997 Expression and RNA splicing of the maize glutathione S-  
891 transferase Bronze2 gene is regulated by cadmium and other stresses. *Plant Physiology*  
892 113: 93-102.

893 Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads.  
894 2011 17: 3.

895 Maruyama, K., Y. Sakuma, M. Kasuga, Y. Ito, M. Seki *et al.*, 2004 Identification of cold-  
896 inducible downstream genes of the Arabidopsis DREB1A/CBF3 transcriptional factor  
897 using two microarray systems. *Plant Journal* 38: 982-993.

898 Matsuoka, Y., Y. Vigouroux, M. M. Goodman, J. Sanchez G, E. Buckler *et al.*, 2002 A single  
899 domestication for maize shown by multilocus microsatellite genotyping. *Proc Natl Acad*  
900 *Sci U S A* 99: 6080-6084.



901 McGrath, J. M., A. M. Betzelberger, S. Wang, E. Shook, X. G. Zhu *et al.*, 2015 An analysis of  
902 ozone damage to historical maize and soybean yields in the United States. *Proc Natl*  
903 *Acad Sci U S A* 112: 14390-14395.

904 McMullen, M. D., S. Kresovich, H. S. Villeda, P. Bradbury, H. Li *et al.*, 2009 Genetic properties  
905 of the maize nested association mapping population. *Science* 325: 737-740.

906 Mills, G., K. Sharps, D. Simpson, H. Pleijel, M. Broberg *et al.*, 2018 Ozone pollution will  
907 compromise efforts to increase global wheat production. *Glob Chang Biol* 24: 3560-  
908 3574.

909 Montenegro, J. D., A. A. Golicz, P. E. Bayer, B. Hurgobin, H. Lee *et al.*, 2017 The pangenome  
910 of hexaploid bread wheat. *Plant Journal* 90: 1007-1013.

911 Moore, G., K. M. Devos, Z. Wang and M. D. Gale, 1995 Cereal genome evolution - grasses, line  
912 up and form a circle. *Current Biology* 5: 737-739.

913 Newman, J., P. Concannon, M. Tardaguila, A. Conesa and L. McIntyre, 2018 Event Analysis:  
914 Using Transcript Events To Improve Estimates of Abundance in RNA-seq Data. *G3-*  
915 *Genes Genomes Genetics* 8: 2923-2940.

916 Nielsen, H., 2017 Predicting Secretory Proteins with SignalP, pp. 59-73 in *Protein Function*  
917 *Prediction: Methods and Protocols*, edited by D. Kihara. Springer New York, New York,  
918 NY.

919 Nielsen, H., J. Engelbrecht, S. Brunak and G. von Heijne, 1997 Identification of prokaryotic and  
920 eukaryotic signal peptides and prediction of their cleavage sites. *Protein Eng* 10: 1-6.

921 Puckette, M. C., Y. Tang and R. Mahalingam, 2008 Transcriptomic changes induced by acute  
922 ozone in resistant and sensitive *Medicago truncatula* accessions. *BMC Plant Biol* 8: 46.

923 Qian, L. W., K. Voss-Fels, Y. X. Cui, H. U. Jan, B. Samans *et al.*, 2016 Deletion of a Stay-Green  
924 Gene Associates with Adaptive Selection in *Brassica napus*. *Molecular Plant* 9: 1559-  
925 1569.

926 Quevillon, E., V. Silventoinen, S. Pillai, N. Harte, N. Mulder *et al.*, 2005 InterProScan: protein  
927 domains identifier. *Nucleic acids research* 33: W116-W120.

928 Rao, M. V., H. Lee and K. R. Davis, 2002 Ozone-induced ethylene production is dependent on  
929 salicylic acid, and both salicylic acid and ethylene act in concert to regulate ozone-  
930 induced cell death. *Plant Journal* 32: 447-456.

931 Ricci, W. A., Z. F. Lu, L. X. Ji, A. P. Marand, C. L. Ethridge *et al.*, 2019 Widespread long-range  
932 cis-regulatory elements in the maize genome. *Nature Plants* 5: 1237-1249.

933 Roessler, K., A. Bousios, E. Meca and B. S. Gaut, 2018 Modeling interactions between  
934 transposable elements and the plant epigenetic response: a surprising reliance on element  
935 retention. *Genome Biol Evol* 10: 803-815.

936 Salika, R., and J. Riffat, 2021 Abiotic stress responses in maize: a review. *Acta Physiologiae*  
937 *Plantarum* 43.

938 Schnable, P. S., D. Ware, R. S. Fulton, J. C. Stein, F. S. Wei *et al.*, 2009 The B73 Maize  
939 Genome: Complexity, Diversity, and Dynamics. *Science* 326: 1112-1115.

940 Sonnhammer, E. L., G. von Heijne and A. Krogh, 1998 A hidden Markov model for predicting  
941 transmembrane helices in protein sequences. *Proc Int Conf Intell Syst Mol Biol* 6: 175-  
942 182.

943 Springer, N. M., K. Ying, Y. Fu, T. Ji, C.-T. Yeh *et al.*, 2009 Maize Inbreds Exhibit High Levels  
944 of Copy Number Variation (CNV) and Presence/Absence Variation (PAV) in Genome  
945 Content. *Plos Genetics* 5.

946 Steige, K. A., B. Laenen, J. Reimegard, D. G. Scofield and T. Slotte, 2017 Genomic analysis  
947 reveals major determinants of cis-regulatory variation in *Capsella grandiflora*.  
948 *Proceedings of the National Academy of Sciences of the United States of America* 114:  
949 1087-1092.

950 Stupar, R. M., and N. M. Springer, 2006 Cis-transcriptional variation in maize inbred lines B73  
951 and Mo17 leads to additive expression patterns in the F1 hybrid. *Genetics* 173: 2199-  
952 2210.

953 Subramanian, A., P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert *et al.*, 2005 Gene set  
954 enrichment analysis: A knowledge-based approach for interpreting genome-wide  
955 expression profiles. *Proceedings of the National Academy of Sciences* 102: 15545-  
956 15550.

957 Sun, S., Y. Zhou, J. Chen, J. Shi, H. Zhao *et al.*, 2018 Extensive intraspecific gene order and  
958 gene structural variations between Mo17 and other maize genomes. *Nature Genetics* 50:  
959 1289-+.

960 Swanson-Wagner, R., S. Eichten, S. Kumari, P. Tiffin, J. Stein *et al.*, 2010 Pervasive gene  
961 content variation and copy number variation in maize and its undomesticated progenitor.  
962 *Genome Research* 20: 1689-1699.

963 Tardaguila, M., L. de la Fuente, C. Marti, C. Pereira, F. J. Pardo-Palacios *et al.*, 2018 SQANTI:  
964 extensive characterization of long-read transcript sequences for quality control in full-  
965 length transcriptome identification and quantification. *Genome Res.*

966 Thatcher, S. R., O. N. Danilevskaya, X. Meng, M. Beatty, G. Zastrow-Hayes *et al.*, 2016  
967 Genome-Wide Analysis of Alternative Splicing during Development and Drought Stress  
968 in Maize. *Plant Physiology* 170: 586-599.

969 Verhoeven, K. J. F., E. H. Verbon, T. P. van Gurp, C. Oplaat, J. F. de Carvalho *et al.*, 2018  
970 Intergenerational environmental effects: functional signals in offspring transcriptomes  
971 and metabolomes after parental jasmonic acid treatment in apomictic dandelion. *New*  
972 *Phytologist* 217: 871-882.

973 Waldeck, N., K. Burkey, T. Carter, D. Dickey, Q. Song *et al.*, 2017 RNA-Seq study reveals  
974 genetic responses of diverse wild soybean accessions to increased ozone levels. BMC  
975 Genomics 18: 498.

976 Wang, B., M. Regulski, E. Tseng, A. Olson, S. Goodwin *et al.*, 2018a A comparative  
977 transcriptional landscape of maize and sorghum obtained by single-molecule sequencing.  
978 Genome Res 28: 921-932.

979 Wang, W. S., R. Mauleon, Z. Q. Hu, D. Chebotarov, S. S. Tai *et al.*, 2018b Genomic variation in  
980 3,010 diverse accessions of Asian cultivated rice. Nature 557: 43-+.

981 Wang, X. C., M. Yang, D. Q. Ren, W. Terzaghi, X. W. Deng *et al.*, 2019 Cis-regulated  
982 alternative splicing divergence and its potential contribution to environmental responses  
983 in Arabidopsis. Plant Journal 97: 555-570.

984 Wedow, J. M., E. A. Ainsworth and S. Li, 2021a Plant biochemistry influences tropospheric  
985 ozone formation, destruction, deposition, and response. Trends in biochemical sciences.

986 Wedow, J. M., C. H. Burroughs, L. R. Acosta, A. D. B. Leakey and E. A. Ainsworth, 2021b  
987 Age-dependent increase in alpha-tocopherol and phytosterols in maize leaves exposed to  
988 elevated ozone pollution. Plant Direct 5.

989 Weirather, J. L., M. de Cesare, Y. Wang, P. Piazza, V. Sebastiano *et al.*, 2017 Comprehensive  
990 comparison of Pacific Biosciences and Oxford Nanopore Technologies and their  
991 applications to transcriptome analysis. F1000Research 6: 100-100.

992 Wittkopp, P. J., 2007 Variable gene expression in eukaryotes: a network perspective. Journal of  
993 Experimental Biology 210: 1567-1575.

994 Woodhouse, M. R., E. K. Cannon, J. L. Portwood, L. C. Harper, J. M. Gardiner *et al.*, 2021 A  
995 pan-genomic approach to genome databases using maize as a model system. BMC Plant  
996 Biology 21.

997 Xu, E., L. Vaahtera, H. Hõrak, D. K. Hinch, A. G. Heyer *et al.*, 2015 Quantitative trait loci  
998 mapping and transcriptome analysis reveal candidate genes regulating the response to  
999 ozone in Arabidopsis thaliana. Plant Cell Environ 38: 1418-1433.

1000 Yang, N., X. Xu, R. Wang, W. Peng, L. Cai *et al.*, 2017a Contributions of Zea mays subspecies  
1001 mexicana haplotypes to modern maize. Nature Communications 8.

1002 Yang, Z., R. Lu, Z. Dai, A. Yan, Q. Tang *et al.*, 2017b Salt-stress response mechanisms using de  
1003 novo transcriptome sequencing of salt-tolerant and sensitive corchorus spp. genotypes.  
1004 Genes 8: 226.

1005 Yao, W., G. W. Li, H. Zhao, G. W. Wang, X. M. Lian *et al.*, 2015 Exploring the rice dispensable  
1006 genome using a metagenome-like assembly strategy. Genome Biology 16.

1007 Yendrek, C., G. Erice, C. Montes, T. Tomaz, C. Sorgini *et al.*, 2017a Elevated ozone reduces  
1008 photosynthetic carbon gain by accelerating leaf senescence of inbred and hybrid maize in  
1009 a genotype-specific manner. *Plant Cell and Environment* 40: 3088-3100.

1010 Yendrek, C. R., T. Tomaz, C. M. Montes, Y. Y. Cao, A. M. Morse *et al.*, 2017b High-  
1011 Throughput Phenotyping of Maize Leaf Physiological and Biochemical Traits Using  
1012 Hyperspectral Reflectance. *Plant Physiology* 173: 614-626.

1013 Yu, J. M., J. B. Holland, M. D. McMullen and E. S. Buckler, 2008 Genetic design and statistical  
1014 power of nested association mapping in maize. *Genetics* 178: 539-551.

1015 Zhao, Q., Q. Feng, H. Y. Lu, Y. Li, A. Wang *et al.*, 2018 Pan-genome analysis highlights the  
1016 extent of genomic variation in cultivated and wild rice. *Nature Genetics* 50: 279-+.

1017 Zou, C., K. Sun, J. D. Mackaluso, A. E. Seddon, R. Jin *et al.*, 2011 Cis-regulatory code of stress-  
1018 responsive transcription in *Arabidopsis thaliana*. *Proceedings of the National Academy of*  
1019 *Sciences of the United States of America* 108: 14992-14997.

1020

1021

1022