

UNIVERSITA' DEGLI STUDI DI  
MILANO-BICOCCA



SCUOLA DI SCIENZE MATEMATICHE FISICHE E NATURALI  
CORSO DI LAUREA TRIENNALE IN FISICA

**PROBING OPTICAL AND RADIO-LOUD AGN FRACTIONS : A  
COMPARATIVE ANALYSIS BETWEEN BCGs AND NON-BCGs  
SAMPLES at  $z < 0.1$**

*Candidate:*  
ANDREA MACCARINELLI

*Supervisor:*  
Prof. SEBASTIANO CANTALUPO

*Co-supervisors:*  
Dott. ANDREA TRAVASCIO

ANNO ACCADEMICO 2022/2023

$\emptyset$

# Abstract

Brightest cluster galaxies (BCGs) are the most massive and luminous galaxies located near the center of relaxed, virialized, and undisturbed galaxy clusters in the local Universe ([12, 16]. According to several observational studies ([6], [13]), these objects experience a special formation process differing from general galaxy evolution. Current theoretical models (e.g., [4, 3]) predict that dry mergers are the dominant mechanisms responsible for their mass assembly at  $z < 1$ . These objects are often observed to host a supermassive black hole (SMBH) in their center ([11]). The process of matter accretion into these SMBHs may release a large amount of energy, resulting in Active Galactic Nuclei (AGN). There are two primary modes in which SMBH accretion can occur: the so-called 'quasar mode' and the 'radio mode'. The quasar mode involves a high accretion rate of the SMBH via an optically-thick and geometrically-thin disk, with most of the energy being released in the form of radiation. In a radio mode scenario, the SMBH accretion of hotter gas occurs at a low rate in an optically-thin and geometrically-thick disk configuration, releasing energy in the form of relativistic particles, i.e. radio jets. The latter is typically observed in BCGs.

The evolutionary processes of BCGs are still not fully understood, and there are no specific studies comparing the frequency of different types of AGN in BCGs with respect to other types of galaxies (e.g., [5]).

The main scientific question driving the project of my thesis is to understand whether the different evolution of BCGs, along with their "special" environment, affects the accretion of SMBHs in their centers compared to other types of galaxies in the local universe, at  $z < 0.1$ . To address this question, I analyzed a sample of BCGs derived from the combination of the Sloan Digital Sky Survey Data Release 7 (SDSS DR7; described in [1]) and the C4 BCGs catalogue produced by [8]. For this BCG sample at  $z=0.02-0.1$ , I use the fluxes of optical lines ( $H\alpha$ , [OIII],  $H\beta$ , [NII], and [SII] doublets)

obtained by the MPA-JHU team using methods described in [9] to obtain a census of optical AGN through the NII and SII BPT diagnostic diagrams. Additionally, by cross-matching the previous catalogs with a collection derived from NVSS and FIRST radio surveys, as presented in [2], I identify the BCGs exhibiting radio loud emission, implying the presence of radio jets.

Following this classification, I finally estimate the fraction of BCGs classified as Optical and Radio Loud AGN. Subsequently, I derive the same fractions for the non-BCG selected galaxies according to the cross-match with the C4 catalogue ([8]).

The analyses reveal that BCGs exhibit a higher fraction of optical AGN  $\sim 50\%$  compared to the non-BCG sample, which shows a percentage of  $\sim 21\%$ , consistent with results found by [14]. Simultaneously, the analysis of Radio Loud emissions indicates that BCGs are more inclined to host Radio Loud Activity, with a fraction of  $\sim 12\%$ . This fraction is found to be 20 times higher than the  $\sim 0.6\%$  observed in the non-BCG sample of selected galaxies.

In conclusion, these results strongly support the idea that BCGs are more likely to exhibit typical ionization due to AGN activity and radio emission. This implies that their privileged position supports frequent accretion of SMBHs in both accretion modes. Previous studies, such as [18, 10, 15], have already shown a prevalence of radio AGN among the BCG population in the local Universe. On the other hand, the larger fraction of BCGs selected as optical AGN compared to normal galaxies is intriguing. Future studies will aim to test the results obtained with this sample and understand if this higher fraction is specifically driven by differences in properties between BCGs and non-BCGs, such as mass, star formation rate, metallicity, and kinematics.

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	The Brightest Cluster Galaxies . . . . .	3
1.2	The Active Galactic Nuclei Feedback . . . . .	3
1.3	The Aim of this thesis . . . . .	3
<b>2</b>	<b>Methods</b>	<b>5</b>
2.1	Data Description . . . . .	5
2.2	Data Analysis . . . . .	5
2.2.1	Optical analysis . . . . .	5
2.2.2	Radio Analysis . . . . .	5
<b>3</b>	<b>Results</b>	<b>7</b>
3.1	The Prevalence of AGN activity in BCGs . . . . .	7
3.2	The Radio activity of BCGs . . . . .	7
	<b>Conclusions</b>	<b>9</b>



# Chapter 1

## Introduction

### 1.1 The Active Galactic Nuclei Feedback

### 1.2 The Brightest Cluster Galaxies

One of the main branches of cosmic structure formation is covered by the Hierarchical model in which galaxies grow in both stellar and mass magnitude, by accreting the surrounding matter.

One of the most extreme examples in this context involves the study of Brightest Cluster Galaxies (BCGs), a unique class of galaxies typically elliptical, often situated at the center and typically standing out as the most luminous and massive objects within the entire cluster.( e.g. [17] ).

Considering the environment, also their evolution is proven via observational studies e.g. [13], to be slightly different to normal galactic ones, resulting in the current main model assumes that BCGs mass assembly is dominated by dry mergers [4, 3].

Thanks to their exclusive environment in which they live, consisting in a BCGs often host a Super Massive Black Hole ( SMBH ) ([11]) resulting in Active Galactic Nuclei ( AGN ) activity such as Radio Jets that feedback into the Intra-Cluster Medium (ICM)[7].

As seen in previous studies ( e.g. [15] ) BCGs more likely host Radio Loud Emission than in other type of galaxies, as a result of the influence of the environment other than its nuclei.

### 1.3 The Aim of this thesis





# Chapter 2

## Methods

### 2.1 Data Description

A description of the Data Samples Utilized, and its peculiarities, starting from The SDSS DR7 description, and following the work provided by the MPA-JHU team.

### 2.2 Data Analysis

How we analyzed the data, where do we defined the fractions calculated ?

#### 2.2.1 Optical analysis

Describe each population of the BPT diagram and its peculiarities, before starting the description of how we collected and processed the data to finally create BPT diagrams . Finally there's the need to explain how we interpreted the diagrams to finally calculate the fractions ( in this case you need to explain which algorithm has been chosen and why )

#### 2.2.2 Radio Analysis

I certainly need to focus on how i have chosen the range of identification, to recognize which element of sdss was effectively recognized as a Radioloud.

Following to this it is necessary to explain how we selected the elements in which define the fractions, by selecting elements of SDSS nearby the location in which best et al finds mostly its radio-emitting elements!



# Chapter 3

## Results

A complete description of the results produced !!

### 3.1 The Prevalence of AGN activity in BCGs

Describe how the evidence of the data analysis we conducted leads to the description of BCGs in a confrontation to non-BCGs, and in how different is still from a pure optical view the response of the AGN feedback.

Present absolutely the Table with all of the percentage ( and the counting ) values of the populations of both BPT NII and BPT SII diagnostics and introduce a confrontation with also the results found in Vitale et AL and others.

### 3.2 The Radio activity of BCGs

Explain the evidences found in the analysis and possible ways to improve results obtained.

There's certainly also the need to add a discussion on how it is possible that we ignore some of the BCGs in the counting, issue that should be resolved by choosing a more complete source of data regarding Radio Emitters galaxies.



# Conclusions

The final chapter of the thesis is a summary of the work done. Therefore, in its first part it resembles much the introduction, adding to it the actual result of the work, its future evolution and prospects, in the view of the writer.



# Bibliography

- [1] Kevork N. Abazajian and Adelman-McCarthy. The Seventh Data Release of the Sloan Digital Sky Survey. *Astrophysical Journal Supplement Series*, 182(2):543–558, June 2009. doi: 10.1088/0067-0049/182/2/543.
- [2] P. N. Best, G. Kauffmann, T. M. Heckman, and Ž. Ivezić. A sample of radio-loud active galactic nuclei in the Sloan Digital Sky Survey. *Monthly Notices of the Royal Astronomical Society*, 362(1):9–24, September 2005. doi: 10.1111/j.1365-2966.2005.09283.x.
- [3] Kevin C. Cooke, Jeyhan S. Kartaltepe, K. D. Tyler, Behnam Darvish, Caitlin M. Casey, Olivier Le Fèvre, Mara Salvato, and Nicholas Scoville. Stellar Mass Growth of Brightest Cluster Galaxy Progenitors in COSMOS Since  $z \sim 3$ . *The Astrophysical Journal*, 881(2):150, August 2019. doi: 10.3847/1538-4357/ab30c9.
- [4] Gabriella De Lucia and Jérémy Blaizot. The hierarchical formation of the brightest cluster galaxies. *Monthly Notices of the Royal Astronomical Society*, 375(1):2–14, February 2007. doi: 10.1111/j.1365-2966.2006.11287.x.
- [5] S. Fisek, S. Alis, E. K. Ülgen, and F. K. Yelkenci. AGN Activity in Brightest Cluster Galaxies (BCGs). *Communications of the Byurakan Astrophysical Observatory*, 66(2):153–158, December 2019. doi: 10.52526/25792776-2019.66.2-153.
- [6] Daniël N. Groenewald, Rosalind E. Skelton, David G. Gilbank, and S. Ilani Loubser. The close pair fraction of BCGs since  $z = 0.5$ : major mergers dominate recent BCG stellar mass growth. *Monthly Notices of the Royal Astronomical Society*, 467(4):4101–4117, June 2017. doi: 10.1093/mnras/stx340.

- 
- [7] B. R. McNamara and P. E. J. Nulsen. Heating Hot Atmospheres with Active Galactic Nuclei. *Annual Review of Astronomy and Astrophysics*, 45(1):117–175, September 2007. doi: 10.1146/an-nurev.astro.45.051806.110625.
- [8] Christopher J. Miller, Robert C. Nichol, Daniel Reichart, Risa H. Wechsler, August E. Evrard, James Annis, Timothy A. McKay, Neta A. Bahcall, Mariangela Bernardi, Hans Boehringer, Andrew J. Connolly, Tomotsugu Goto, Alexie Kniazev, Donald Lamb, Marc Postman, Donald P. Schneider, Ravi K. Sheth, and Wolfgang Voges. The C4 Clustering Algorithm: Clusters of Galaxies in the Sloan Digital Sky Survey. *The Astronomical Journal*, 130(3):968–1001, September 2005. doi: 10.1086/431357.
- [9] James E. O’Donnell. R v-dependent Optical and Near-Ultraviolet Extinction. *The Astrophysical Journal*, 422:158, February 1994. doi: 10.1086/173713.
- [10] P. Oliva-Altamirano, S. Brough, C. Lidman, W. J. Couch, A. M. Hopkins, M. Colless, E. Taylor, A. S. G. Robotham, M. L. P. Gunawardhana, T. Ponman, I. Baldry, A. E. Bauer, J. Bland-Hawthorn, M. Cluver, E. Cameron, C. J. Conselice, S. Driver, A. C. Edge, A. W. Graham, E. van Kampen, M. A. Lara-López, J. Liske, A. R. López-Sánchez, J. Loveday, S. Mahajan, J. Peacock, S. Phillipps, K. A. Pimbblet, and R. G. Sharp. Galaxy And Mass Assembly (GAMA): testing galaxy formation models through the most massive galaxies in the Universe. *Monthly Notices of the Royal Astronomical Society*, 440(1):762–775, May 2014. doi: 10.1093/mnras/stu277.
- [11] D. A. Rafferty, B. R. McNamara, P. E. J. Nulsen, and M. W. Wise. The Feedback-regulated Growth of Black Holes and Bulges through Gas Accretion and Starbursts in Cluster Central Dominant Galaxies. *The Astrophysical Journal*, 652(1):216–231, November 2006. doi: 10.1086/507672.
- [12] A. Sandage, J. Kristian, and J. A. Westphal. The extension of the Hubble diagram. I. New redshifts and BVR photometry of remote cluster galaxies, and an improved richness correction. *The Astrophysical Journal*, 205:688–695, May 1976. doi: 10.1086/154324.



- 
- [13] A. Travascio, A. Bongiorno, P. Tozzi, R. Fassbender, F. De Gasperin, V. F. Cardone, L. Zappacosta, G. Vietri, E. Merlin, M. Bischetti, E. Piconcelli, F. Duras, F. Fiore, N. Menci, P. Mazzotta, and A. Nastasi. Multiple AGN activity during the BCG assembly of XDCPJ0044.0-2033 at  $z \sim 1.6$ . *Monthly Notices of the Royal Astronomical Society*, 498(2):2719–2733, October 2020. doi: 10.1093/mnras/staa2495.
- [14] M. Vitale, J. Zuther, M. García-Marín, A. Eckart, M. Bremer, M. Valencia-S., and A. Zensus. Classifying radio emitters from the Sloan Digital Sky Survey. Spectroscopy and diagnostics. *Astronomy & Astrophysics*, 546:A17, October 2012. doi: 10.1051/0004-6361/201219290.
- [15] Anja Von Der Linden, Philip N. Best, Guinevere Kauffmann, and Simon D. M. White. How special are brightest group and cluster galaxies? *Monthly Notices of the Royal Astronomical Society*, 379(3):867–893, August 2007. doi: 10.1111/j.1365-2966.2007.11940.x.
- [16] Anja von der Linden, Vivienne Wild, Guinevere Kauffmann, Simon D. M. White, and Simone Weinmann. Star formation and AGN activity in SDSS cluster galaxies. *Monthly Notices of the Royal Astronomical Society*, 404(3):1231–1246, May 2010. doi: 10.1111/j.1365-2966.2010.16375.x.
- [17] Z. L. Wen and J. L. Han. Dependence of the bright end of composite galaxy luminosity functions on cluster dynamical states. *Monthly Notices of the Royal Astronomical Society*, 448(1):2–8, March 2015. doi: 10.1093/mnras/stu2722.
- [18] Z. S. Yuan, J. L. Han, and Z. L. Wen. Radio luminosity function of brightest cluster galaxies. *Monthly Notices of the Royal Astronomical Society*, 460(4):3669–3678, August 2016. doi: 10.1093/mnras/stw1125.