



McStas

—
n—→



2021
HighNESS
Virtual
McStas
School

McStas Introduction and general concepts






McStas



2021
HighNESS
Virtual
McStas
School



Agenda

- A (very) brief introduction to McStas
- Components of neutron instruments
- How McStas works under the hood
- Components and instruments
- A demo



Neutron

The **quark** structure of the neutron. (The color assignment of individual quarks is not important; what is important is that all three colors are present.)

Classification	Baryon
Composition	1 up quark, 2 down quarks
Statistics	Fermionic
Interactions	Gravity, Weak, Strong, Electromagnetic
Symbol	$n^- n^0 N^0$
Antiparticle	Antineutron
Theorized	Ernest Rutherford ^{[1][2]} (1920)
Discovered	James Chadwick ^[1] (1932)
Mass	$1.674\,927\,351(74) \times 10^{-27} \text{ kg}$ $939.565\,378(21) \text{ MeV}/c^2$ ^[3] $1.008\,664\,916\,00(43) \text{ u}$ ^[3]
Mean lifetime	$881.5(15) \text{ s}$ (free)
Electric charge	0 e
Electric dipole moment	$< 2.9 \times 10^{-26} \text{ e-cm}$
Electric polarizability	$1.16(15) \times 10^{-3} \text{ fm}^3$

Life time:

$$\tau_{1/2} = 890 \text{ s}$$

Mass:

$$m = 1.675 \times 10^{-27} \text{ kg}$$

Charge:

$$Q = 0$$

Spin:

$$s = \hbar/2$$

Magnetic moment:

$$\mu/\mu_n = -1.913$$

$$E = \frac{1}{2}mv^2 = \frac{\hbar^2 k^2}{2m} \quad \lambda = 2\pi/k$$

$$E = 81.81 \cdot \lambda^{-2} = 2.07 \cdot k^2 = 5.23 \cdot v^2$$

Subatomic particle discovered by Sir James Chadwick in 1932



	Energy	Wavelength	n-Wavevector	Velocity	Frequency
cold neutrons:	$E = 1 \text{ meV}$ $E = 5 \text{ meV}$	$\lambda = 9.0446 \text{ \AA}$ $\lambda = 4.0449 \text{ \AA}$	$k = 0.6947 \text{ 1/\AA}$ $k = 1.5534 \text{ 1/\AA}$	$v = 437 \text{ m/s}$ $v = 978 \text{ m/s}$	$v = 0.2418 \text{ THz}$ $v = 1.2090 \text{ THz}$
thermal neutrons:	$E = 25 \text{ meV}$ $E = 50 \text{ meV}$	$\lambda = 1.8089 \text{ \AA}$ $\lambda = 1.2791 \text{ \AA}$	$k = 3.4734 \text{ 1/\AA}$ $k = 4.9122 \text{ 1/\AA}$	$v = 2187 \text{ m/s}$ $v = 3093 \text{ m/s}$	$v = 6.045 \text{ THz}$ $v = 12.090 \text{ THz}$







2021
HighNESS
Virtual
McStas
School

Monte Carlo techniques

- Los Alamos has since then developed and perfected many different monte carlo codes leading to what is today known as the codes MCNP5 and MCNPX
- State of the art is MCNP6 that features numerous (even exotic) particles
- MCNP was originally Monte Carlo Neutron Photon, later N-Particle
- Mainly used for high-energy particle descriptions in weapons, power reactors and routinely used for estimating dose rates and needed shielding
- Not much focus on crystalline / ordered material and coherent scattering of neutrons due to the focus on high energies



Examples of Monte Carlo programs



McStas



2021
HighNESS
Virtual
McStas
School



Each time physics takes place (scattering, absorption, ...) random choices are made.

Light ray-tracing: PoV-RAY and others ...

*Nuclear reactor simulations etc. (general particle transport):
MCNP, Tripoli, GEANT4, FLUKA*

Neutron Monte Carlo ray-tracing propagation:

McStas <www.mcstas.org>, Vitess, Restrax, NISP, IDEAS, McVine

Neutrons are described as (\mathbf{r} , \mathbf{v} , \mathbf{s} , t), and are transported along instrument models.

Propagation simply uses Newton rules, incl. gravitation.

X-ray tracing

Shadow, McXtrace, RAY, ...




HighNess

McStas




2021
HighNESS
Virtual
McStas
School


In the big picture, McStas is this...



- Classical Newtonian mechanics, i.e.
- (independent, particles though...)



and




HighNess

McStas



2021
HighNESS
Virtual
McStas
School

In the big picture, McStas is this...



The instrument defines our “lab coordinate system”






McStas



2021
HighNESS
Virtual
McStas
School

In the big picture, McStas is this...



The instrument defines our “lab coordinate system”

The components define devices or features available in our instrument






McStas



2021
HighNESS
Virtual
McStas
School

In the big picture, McStas is this...



The instrument defines our “lab coordinate system”

The components define devices or features available in our instrument - they have different function




HighNess

McStas



2021
HighNESS
Virtual
McStas
School

In the big picture, McStas is this...



The instrument defines our “lab coordinate system”

The components define devices or features available in our instrument - they have different function

Neutron particles are passed on from one component to the next, changing state under way




HighNess

McStas



2021
HighNESS
Virtual
McStas
School

McStas is a Monte Carlo ray-tracer






McStas




2021
HighNESS
Virtual
McStas
School



Elements of Monte-Carlo raytracing

- Instrument Monte Carlo methods implement coherent scattering effects
- Uses deterministic propagation where this can be done
- Uses Monte Carlo sampling of “complicated” distributions and stochastic processes and multiple outcomes with known probabilities are involved
 - I.e. inside scattering matter
- Uses the particle-wave duality of the neutron to switch back and forward between deterministic ray tracing and Monte Carlo approach



- Result: A realistic and efficient transport of neutrons in the thermal and cold range






McStas



2021
HighNESS
Virtual
McStas
School

Components of neutron instruments




HighNess

McStas



2021
HighNESS
Virtual
McStas
School

Components of neutron instruments




In McStas the moderator is the “source”






2021
HighNESS
Virtual
McStas
School


Components of neutron instruments



In McStas the moderator is the “source”





Components of neutron instruments




The sample:


Crystalline, powders, liquids, micelles, structures to image, inelastic features like phonons...



In McStas the moderator is the “source”



Components of neutron instruments







The sample:

Crystalline, powders, liquids, micelles, structures to image, inelastic features like phonons...

In McStas the moderator is the “source”

Detectors are “monitors” in McStas. Mostly they act as “perfect probes” and can be positioned thought your instrument gathering 1D/2D/ event lists...





2021
HighNESS
Virtual
McStas
School

McStas Introduction

- Flexible, general simulation utility for neutron scattering experiments.
- Original design for **Monte carlo Simulation of triple axis spectrometers**
- Developed at DTU Physics, ILL, PSI, Uni CPH, ESS DMSC
- V. 1.0 by K Nielsen & K Lefmann (1998) RISØ
- Currently 2.5+1 people full time plus students




GNU GPL license
Open Source

The screenshot shows the McStas homepage. The header features the McStas logo and navigation links for Getting Started, Latest Headlines, About McStas, Documentation, Download, Mailing list, and Workshops/conferences. The main content area is titled "McStas - A neutron ray-trace simulation package". It includes a plot showing simulated scattering from a hollow cylinder vanadium sample, a "Recent news" section with entries for May 18th, 2009, and April 14th, 2009, and a sidebar with links for Developments, Links, Report bugs, CVS, and McStas Ubuntu live-dvd.

Project website at
<http://www.mcstas.org>

mcstas-users@mcstas.org mailinglist





McStas Introduction

Main Page – McXtraceWiki

Most Visited Getting Started Latest Headlines http://www.google... Geekblog Nyheder http://www.google... dr.dk open streami... Log in / create account

article discussion edit history

Main Page

McXtrace

[edit]

McXtrace - Monte Carlo Xray ray-tracing is a joint venture by

Risø DTU DTU ESRF JJ X-RAY Danish Science Design Engineering Production

Funding from NABIIT, DSF and the above parties.

Our code will be based on technology from

For information on our progress, please subscribe to our user mailinglist.

mailto:webmaster@mcxtrace.org


This page was last modified 13:15, 25 February 2009. This page has been accessed 2,049 times. Privacy policy About McXtraceWiki Disclaimers Powered By MediaWiki

- Synergy, knowledge transfer, shared infrastructure










Used in many places





What is McStas used for?

- Instrumentation
- Planning
- Construction
- Virtual experiments
- Data analysis
- Teaching
(KU, DTU)






2021
HighNESS
Virtual
McStas
School


Reliability - cross comparisons

- Much effort has gone into this
- Here: simulations vs. exp. at powder diffract. DMC, PSI
- The bottom line is
- McStas agree very well with other packages (NISP, Vitess, IDEAS, RESTRAX, ...)
- Experimental line shapes are within 5%
- All
- C



E. Farhi, P. Willendrup

des ε



P. Willendrup et al., Physica B, 386, (2006), 1032.






McStas



2021
HighNESS
Virtual
McStas
School




McStas overview

- Portable code (Unix/Linux/Mac/Windoze)
 - Ran on everything from iPhone to 1000+ node cluster!
- 'Component' files (~100) inserted from library
 - Sources
 - Optics
 - Samples
 - Monitors
 - If needed, write your own comps
- DSL + ISO-C code gen.

CPU's



+ NVIDIA GPU's







McStas




2021
HighNESS
Virtual
McStas
School




Under-the-hood / inner workings

- Domain-specific-language (DSL) based on compiler technology (LeX+Yacc)
 - Simple Instrument language  ISO C
- Component codes realizing beamline parts (including user contribs)
- Library of common functions for e.g.
 - I/O
 - Random numbers
 - Physical constants
 - Propagation
 - Precession in fields
 - ...





McStas




HighNess




2021
HighNESS
Virtual
McStas
School

Placing components - source

- One of the first components in your instrument is typically a source, which has a coordinate system like this....



- z is along neutron beam direction
- y is vertical
- x at an angle of 90° wrt. z,y



Right-handed
coordinate system






McStas




2021
HighNESS
Virtual
McStas
School



Placing components - source

- Often the source coordinate system coincides with the “lab” coordinate system, denoted ABSOLUTE in McStas language, i.e.



- COMPONENT Source = Source_simple(...)
AT (0,0,0) ABSOLUTE





McStas




2021
HighNESS
Virtual
McStas
School


Placing further components - RELATIVE

Placing further components is done by order of

1. Location, i.e



COMPONENT Source = Source_simple(...)
AT (0,0,0) ABSOLUTE




COMPONENT Guide = Guide(...)
AT (0,0,1) RELATIVE Source




Placing further components - RELATIVE

Placing further components is done by order of
2. Rotation, i.e





COMPONENT Source = Source_simple(...)
AT (0,0,0) ABSOLUTE



COMPONENT Guide = Guide(...)
AT (0,0,1) RELATIVE Source
ROTATED (0,30,0) RELATIVE Source

(Reference labels can also be PREVIOUS or PREVIOUS+1 etc.)

Components often have their origin at the centre of mass, i.e. for samples ... but not for neutron guides



Generally speaking, the component author can choose **the meaningful coordinate system for the given problem!**

- The McStas system takes care of the transformation between them....

Component geometries are typically simple objects... But some have polygon-description of the surface

HighNess

McStas

n



DTU
ESS
EUROPEAN SPALLATION SOURCE

2021
HighNESS
Virtual
McStas
School

2D



3D





McStas




2021
HighNESS
Virtual
McStas
School

Component classes

- Sources - these define MC starting conditions / “inject” neutrons to our simulation
- Optics - used to tailor properties of the neutron beam
 - Examples are mirrors, guides, choppers, collimators, slits, ...
- Samples - “matter” of some form
 - Powders, single crystals, liquids, micelles in solution, reflecting surfaces...
- Monitors - may probe the state of the neutron beam and store histograms / event lists
- Misc, obsolete
 - “Other stuff” and “Old stuff”

*Other
Stuff*





McStas



2021
HighNESS
Virtual
McStas
School


Component classes

- Sources - these define MC starting conditions / “inject” neutrons to our simulation
- Optics - used to tailor properties of the neutron beam
 - Examples are mirrors, guides, choppers, collimators, slits, ...
- Samples - “matter” of some form
 - Powders, single crystals, liquids, micelles in solution, reflecting surfaces
- Monitors - may probe
- Misc, obsolete
 - “Other stuff” and “Old stuff”




Common to all components:
They set, manipulate/interact with
or measure the state of the neutron (ray)

*Other
Stuff*



Neutron sources, i.e. moderators

- To first order emit uniformly into 4π steradian






McStas



2021
HighNESS
Virtual
McStas
School

Neutron sources, i.e. moderators




- Generally we are interested in the input to a single instrument, characterised by a certain solid angle Ω , often corresponding to a rectangle $xw \times yh$ at a distance $dist$ from the source

$xw \quad yh \quad dist$

E.g. the beamport?

Neutron sources, i.e. moderators

- The emission intensity into our chosen solid angle Ω can be a function of wavelength, time (pulsed sources) and possibly point of origin on the source surface



$$I(\lambda)$$

$$I(\lambda, t)$$

$$I(\lambda, t, \vec{r})$$

$$\Omega [n/s/str]$$

$$\Omega [n/s/str]$$

$$\Omega [n/s/str]$$


- The emission of particles into the solid angle Ω is in fact an integration and leads to a simulated “intensity” of $I_\Omega [n/s]$
- In McStas, that integrated intensity is partitioned over a given set of particle rays referred to as **ncount**, -n or --ncount
- The default **ncount** is 1e6 rays



2021
HighNESS
Virtual
McStas
School

Neutron sources, i.e. moderators

- Our neutron rays are emitted randomly, sampling source “spectrum”, i.e. wavelength, time and area and all variables of the
- assigning neutron weights such that



$$I_{\Omega}(\lambda, t, \vec{r}) [n/s]$$

p

$$\sum_{j=1}^{\text{ncount}} p_j = \int_{d\lambda, dt, d\vec{r}} I_{\Omega}(\lambda, t, \vec{r})$$





2021
HighNESS
Virtual
McStas
School

Neutron rays in McStas - what are they?

- Defining the neutron starting conditions imply setting:
 - The **starting point** on the surface, i.e. \vec{r} (in the code variables x, y, z)
 - The **direction** into Ω and our λ/E_{kin} (in the code variables vx, vy, vz)
 - The **starting time** (in the code the variable t)
 - The initial **intensity** / weight of the neutron ray (in the code the variable p)
 - If needed the initial **polarisation** (in the code the variables sx, sy, sz)

Neutron **ray** in McStas:

Location x, y, z

Velocity vx, vy, vz

Time t

Polarisation. sx, sy, sz


Intensity p



Neutron (ray)-matter interaction 1: reflecting surface


- 1 starting situation

Neutron ray



$$(x, y, z, vx, vy, vz, t, sx, sy, sz, p)_0$$


Neutron (ray)-matter interaction 1: reflecting surface

- 2. Propagate to the mirror surface




McStas
 n →
DTU
ESS
EUROPEAN SPALLATION SOURCE
2021
HighNESS
Virtual
McStas
School




Neutron (ray)-matter interaction 1: reflecting surface

- 3. Checks (are we on surface, what is probability of reflection etc.)



2021
HighNESS
Virtual
McStas
School


$(x, y, z, vx, vy, vz, t, sx, sy, sz, p)_0$



Neutron (ray)-matter interaction 1: reflecting surface


- 4. Reflect

$(x, y, z, vx, vy, vz, t, sx, sy, sz, p)_0$



Weight of final ray is adjusted according to reflectivity, see next slide


Parametrisation of reflectivity on mirrors etc.




$$V = \frac{2\pi\hbar^2}{m} b N \quad \sin\theta < \sqrt{\frac{mV}{2\pi^2\hbar^2}} \lambda$$

$$m = \frac{\theta_{mirror}}{\theta_{Ni}}$$

$$R_0 \cdot \left(1 - \frac{\tanh(Q - mQ_c)}{W}\right) \cdot (1 - \alpha(Q - Q_c))$$





(i.e. Q is calculated for given neutron, reflectivity encoded in **changed p value**)



absorbed, transmitted, or scattered

Neutron (ray)-matter interaction in General





For a **non-thin** sample the probabilities for **absorption**, **transmission** or **scattering** are given by

$$p_A = (1 - e^{-\Sigma_T t})(\Sigma_A / \Sigma_T)$$

$$p_S = (1 - e^{-\Sigma_T t})(\Sigma_S / \Sigma_T)$$

$$p_T = 1 - p_S - p_A = e^{-\Sigma_T t}$$

$$\Sigma_* = \rho \sigma_*$$

$\mathbf{t} =$ sample thickness


macroscopic cross section [cm⁻¹]

number density [atoms/cm³]

microscopic cross section [barn/atom]
1 barn = 10⁻²⁴cm²




Samples/Matter interaction in General in McStas



transmitted+absorption, or scattered+absorption

Transport of weight through the instrument...



p_0

p_j

p_n

$$p_j = w_j p_{j-1}$$

$$p_j = p_0 \prod_{k=1}^j w_k$$


The weight multiplier of the j 'th component, w_j , is calculated by the probability rule $f_{MC,b}w_j = P_b$ where P_b is the physical probability for the event "b", and $f_{MC,b}$ is the probability that the Monte Carlo simulation selects this event.

In case of "branching", i.e. multiple outcomes, it is clear that

$$\sum_b f_{MC,b} = 1$$



To first order, McStas is linear and follows sequence of components in your file...



Starting at the source






McStas



2021
HighNESS
Virtual
McStas
School

To first order, McStas is linear and follows sequence of components in your file...



Moving to first comp in the list






McStas



2021
HighNESS
Virtual
McStas
School

To first order, McStas is linear and follows sequence of components in your file...



Moving to 3rd comp in list requires “moving back in time”.


Default behavior is to ABSORB this type of neutron.

For monitors use `restore_neutron=1` in this case.

For homegrown comps use `ALLOW_BACKPROP` macro.



To first order, McStas is linear and follows sequence of components in your file...



The order of components is important,
and in general overlaps should be avoided!

Moving to 3rd comp in list requires “moving back in time”.

Default behavior is to ABSORB this type of neutron.

For monitors use `restore_neutron=1` in this case.

For homegrown comps use `ALLOW_BACKPROP` macro.






McStas



2021
HighNESS
Virtual
McStas
School



Instrument file

```
DEFINE INSTRUMENT My_Instrument(DIST=10)

/* Here comes the TRACE section, where the actual      */
/* instrument is defined as a sequence of components.  */
TRACE

/* The Arm() class component defines reference points and orientations */
/* in 3D space.                                                       */
COMPONENT Origin = Arm()
    AT (0,0,0) ABSOLUTE

COMPONENT Source = Source_simple(
    radius = 0.1, dist = 10, xw = 0.1, yh = 0.1, E0 = 5, dE = 1)
    AT (0, 0, 0) RELATIVE Origin

COMPONENT Emon = E_monitor(
    filename = "Emon.dat", xmin = -0.1, xmax = 0.1, ymin = -0.1,
    ymax = 0.1, Emin = 0, Emax = 10)
    AT (0, 0, DIST) RELATIVE Origin

COMPONENT PSD = PSD_monitor(
    nx = 128, ny = 128, filename = "PSD.dat", xmin = -0.1,
    xmax = 0.1, ymin = -0.1, ymax = 0.1)
    AT (0, 0, 1e-10) RELATIVE Emon

/* The END token marks the instrument definition end */
END
```

Written by you!






McStas



2021
HighNESS
Virtual
McStas
School



Component file

```
*****
* Mcstas, neutron ray-tracing package
* Copyright 1997-2002, All rights reserved
* Risoe National Laboratory, Roskilde, Denmark
* Institut Laue Langevin, Grenoble, France
*
* Component: Source_flat
*
* %I
* Written by: Kim Lefmann
* Date: October 30, 1997
* Modified by: KL, October 4, 2001
* Modified by: Emmanuel Farhi, October 30, 2001. Serious bug corrected.
* Version: $Revision: 1.22 $
* Origin: Risoe
* Release: McStas 1.6
*
* A circular neutron source with flat energy spectrum and arbitrary flux
*
* %D
* The routine is a circular neutron source, which aims at a square target
* centered at the beam (in order to improve MU-acceptance rate). The angular
* divergence is then given by the dimensions of the target.
* The neutron energy is uniformly distributed between E0-dE and E0+dE.
*
* Example: Source_flat(radius=0.1, dist=2, xw=.1, yh=.1, E0=14, dE=2)
*
* %P
* radius: (m) Radius of circle in (x,y,0) plane where neutrons
* are generated.
* dist: (m) Distance to target along z axis.
* xw: (m) Width(x) of target
* yh: (m) Height(y) of target
* E0: (meV) Mean energy of neutrons.
* dE: (meV) Energy spread of neutrons.
* Lambda0 (AA) Mean wavelength of neutrons.
* dLambda (AA) Wavelength spread of neutrons.
* flux (1/(s*cm**2*s)) Energy integrated flux
*
* %E
*****
```

```
DEFINE COMPONENT Source_simple
DEFINITION PARAMETERS ()
SETTING PARAMETERS (radius, dist, xw, yh, E0=0, dE=0, Lambda0=0, dLambda=0, flux=1)
OUTPUT PARAMETERS ()
STATE PARAMETERS (x, y, z, vx, vy, vz, t, s1, s2, p)
DECLARE
|{
    double pmul, pdir;
}
INITIALIZE
|{
    pmul=flux*PI*1e4*radius*radius/mcget_ncount();
}
```

```
TRACE
|(
    double chi,E,Lambda,v,r, xf, yf, rf, dx, dy;
    t=0;
    z=0;

    chi=2*PI*rand01();
    r=sqrt(rand01())*radius;
    x=r*cos(chi);
    y=r*sin(chi);
)
randvec_target_rect(&xf, &yf, &rf, &pdir,
    0, 0, dist, xw, yh, ROT_A_CURRENT_COMP);

dx = xf-x;
dy = yf-y;
rf = sqrt(dx*dx+dy*dy+dist*dist);

p = pdir*pmul;

if(Lambda0==0) {
    E=E0+dr*randpm1(); /* Choose from uniform distribution */
    v=sqrt(E)*SE2V;
} else {
    Lambda=Lambda0+dLambda*randpm1();
    v = K2V*(2*PI/Lambda);
}

vz=v*dist/rf;
vy=v*dy/rf;
vx=v*dx/rf;
}

MCDISPLAY
|(
    magnify("xy");
    circle("xy",0,0,0, radius);
)
END
```

Written by developers
and possibly you!






McStas



2021
HighNESS
Virtual
McStas
School



Generated c-code

```

/* Automatically generated file. Do not edit.
 * Format: ANSI C source code
 * Creator: McStas <http://neutron.risoe.dk>
 * Instrument: My_Instrument.instr (My Instrument)
 * Date: Sat Apr 9 15:27:56 2005
 */

/* THOUSANDS of lines removed here.... */

/* TRACE Component Source. */
SIG_MESSAGE("Source (Trace)");
mcDEBUG_COMP("Source")
mccoordschange(mcpoSource, mcrotrSource,
    &mcnlx, &mcnly, &mcnlz,
    &mcnlvx, &mcnlvy, &mcnlvz,
    &mcnlt, &mcnlsx, &mcnlsy);
mcDEBUG_STATE(mcnlx, mcnly, mcnlz, mcnlvx, mcnlvy, mcnlvz, mcnlt, mcnlsx, mcnlsy, mcnlp)
#define x mcnlx
#define y mcnly
#define z mcnlz
#define vx mcnlvx
#define vy mcnlvy
#define vz mcnlvz
#define t mcnlt
#define s1 mcnlsx
#define s2 mcnlsy
#define p mcnlp
STORE_NEUTRON(2,mcnlx, mcnly, mcnlz, mcnlvx,mcnlvy,mcnlvz,mcnlt,mcnlsx,mcnley, mcnlsz, mcnlp);
mcScattered=0;
mcNCounter[2]++;
#define mcompcurname Source
#define mcompcurindex 2
{
    /* Declarations of SETTING parameters. */
MCNUM radius = mccSource_radius;
MCNUM dist = mccSource_dist;
MCNUM xv = mccSource_xv;
MCNUM yh = mccSource_yh;
MCNUM EO = mccSource_EO;
MCNUM dr = mccSource_dE;
MCNUM Lambda0 = mccSource_Lambda0;
MCNUM dLambda = mccSource_dLambda;
MCNUM flux = mccSource_flux;
#line 58 "Source_simple.comp"
{
    double chi,E,Lambda,v,r, xf, yf, rf, dx, dy;

t=0;
z=0;

chi=2*PI*rand01();                                /* Choose point on source */
r=sqrt(rand01())*radius;                          /* with uniform distribution. */
x=r*cos(chi);
y=r*sin(chi);

randvec_target_rect(&xf, &yf, &rf, &dir,
    0, 0, dist, xv, yh, ROT_A_CURRENT_COMP);
}

```

Written by mcstas!

McStas is a (pre)compiler!

Input is .comp and .instr files +
runtime functions for e.g. random
numbers

Output is a single c-file, which can
be compiled using e.g. gcc.



Can take input arguments if
needed.



Writing new comps or understanding existing is not that complex...

- Check our long list of components and look inside... Most of them are quite simple and short... Statistics:

Number of lines of code per component - 199 comps in total





McStas




2021

HighNESS
Virtual
McStas
School

Writing new comps or understanding existing is not that complex...

- Check our long list of components and look inside... Most of them are quite simple and short... Statistics:

Number of lines of code per component - 199 comps in total











- Well-developed community support
 - 30-40% of existing and new additions are from users
 - No direct refereeing of the code, but these requirements:
 - At least one test-instrument
 - Meaningful documentation headers (in-code docs)
 - Contributions go in dedicated contrib/ section of library




Example suite: ~140 instruments

Spin-echo B scan dependence of wavelength





2021 HighNESS Virtual McStas School



Your HighNESS...



**THIS IS NOT
THE END**

**IT'S JUST
THE BEGINNING**

