

Peter Willendrup

Further samples...



Slides adapted from Mads Bertelsen, ESS DMSC

Further samples in McStas

- Models for SANS and reflectometry
- Inelastic scattering, examples:
 - Phonon_simple
 - Isotropic_sqw
- McStas performance, TAS / Chopper

Small angle scattering SANS

- SANS method can be used for many types of material
- Often: Molecule + Liquid (buffer solution)
- Isotropic scattering



 **HighNess**

McStas




2021
HighNESS
Virtual
McStas
School


SANS

Small Angle Neutron Scattering

- Elastic Scattering
- Small angle -> small q -> big r
- Gain information on the molecular scale 10-100Å
- Low signal to noise
- Contrast method
- Instrument requirements: good collimation, long flight distance after detector.



$$q = \frac{4\pi}{\lambda} \sin(\theta)$$



HighNess

McStas



**2021
HighNESS
Virtual
McStas
School**

McStas has a suite of
SANS-models:

Try ellipsoidal and
cylindrical particles

-or-

Elliptic cylinders

Go for Nanodiscs and
Liposomes

*Also – SASmodels
from SASview*

SANS other samples

- SANS_AnySamp.comp
- SANS_DebyeS.comp
- SANS_Cylinders.comp
- SANSEllipticCylinders.comp
- SANS_Guinier.comp
- SANSLiposomes.comp
- SANSNanodiscs.comp
- SANSNanodiscsFast.comp
- SANSNanodiscsWithTags.
- SANSNanodiscsWithTagsFast
- SANSPDB.comp
- SANSPDBFAST.comp
- SANSShells.comp
- SANSSpheres.comp

HighNess

McStas



2021
HighNESS
Virtual
McStas
School

SANS spheres

Input parameters

Parameters in **boldface** are required; the others are optional.

Name	Unit	Description	Default
R	AA	Radius of scattering hard spheres	100
Phi	1	Particle volume fraction	1e-3
Delta_rho	fm/AA^3	Excess scattering length density	0.6
sigma_abs	m^-1	Absorption cross section density at 2200 m/s	0.05
xwidth	m	horiz. dimension of sample, as a width	0
yheight	m	vert. dimension of sample, as a height for cylinder/box	0
zdepth	m	depth of sample	0
radius	m	Outer radius of sample in (x,z) plane for cylinder/sphere	0
target_x			0
target_y	m	position of target to focus at	0
target_z			6
target_index	1	Relative index of component to focus at, e.g. next is +1	0
focus_xw	m	horiz. dimension of a rectangular area	0
focus_yh	m	vert. dimension of a rectangular area	0
focus_aw	deg	horiz. angular dimension of a rectangular area	0
focus_ah	deg	vert. angular dimension of a rectangular area	0
focus_r	m	Detector (disk-shaped) radius	0

Dilute, monodisperse, hard spheres in solution, with given contrast and radius

HighNess

McStas



2021
HighNESS
Virtual
McStas
School

SasView_models

Input parameters

Parameters in **boldface** are required; the others are optional.

Name	Unit	Description	Default
model_index			21
model_scale			1.0
model_pars			{60}
model_abs	1/m	Absorption cross section density at 2200 m/s	0.5
xwidth	m	horiz. dimension of sample, as a width	0
yheight	m	vert. dimension of sample, as a height for cylinder/box	0
zdepth	m	depth of sample	0
radius	m	Outer radius of sample in (x,z) plane for cylinder/sphere	0
target_x	m	relative focus target position	0
target_y	m	relative focus target position	0
target_z	m	relative focus target position	6
target_index	1	Relative index of component to focus at, e.g. next is +1	0
focus_xw	m	horiz. dimension of a rectangular area	0
focus_yh	m	vert. dimension of a rectangular area	0
focus_aw	deg	horiz. angular dimension of a rectangular area	0
focus_ah	deg	vert. angular dimension of a rectangular area	0
focus_r	m	Detector (disk-shaped) radius	0

HighNess

McStas



2021
HighNESS
Virtual
McStas
School

SasView_models

47	parallelepiped	(sld, solvent_sld, a_side, b_side, c_side)
48	parallelepiped_xy	(sld, solvent_sld, a_side, b_side, c_side, theta, phi, psi)
49	pearl_necklace	(radius, edge_separation, string_thickness, number_of_pearls, sld, string_sld, solvent_sld)
50	pearl_necklace_xy	(radius, edge_separation, string_thickness, number_of_pearls, sld, string_sld, solvent_sld)
51	sphere	(sld, solvent_sld, radius)
52	sphere_xy	(sld, solvent_sld, radius)
53	star_polymer	(radius2, arms)
54	star_polymer_xy	(radius2, arms)
55	stickyhardsphere	(effect_radius, volfraction, perturb, stickiness)
56	stickyhardsphere_xy	(effect_radius, volfraction, perturb, stickiness)
57	triaxial_ellipsoid	(sld, solvent_sld, req_minor, req_major, rpolar)
58	triaxial_ellipsoid_xy	(sld, solvent_sld, req_minor, req_major, rpolar, theta, phi, psi)

Inelastic scattering $S(\mathbf{q}, \omega)$

- partial differential cross section
- Scattering function
- Phonons, Spin waves, ...

$$\left(\frac{d^2\sigma}{d\Omega dE_f} \right)_{coh} = \frac{\sigma_{coh}}{4\pi} \frac{k_f}{k_i} N S(\mathbf{q}, \omega)$$

$$S(\mathbf{q}, \omega)_{coh} = \frac{1}{2\pi\hbar} \int \frac{1}{N} \sum_{jj'} \left\langle e^{-i\mathbf{q} \cdot \mathbf{R}_{j'}(0)} e^{-i\mathbf{q} \cdot \mathbf{R}_j(t)} \right\rangle e^{-i\omega t} dt$$

Popular component: Phonon_simple

- One isotropic acoustic phonon branch in all Brillouin zones on FCC Bravais single crystal

$$\frac{d^2\sigma'}{d\Omega dE_f} = b^2 \frac{k_f}{k_i} \frac{(2\pi)^3}{V_0} \frac{1}{2M} \exp(-2W) \times \sum_{\tau, q, p} \frac{(\boldsymbol{\kappa} \cdot \mathbf{e}_{q,p})^2}{\omega_{q,p}} \left\langle n_{q,p} + \frac{1}{2} \mp \frac{1}{2} \right\rangle \delta(\omega \pm \omega_{q,p}) \delta(\boldsymbol{\kappa} \pm \mathbf{q} - \boldsymbol{\tau})$$

Dispersion $d_1(\mathbf{q}) = c_1/a \sqrt{z - s_q}$

For FCC Bravis $z = 12 \quad s_q = \sum_{\mathbf{nn}} \cos(\mathbf{q} \cdot \mathbf{r}_{\mathbf{nn}})$

- M - Atomic mass
- b – scattering length
- n – bose factor
- a – fcc lattice spacing
- c - speed of sound
- $\boldsymbol{\kappa}$ – measured q vector
- \mathbf{q} – Phonon scattering vector

HighNess

McStas




2021
HighNESS
Virtual
McStas
School


Popu

- Dispers

ω




_simple




Popular component: Phonon_simple

- Example of the output



Popular component: Isotropic_sqw

- Isotropic processes (powder, liquid, ...)
- Use data files to describe $S(q,w)$ directly, coherent and incoherent
- Supports concentric



Popular component: Isotropic_sqw

- Isotropic processes (powder, liquid, ...)
- Use data files to describe $S(q, \omega)$ directly, coherent and incoherent
- Supports concentric

 HighNess

McStas






2021

HighNESS

Virtual


McStas

School




Popular component: Isotropic_sqw

- Rb liquid in time of flight
- Coherent and incoherent




Inelastic scattering in McStas

- Monte carlo sampling issues
- Need to sum over large amount of possible final states to find cross section
- Need large amount of rays to sample all the options




TAS

- Only a small fraction of neutrons arrive, r



Chopper spectrometers

- Only a small fraction of neutrons arrive



Conclusion

- Inelastic scattering supported in McStas, but could use more sample components
- Longer computational times required
- Advantages from simulation especially important for spectroscopy (resolution function)



McStas



**2021
HighNESS
Virtual
McStas
School**