

RESEARCH

A sample article title

Jane E. Doe^{1*} and John R.S. Smith^{1,2}

*Correspondence:

jane.e.doe@cambridge.co.uk

¹Department of Science,
University of Cambridge, London,
UKFull list of author information is
available at the end of the article**Abstract****First part title:** Text for this section.**Second part title:** Text for this section.**Keywords:** sample; article; author**Content**

Text and results for this section, as per the individual journal's instructions for authors.

Section title

Text for this section...

Sub-heading for section

Text for this sub-heading...

Sub-sub heading for section

Text for this sub-sub-heading...

Sub-sub-sub heading for section Text for this sub-sub-sub-heading...

In this section we examine the growth rate of the mean of Z_0 , Z_1 and Z_2 . In addition, we examine a common modeling assumption and note the importance of considering the tails of the extinction time T_x in studies of escape dynamics. We will first consider the expected resistant population at vT_x for some $v > 0$, (and temporarily assume $\alpha = 0$)

$$E[Z_1(vT_x)] = \int_0^{v \wedge 1} Z_0(uT_x) \exp(\lambda_1) du.$$

If we assume that sensitive cells follow a deterministic decay $Z_0(t) = xe^{\lambda_0 t}$ and approximate their extinction time as $T_x \approx -\frac{1}{\lambda_0} \log x$, then we can heuristically estimate the expected value as

$$\begin{aligned} E[Z_1(vT_x)] \\ = \frac{\mu}{r} \log x \int_0^{v \wedge 1} x^{1-u} x^{(\lambda_1/r)(v-u)} du. \end{aligned} \quad (1)$$

Thus we observe that this expected value is finite for all $v > 0$ (also see [1, 2, 3, 4, 5, 6]).

Appendix

Text for this section. . .

Acknowledgements

Text for this section. . .

Funding

Text for this section. . .

Abbreviations

Text for this section. . .

Availability of data and materials

Text for this section. . .

Ethics approval and consent to participate

Text for this section. . .

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Text for this section. . .

Authors' contributions

Text for this section . . .

Authors' information

Text for this section. . .

Author details

¹Department of Science, University of Cambridge, London, UK. ²Institute of Biology, National University of Sciences, Kiel, Germany.

References

1. Koonin, E.V., Altschul, S.F., Bork, P.: Brca1 protein products: functional motifs. Nat. Genet. **13**, 266–267 (1996)

2. Jones, X.: Zeolites and synthetic mechanisms. In: Smith, Y. (ed.) Proceedings of the First National Conference on Porous Sieves: 27–30 June 1996; Baltimore, pp. 16–27 (1996)


3. Margulis, L.: Origin of Eukaryotic Cells. Yale University Press, New Haven (1970)

4. Schnepf, E.: From prey via endosymbiont to plastids: comparative studies in dinoflagellates. In: Lewin, R.A. (ed.) Origins of Plastids, 2nd edn., pp. 53–76. Chapman and Hall, New York (1993)

5. Kohavi, R.: Wrappers for performance enhancement and obvious decision graphs. PhD thesis, Stanford University, Computer Science Department (1995)

6. ISSN International Centre: The ISSN register (2006). <http://www.issn.org> Accessed Accessed 20 Feb 2007

Figures



Tables

Table 1 Sample table title. This is where the description of the table should go

	B1	B2	B3
A1	0.1	0.2	0.3
A2
A3

Additional Files

Additional file 1 — Sample additional file title

Additional file descriptions text (including details of how to view the file, if it is in a non-standard format or the file extension). This might refer to a multi-page table or a figure.

Additional file 2 — Sample additional file title
Additional file descriptions text.