

## 4.7: Covalent Bonding: Simple Lewis Structures

In the Lewis model, we represent covalent bonding with a *Lewis structure*, which depicts neighboring atoms as sharing some (or all) of their valence electrons in order to attain octets (or duets for hydrogen).

### Single Covalent Bonds

Consider hydrogen and oxygen, which have the following Lewis symbols:



In water, these atoms share their unpaired valence electrons so that each hydrogen atom gets a duet and the oxygen atom gets an octet as represented with this Lewis structure:




The shared electrons—those that appear in the space between the two atoms—count toward the octets (or duets) of *both of the atoms*.



A shared pair of electrons is called a **bonding pair**, while a pair that is associated with only one atom—and therefore not involved in bonding—is called a **lone pair**. Lone pair electrons are also called **nonbonding electrons**.




We often represent a bonding pair of electrons by a dash to emphasize that the pair constitutes a chemical bond.



The Lewis model helps explain the tendency of some elements to form diatomic molecules. For example, consider the Lewis symbol for chlorine:




If two Cl atoms pair together, they each get an octet.






Atoms can also share three electron pairs. Consider the Lewis structure of another diatomic molecule,  $\text{N}_2$ . Because each N atom has five valence electrons, the Lewis structure for  $\text{N}_2$  has 10 electrons. Both nitrogen atoms attain octets by sharing three electron pairs:




In this case, the bond is a **triple bond**. Triple bonds are even shorter and stronger than double bonds. When we examine nitrogen in nature, we find that it exists as a diatomic molecule with a very strong bond between the two nitrogen atoms. The bond is so strong that it is difficult to break, making  $\text{N}_2$  a relatively unreactive molecule.

## Covalent Bonding: Models and Reality


The Lewis model predicts the properties of molecular compounds in many ways. We have already seen how it explains the existence of several diatomic elements. The Lewis model also accounts for why particular combinations of atoms form molecules and others do not. For example, why is water  $\text{H}_2\text{O}$  and not  $\text{H}_3\text{O}$ ? We can write a good Lewis structure for  $\text{H}_2\text{O}$  but not for  $\text{H}_3\text{O}$ .



In this way, the Lewis model predicts that  $\text{H}_2\text{O}$  should be stable, while  $\text{H}_3\text{O}$  should not be, and that is in fact the case. However, if we remove an electron from  $\text{H}_3\text{O}$ , we get  $\text{H}_3\text{O}^+$ , which should be stable (according to the Lewis model) because, when we remove the extra electron, oxygen gets an octet.



This ion, called the hydronium ion, is in fact stable in aqueous solutions (see Section 8.7). The Lewis model predicts other possible combinations for hydrogen and oxygen as well. For example, we can write a Lewis structure for  $\text{H}_2\text{O}_2$  as follows:



Indeed,  $\text{H}_2\text{O}_2$ , or hydrogen peroxide, exists and is often used as a disinfectant and a bleach.


The Lewis model also accounts for why covalent bonds are highly *directional*. The attraction between two covalently bonded atoms is due to the sharing of one or more electron pairs in the space between them. Thus, each bond links just one specific pair of atoms—in contrast to ionic bonds, which are *nondirectional* and hold together an entire array of ions. As a result, the fundamental units of covalently bonded compounds are individual molecules. These molecules can interact with one another in a number of different ways that we will cover in Chapter 11. However, in covalently bonded molecular compounds, the interactions *between* molecules (intermolecular forces) are generally much weaker than the bonding interactions within a molecule (intramolecular forces), as shown in Figure 4.10. When a molecular compound melts or boils, the molecules themselves remain intact—only the relatively weak interactions between molecules must be overcome. Consequently, molecular compounds tend to have lower melting and boiling points than ionic compounds.

**Figure 4.10 Intermolecular and Intramolecular Forces**

The covalent bonds between atoms of a molecule are much stronger than the interactions between

molecules. To bring a molecular substance to a boil, only the relatively weak intermolecular forces have to be overcome, so molecular compounds often have low boiling points.

### Molecular Compound



#### Conceptual Connection 4.5 Energy and the Octet Rule

#### Conceptual Connection 4.6 Ionic and Molecular Compounds

Interactive Interac

Not for Distribution

*Not for Distribution*