

Miyaoka-Yau inequality for hyperplane arrangements

Martin de Borbon

arXiv: 2411.09573 (joint with Dmitri Panov)

Loughborough University

29/11/2024

- Main result: $Q \leq 0$ on C
- Proof: Bogomolov-Gieseker inequality for stable parabolic bundles
- Application: reflection arrangements

Basic definitions

- This is a paper about complex projective space

$$\mathbb{CP}^n = (\mathbb{C}^{n+1} \setminus \{0\}) / \mathbb{C}^*.$$

- Let \mathcal{H} be a hyperplane arrangement in \mathbb{CP}^n . That is, \mathcal{H} is a finite set of pairwise distinct complex hyperplanes

$$H \subset \mathbb{CP}^n.$$

- Let $L \subset \mathbb{CP}^n$ be a linear subspace obtained as intersection of hyperplanes $H \in \mathcal{H}$. The **multiplicity** of L is

$$m_L = |\{H \in \mathcal{H} \mid H \supset L\}|.$$

Note: $m_L \geq \text{codim } L$

Codimension 2 subspaces

Let $L \subset \mathbb{CP}^n$ be a codimension 2 intersection of hyperplanes in \mathcal{H} .
We say that

- L is **reducible** if $m_L = 2$
- L is **irreducible** if $m_L \geq 3$




Figure 1: Reducible (left) and irreducible (right).

The Hirzebruch quadratic form

- $\mathcal{H} = \{H_1, \dots, H_N\}$ hyperplane arrangement in \mathbb{CP}^n
- σ_i = number of irreducible codimension 2 subspaces $L \subset H_i$
- The **Hirzebruch quadratic form** of \mathcal{H} is the homogeneous degree 2 polynomial on \mathbb{R}^N given by

$$Q(a_1, \dots, a_N) = \sum_{i,j=1}^N Q_{ij} a_i a_j$$

$$Q_{ij} = \begin{cases} -(n+1)\sigma_i + 2n & \text{if } i = j \\ -2 & \text{if } i \neq j \text{ and } L = H_i \cap H_j \text{ is reducible} \\ n-1 & \text{if } i \neq j \text{ and } L = H_i \cap H_j \text{ is irreducible} \end{cases}$$

Let σ_i be the number of points p with $r_p \geq 3$ lying on the i -th line of the given arrangement of k lines in the plane. We consider the $(k \times k)$ -symmetric matrix A with

$$(3) \quad A_{ij} = \begin{cases} 3\sigma_i - 4 & (i = j), \\ 2 & (i \neq j, \quad p \in L_i \cap L_j \text{ with } r_p = 2), \\ -1 & (i \neq j, \quad p \in L_i \cap L_j \text{ with } r_p \geq 3). \end{cases}$$

With the k lines we associate real variables x_i and let x be the column vector (x_1, \dots, x_k) . With the s points p_j with $r_{p_j} \geq 3$ we associate real variables y_j . For each point p_j with $r_{p_j} \geq 3$ we consider the linear form

$$P_j(x, y) = 2y_j + \sum_{p_j \in L_i} x_i, \quad \text{where } y = (y_1, \dots, y_s).$$

Höfer's formula. For the algebraic surface Y (a good covering of S of degree d with respect to $L_1, \dots, L_k, E_1, \dots, E_s$ and the given branching numbers $n_1, \dots, n_k, m_1, \dots, m_s$) we have

$$(4) \quad (3c_2(Y) - c_1^2(Y))/d = \frac{1}{4} \left(x^t A x + \sum_{j=1}^s P_j(x, y)^2 \right),$$

where $x_i = 1 - \frac{1}{n_i}$ and $y_j = -1 - \frac{1}{m_j}$.

The matroid polytope

\mathcal{H} is an *essential* and *irreducible* hyperplane arrangement in \mathbb{CP}^n

- A **basis** of $\mathcal{H} = \{H_1, \dots, H_N\}$ is a subset $\mathcal{B} \subset \mathcal{H}$ consisting of $n + 1$ linearly independent hyperplanes
- The indicator vector of \mathcal{B} is the 1/0 vector

$$\mathbf{e}_{\mathcal{B}} = \sum_{i | H_i \in \mathcal{B}} \mathbf{e}_i$$

where $\mathbf{e}_1, \dots, \mathbf{e}_N$ are the standard basis vectors of \mathbb{R}^N

- The **matroid polytope** is the convex hull of the vectors $\mathbf{e}_{\mathcal{B}}$

$$P = \text{conv}\{ \mathbf{e}_{\mathcal{B}} \mid \mathcal{B} \text{ is a basis of } \mathcal{H} \}.$$

Note: P is contained in the $(N - 1)$ -simplex $\Delta \subset \mathbb{R}^N$ with

$$\Delta = \left\{ (a_1, \dots, a_N) \in \mathbb{R}^N \mid a_i \geq 0, \sum_i a_i = n + 1 \right\}$$

The semistable and stable cones

- The **semistable cone** is the cone over the matroid polytope

$$C = \mathbb{R}_{\geq 0} \cdot P = \text{cone} \{ \mathbf{e}_{\mathcal{B}} \mid \mathcal{B} \text{ is a basis of } \mathcal{H} \}.$$

It is a convex polyhedral cone contained in the octant $\mathbb{R}_{\geq 0}^N$

- The **stable cone** is the interior of $C \subset \mathbb{R}^N$

$$C^\circ = \text{int}(C).$$

- \mathcal{H} is essential and irreducible $\iff \dim P = N - 1$
 $\iff C^\circ$ is non empty

Defining linear inequalities of the stable cone

- Let \mathcal{L} be the finite set of *non-empty* and *proper* linear subspaces $L \subset \mathbb{CP}^n$ obtained by intersecting members of \mathcal{H}
- $(a_1, \dots, a_N) \in C^\circ \iff \forall i : a_i > 0 \text{ and } \forall L \in \mathcal{L} :$

$$\sum_{i \mid L \subset H_i} a_i < \frac{\text{codim } L}{n+1} \cdot \sum_{i=1}^N a_i$$

Relation to Geometric Invariant Theory:

- Standard embedding $\mathbb{CP}^n \subset \mathfrak{su}(n+1)^*$ as a coadjoint orbit.
Let $p_i \in (\mathbb{CP}^n)^*$ be the annihilator of H_i
- $(a_1, \dots, a_N) \in C^\circ \iff \exists F \in SL(n+1, \mathbb{C})$ such that the centre of mass of the points $F(p_i)$ with weights a_i is $0 \in \mathfrak{su}(n+1)^*$
- Familiar case $n = 1$, then $\mathbb{CP}^1 = S^2$ and $\mathfrak{su}(2)^* = \mathbb{R}^3$

Main Result

Theorem (Miyaoka-Yau inequality, dB-Panov 2024)

The Hirzebruch quadratic form is non-positive on the semistable cone:

$$C \subset \{Q \leq 0\}$$

- If $n = 1$ then $Q \equiv 0$
- If $n = 2$ this follows from Panov's *Polyhedral Kähler Manifolds*, Geometry & Topology, 2009
- **Conjecture:** if $\mathbf{a} = (a_1, \dots, a_N) \in C^\circ$ is such that $Q(\mathbf{a}) = 0$ and $a_i \in (0, 1)$. Then there is a Kähler metric on \mathbb{CP}^n of constant holomorphic sectional curvature with cone angles $2\pi\alpha_i$ in transverse directions to the hyperplanes $H_i \in \mathcal{H}$, with $\alpha_i = 1 - a_i$.

klt and CY arrangements

A weighted arrangement is a pair $(\mathcal{H}, \mathbf{a})$ consisting of:

- a hyperplane arrangement \mathcal{H} in \mathbb{CP}^n ;
- a weight vector $\mathbf{a} \in \mathbb{R}^{\mathcal{H}}$ with components $a_H > 0$.

The weighted arrangement $(\mathcal{H}, \mathbf{a})$ is

- klt if

$$\forall L \in \mathcal{L} : \sum_{H \mid H \supset L} a_H < \operatorname{codim} L$$

where \mathcal{L} is the set of non-empty and proper subspaces $L \subset \mathbb{CP}^n$ obtained by intersecting hyperplanes in \mathcal{H} . In particular,

$$0 < a_H < 1$$

- Calabi-Yau (CY) if

$$\sum_{H \in \mathcal{H}} a_H = n + 1$$

Restatement of the main theorem

Theorem (Miyaoka-Yau inequality, dB-Panov 2024)

Suppose that the weighted arrangement $(\mathcal{H}, \mathbf{a})$ is klt and CY. Then

$$Q(\mathbf{a}) = \sum_{L \in \mathcal{L}_{\text{irr}}^{n-2}} a_L^2 - \frac{1}{2} \sum_{H \in \mathcal{H}} B_H \cdot a_H^2 - \frac{n+1}{2} \leq 0.$$

- $\mathcal{L}_{\text{irr}}^{n-2}$ is the set of irreducible codimension 2 subspaces $L \subset \mathbb{CP}^n$
- The weight a_L at $L \in \mathcal{L}_{\text{irr}}^{n-2}$ is given by

$$a_L = \frac{1}{2} \cdot \sum_{H \supset L} a_H$$

- $B_H + 1$ is the number of $L \in \mathcal{L}_{\text{irr}}^{n-2}$ with $L \subset H$

Sketch proof

- Logarithmic resolution

$$X \xrightarrow{\pi} \mathbb{CP}^n$$

with $D = \pi^{-1}(\mathcal{H})$ a simple normal crossing divisor.

- X is the *minimal De Concini-Procesi wonderful model* of \mathcal{H} .
- The irreducible components of D are in bijective correspondence with non-empty and proper irreducible subspaces $L \in \mathcal{L}_{\text{irr}}$

$$D = \bigcup_{L \in \mathcal{L}_{\text{irr}}} D_L$$

where D_L is the unique irreducible component of D such that

$$\pi(D_L) = L.$$