

A Miyaoka-Yau inequality for hyperplane arrangements in $\mathbb{CP}^d$

Martin de Borbon (joint with Dmitri Panov)

Living on the edge of the moduli space, Aarhus


18-22 May 2026

- Introduction: flat metrics on surfaces with conical singularities
- Main result: polyhedral Kähler metrics on $\mathbb{CP}^d$
- Sketch proof: Kobayashi-Hitchin correspondence for parabolic bundles
- Conjectures: Kähler-Einstein metrics and Miyaoka-Yau inequality for log pairs

The 2-cone


Let $\alpha > 0$. Denote by $C(2\pi\alpha)$ the cone over a circle of length $2\pi\alpha$.
In polar coordinates,

$$dr^2 + \alpha^2 r^2 d\theta^2$$



Flat metrics on surfaces with conical singularities

- Let Σ be a compact orientable 2-dimensional surface.
- A flat metric on Σ with conical singularities is a flat metric on $\Sigma \setminus \{p_1, \dots, p_n\}$ that is locally isometric to $C(2\pi\alpha_i)$ near p_i
- Examples:



Surface of a cube




Double of a triangle

Gauss-Bonnet: $\sum_{i=1}^n (1 - \alpha_i) = \chi(\Sigma)$

- In 2D: metric + orientation $\implies$ complex structure
- The complex structure extends smoothly across conical singularities: if $z = r^{1/\alpha}e^{i\theta}$ then

$$dr^2 + \alpha^2 r^2 d\theta^2 = \alpha^2 |z|^{2\alpha-2} |dz|^2$$

Flat metrics on the 2-sphere with cone points



Surface of a cube. Flat metric on S^2 with 8 cone points of total angle $2\pi\beta = 3(\pi/2)$.



Double of a triangle with interior angles $\pi\beta_i$. Flat metric on S^2 with 3 cone points of total angle $2\pi\beta_i$

$$\sum_{i=1}^n (1 - \beta_i) = 2$$

(Gauss-Bonnet)

- Surface of a cube: $8 \cdot (1 - 3/4) = 2$
- Double of a triangle: $3 - (\beta_1 + \beta_2 + \beta_3) = 2$

McOwen (1988), Troyanov (1991)

Let X be a compact Riemann surface. Let $x_1, \dots, x_n \in X$ and let $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbb{R}_{>0})^n$. Define

$$\chi(X, \alpha) = \chi(X) + \sum_{i=1}^n (\alpha_i - 1)$$

- If $\chi(X, \alpha) < 0$ then there exists a unique Kähler metric on X with $K_g = -1$ and cone angle $2\pi\alpha_i$ at x_i
- If $\chi(X, \alpha) = 0$ then there exists a unique up to scale Kähler metric on X with $K_g = 0$ and cone angle $2\pi\alpha_i$ at x_i
- If $\chi(X, \alpha) > 0$ and $\alpha_i \in (0, 1)$ then there exists a unique Kähler metric on X with $K_g = 1$ and cone angle $2\pi\alpha_i$ at x_i if and only if

$$\forall i \quad a_i < \frac{1}{2} \cdot \sum_{j=1}^n a_j \quad \text{where } a_i = 1 - \alpha_i$$

Main result

Theorem (dB-Panov, 2025-26)

Let $\mathcal{H} = \{H_1, \dots, H_n\}$ be a hyperplane arrangement in $\mathbb{CP}^d$ with $d \geq 1$. Let $\alpha_1, \dots, \alpha_n \in (0, 1)$. Write $a_i = 1 - \alpha_i$. Assume that:

- ① $\sum_{i=1}^n a_i = d + 1$
- ② the weighted arrangement $\{(H_i, a_i)\}_{i=1}^n$ is **stable**

Then the following inequality holds:

$$Q(a_1, \dots, a_n) \leq 0 \tag{1}$$

where Q is the **Hirzebruch quadratic form** of $\mathcal{H}$. Moreover, the equality holds if and only if there exists a **polyhedral Kähler metric** on $\mathbb{CP}^d$ with cone angles $2\pi\alpha_i$ along H_i .

Let σ_i be the number of points p with $r_p \geq 3$ lying on the i -th line of the given arrangement of k lines in the plane. We consider the $(k \times k)$ -symmetric matrix A with

$$(3) \quad A_{ij} = \begin{cases} 3\sigma_i - 4 & (i = j), \\ 2 & (i \neq j, \quad p \in L_i \cap L_j \text{ with } r_p = 2), \\ -1 & (i \neq j, \quad p \in L_i \cap L_j \text{ with } r_p \geq 3). \end{cases}$$

With the k lines we associate real variables x_i and let x be the column vector $(x_1, \dots, x_k)$. With the s points p_j with $r_{p_j} \geq 3$ we associate real variables y_j . For each point p_j with $r_{p_j} \geq 3$ we consider the linear form

$$P_j(x, y) = 2y_j + \sum_{p_j \in L_i} x_i, \quad \text{where } y = (y_1, \dots, y_s).$$

Höfer's formula. For the algebraic surface Y (a good covering of S of degree d with respect to $L_1, \dots, L_k, E_1, \dots, E_s$ and the given branching numbers $n_1, \dots, n_k, m_1, \dots, m_s$) we have

$$(4) \quad (3c_2(Y) - c_1^2(Y))/d = \frac{1}{4} \left(x^t A x + \sum_{j=1}^s P_j(x, y)^2 \right),$$

where $x_i = 1 - \frac{1}{n_i}$ and $y_j = -1 - \frac{1}{m_j}$.

Hyperplane arrangements: basics and notation

- A **hyperplane arrangement** is a finite set $\mathcal{H} = \{H_1, \dots, H_n\}$ of pairwise distinct complex hyperplanes $H_i \subset \mathbb{CP}^d$
- Index set $[n] = \{1, \dots, n\}$
- The **intersection poset** of $\mathcal{H}$ is the set of all proper linear subspaces obtained by intersecting hyperplanes in $\mathcal{H}$

$$\mathcal{L} = \left\{ L \subset \mathbb{CP}^d \mid L = \bigcap_{i \in I} H_i \text{ for some non-empty } I \subset [n] \right\}$$

- The **rank** of $L \in \mathcal{L}$ is defined as

$$r(L) = \text{codim}(L \subset \mathbb{CP}^d)$$

Take the convention that $r(\emptyset) = d + 1$

- $\mathcal{L}^k = \{L \in \mathcal{L} \mid r(L) = k\}$
- The **multiplicity** of an element $L \in \mathcal{L}$ is defined as

$$\text{mult}(L) = |\{i \mid H_i \supset L\}|$$

Stability

A weighted arrangement $\{(H_i, a_i)\}_{i=1}^n$ consisting of hyperplanes $H_i \subset \mathbb{CP}^d$ together with weights $a_i > 0$ is **stable** if

$$\forall L \in \mathcal{L} \setminus \{\emptyset\} : \quad \frac{1}{r(L)} \cdot \sum_{i|H_i \supset L} a_i < \frac{1}{d+1} \cdot \sum_{i=1}^n a_i$$

Suppose that

$$\sum_{i=1}^n a_i = d+1 \tag{2}$$

Then stability is equivalent to

$$\forall L \in \mathcal{L} \setminus \{\emptyset\} : \quad \sum_{i|H_i \supset L} a_i < r(L) \tag{3}$$

Let (X, Δ) be the log pair $(\mathbb{CP}^d, \sum a_i H_i)$.

Equation (2) $\iff (X, \Delta)$ is Calabi-Yau

Equation (3) $\iff (X, \Delta)$ is klt

The Hirzebruch quadratic form

Let $\mathcal{H} = \{H_1, \dots, H_n\}$ be a hyperplane arrangement in $\mathbb{CP}^d$. The **Hirzebruch quadratic form** of $\mathcal{H}$ is the homogeneous degree 2 polynomial on $\mathbb{R}^n$ given by

$$Q(x_1, \dots, x_n) = \sum_{i,j=1}^k Q_{ij} x_i x_j$$

$$Q_{ij} = \begin{cases} -(d+1)\sigma_i + 2d & \text{if } i = j \\ -2 & \text{if } i \neq j \text{ and } \text{mult}(H_i \cap H_j) = 2 \\ d-1 & \text{if } i \neq j \text{ and } \text{mult}(H_i \cap H_j) \geq 3 \end{cases}$$




where $\sigma_i = |\{L \in \mathcal{L}^2 \mid L \subset H_i, \text{mult}(L) \geq 3\}|$

Constant curvature surfaces with cone singularities

$$\begin{cases} dr^2 + \beta^2 \sin^2 r d\theta^2 \\ dr^2 + \beta^2 r^2 d\theta^2 \\ dr^2 + \beta^2 \sinh^2 r d\theta^2 \end{cases}$$



Double of spherical triangle.
Constant curvature 1 metric on the
2-sphere with 3 cone points.



- Orbifolds: $\beta = 1/k$ for $k \geq 2$ integer
- Modular curve $\mathbb{H}/PSL(2, \mathbb{Z})$.
Hyperbolic metric with two
cone points $1/2, 1/3$

Existence and uniqueness

- Fix $\vec{\beta} = (\beta_1, \dots, \beta_n)$ with $\sum_{i=1}^n (1 - \beta_i) = 2$
- $\text{Met}(\vec{\beta}) =$ flat metrics on S^2 with n cone points x_i of total angle $2\pi\beta_i$ modulo marked (orientation preserving) isometry and scale
- $\mathcal{M}_{0,n} =$ configuration of n distinct marked points in $\mathbb{CP}^1$ up to the action of Möbius transformations $PSL(2, \mathbb{C})$

Theorem (Troyanov)


The forgetful map $F : \text{Met}(\vec{\beta}) \rightarrow \mathcal{M}_{0,n}$ is a bijection

Proof: If $x_1, \dots, x_n \in \mathbb{C}$ then

$$\left(\prod_{i=1}^n |z - x_i|^{2\beta_i - 2} \right) |dz|^2$$

extends smoothly over ∞ and defines a flat (Kähler) metric on $\mathbb{CP}^1$ with cone angles $2\pi\beta_i$ at the points x_i

Collision of two cone points



$$\gamma + (1 - \beta_1) + (1 - \beta_2) = 1$$


Model flat metrics on $\mathbb{C}$ with cone points

- $$\sum_{i=1}^p (1 - \beta_i) < 1$$

$$g_F = \left(\prod_{i=1}^p |z - x_i|^{2\beta_i - 2} \right) |dz|^2$$

- flat Kähler metric on $\mathbb{C}$ with cone angles $2\pi\beta_i$ at x_i
- isometric to the 2-cone $C(2\pi\gamma)$ outside a compact set

$$1 - \gamma = \sum_{i=1}^p (1 - \beta_i)$$



$$\lim_{\lambda \rightarrow 0} \lambda \cdot g_F = C(2\pi\gamma)$$

Bubble trees

Consider a family of flat metrics g_t for which a cluster of cone points $x_1(t), \dots, x_p(t)$ collides to 0 as $t \rightarrow 0$


- For $f \in \mathcal{O}_{\mathbb{C},0}$ let $\nu(f)$ = order of vanishing of f at 0
- For $r \geq 0$ we have an equivalence relation $f \sim_r g$ if $\nu(f - g) \geq r$
- If $S = \{x_1(t), \dots, x_p(t)\} \subset \mathcal{O}_{\mathbb{C},0}$ is a finite set then the equivalence classes of $\sim_r$ make a tree $\mathcal{T}$ with root S and leaves $\{x_i(t)\}$
- To every non-leaf vertex $\mathbf{v} \in \mathcal{T}$ we associate a model flat metric $B_{\mathbf{v}}$ on $\mathbb{C}$. The number of cone points of $B_{\mathbf{v}}$ is equal to the number of children of $\mathbf{v}$.
- The cone angles are $2\pi\gamma_{\mathbf{w}}$ with $1 - \gamma_{\mathbf{w}} = \sum_{i|x_i \in \mathbf{w}} (1 - \beta_i)$ and the position of the cone points are $x_{\mathbf{w}} = \lim_{t \rightarrow 0} t^{-k} x_i(t)$ where k is the smallest integer such that the elements of $\mathbf{v}$ are *not* $\sim_k$ equivalent.

Rescaled limits

- Let $s \in \mathcal{O}_{\mathbb{C},0}$ be a section
- For $\alpha > 0$ let h_α be the pointed Gromov-Hausdorff limit

$$h_\alpha = \lim_{t \rightarrow 0} (|t|^{-2\alpha} \cdot g_t, s(t))$$

- The section determines a path in the tree with vertices $\mathbf{v}_1, \dots, \mathbf{v}_\ell$
- $0 = \alpha_0 < \alpha_1 < \dots < \alpha_\ell$ such that $h_\alpha = C(2\pi\gamma_{\mathbf{v}_i})$ (with base point its vertex) if $\alpha_{i-1} < \alpha < \alpha_i$ and $h_\alpha = B_{\mathbf{v}_i}$ if $\alpha = \alpha_i$ (with base point $\lim_{t \rightarrow 0} t^{-k} s(t)$).



$s(t) = t - t^4 + (\text{h.o.t.})$ shown in red where $\mathcal{T}$



- Thurston: complex hyperbolic cone metric on $\overline{\text{Met}}(\vec{\beta})$, divisors represent collision of pair of points
- Deligne-Mumford-Knudsen compactification $\overline{\mathcal{M}}_{0,n}$, divisors correspond to partitions of $\{1, \dots, n\}$ into two disjoint sets, its points represent nodal curves with 2 irreducible components
- Logarithmic resolution $\overline{\mathcal{M}}_{0,n} \rightarrow \overline{\text{Met}}(\vec{\beta})$ (Koziarz-Nguyen), points in $\overline{\mathcal{M}}_{0,n}$ represent bubble trees

Kähler-Einstein metrics with cone singularities

$D \subset X$ smooth complex
hypersurface and $0 < \beta < 1$

$\longleftrightarrow$

$(X, (1 - \beta)D)$





- $\text{Ric}(g_{KE}) = \lambda \cdot g_{KE}$ on $X \setminus D$
- $g_{KE} \sim C(2\pi\beta) \times \mathbb{C}^{n-1}$ near D

- Donaldson's interior Schauder estimate for the Laplace operator
- Existence and regularity theory, polyhomogeneous expansion (Jeffress-Mazzeo-Rubinstein)
- Chern-Weil formulas (Jian Song and Xiaowei Wang)
- Algebraic structure on non-collapsed Gromov-Hausdorff limits (Chen-Donaldson-Sun)

Polyhedral Kähler cones



$\bar{g}$ = constant curvature 1
metric on the 3-sphere



$\xrightarrow{\text{Hopf}}$


Cone:

$$d\rho^2 + \rho^2 \bar{g}$$




Proof of Theorem

$$\text{continuity path: } (\omega_0 + i\partial\bar{\partial}\varphi_t)^2 = e^{tf}\omega_0^2$$



$d = 2 \rightarrow S^1$ -symmetry

Gibbons-Hawking ansatz




$$g_{RF} = V g_{\mathbb{R}^3_\beta} + (1/V) \alpha^2, \quad d\alpha = -\star_\beta dV$$


L^2 -norm of the curvature

Chern-Weil: the energy of a KE metric depends only on topology.

$$E(g) := \frac{1}{8\pi^2} \int |\text{Riem}(g)|^2$$



Blow-up limits



$C_\epsilon = \{P_d(z, w) + (\text{h. o. t.}) = \epsilon\}$ as $\epsilon \rightarrow 0$. If we rescale coordinates by $z = \epsilon^{1/d}\tilde{z}$, $w = \epsilon^{1/d}\tilde{w}$ then C_ϵ converges to $C = \{P_d(\tilde{z}, \tilde{w}) = 1\}$ as $\epsilon \rightarrow 0$.

Conjecture:

- For every $1/2 < \beta < 1$ there is a Calabi-Yau metric on $\mathbb{C}^2$ with cone angle $2\pi\beta$ along the parabola $\{w = z^2\}$
- The tangent cone at infinity equal to $\mathbb{C}_\gamma \times \mathbb{C}$, where $\gamma = 2\beta - 1$
- The energy of the metric is finite and given by $E = 1 - \beta$

Moduli spaces of pairs

- Ascher-DeVleming-Liu: K -moduli space of pairs $(\mathbb{CP}^2, (1 - \beta)C)$ with $\deg C = 4$ is the GIT quotient for $5/8 < \beta < 1$
- If $\gamma = 2\beta - 1$ then $1/4 < \gamma < 1$. By Li-Sun there is a KE metric with cone angle $2\pi\gamma$ along $C_0 = \{Q = 0\}$ where $Q = X_0^2 + X_1^2 + X_2^2$.
- Let P be a generic homogeneous degree 4 and $C_\epsilon = \{\epsilon P + Q^2 = 0\}$. There are eight branch points $\{P = 0\} \cap \{Q = 0\}$
- Family of KE metrics g_ϵ with cone angle $2\pi\beta$ along C_ϵ and a limiting KE metric g_0 with cone angle $2\pi\gamma$ along C_0
- The convergence $g_\epsilon \rightarrow g_0$ is modelled in directions transverse to C_0 by two cone points of angle $2\pi\beta$ which collide into a single cone point of angle $2\pi\gamma$
- It is expected that the parabola CY metric is a re-scaled limit of the sequence g_ϵ at each of the eight branch points
- The Song-Wang energy formula $E = \chi(X) + (\beta - 1)\chi(D)$ gives $E(g_\epsilon) - E(g_0) = 8(1 - \beta)$

A general theory

Non-collapsed polarized KE limit space (X, d_{KE})

$$C_p X := \lim_{\lambda \rightarrow 0} (X, \lambda^{-1} d_{KE}, p)$$

- Uniqueness, $C_p X$ is an affine algebraic variety (Donaldson-Sun).
Order of vanishing w.r.t. $d_{KE} \implies$ two steps degeneration.
- $C_p X$ depends only on $\mathcal{O}_{X,p}$ (the germ of X at p).
Chi Li's normalized volumes of valuations.

Sasaki geometry precedents: Martelli-Sparks-Yau (volume minimization), Collins-Székelyhidi (K-stability) .

Question

Describe all possible rescaled limits of a family.

Song Sun's *minimal bubbles*.

THANK YOU!