

A.P. SHAH INSTITUTE OF TECHNOLOGY

Department of Computer Science and Engineering
Data Science

Homeomorphic Graph


A graph homomorphism F from a graph G = (V, E) to a graph G' = (V', E') is written as: $f: G \rightarrow G'$. It is a mapping $f: V \rightarrow V'$ from the vertex set of G to the vertex set of G' such that $\{u, v\} \in E \Rightarrow \{f(u), f(v) \in E'\}$.

In simple words, If another graph G* can be formed by dividing the edge of G with additional vertices, or if a Graph G* can be obtained by introducing vertices of degree 2 in any edge of a Graph G, then the graph G* is complete. Both the graphs G and G* are known as Homeomorphic graphs.

Properties of Homeomorphic Graph

- Homomorphism always retains a graph's edges and connectedness.
- If a homomorphism is a bijective mapping, then it is an isomorphism.
- The compositions of homomorphisms are also homomorphisms.
- Finding out if there is any homomorphic graph of another graph is an NP-complete problem.


Example




PARSHVANATH CHARITABLE TRUST'S

A.P. SHAH INSTITUTE OF TECHNOLOGY

Department of Computer Science and Engineering
Data Science





As we can see in both examples, G' can be derived from G,' and H' can be derived from H by introducing a vertex of degree 2.