

PARSHWANATH CHARITABLE TRUSTS



A.P. SHAH INSTITUTE OF TECHNOLOGY

Department of Computer Science and Engineering Data Science




Semester:

Subject Blatistics for AIBDS

Academic Year: 2023 2024


NORMAL DISTRIBUTION:



is called as Gauerian * The Normal Distribution Distribution.

- * It is also called as bell-curre.
- * The major distailbution lieu in Mean.
- * In normal distribution Mean = Median = Mode.
- * The Normal Curve is symmetric about the Mean.
- * The Mean parameter server as a location

parameter.




The standard Deviation is a scale parameter :

Different conve gives différent scale.

Subject Incharge: Prof. Sarala Mary

Page No. 3

Department of CSE-Data Science | APSIT



A.P. SHAH INSTITUTE OF TECHNOLOGY

PARSHWANALH CHARITABLE TROST S

Department of Computer Science and Engineering Data Science



Semester:

Subject & Tatistics for AIDDS

Academic Year: 2023 2024

* From the diagram (1416) is one standard deviation away from the mean in night side. M-126 is two standard deviation away from the night side. M+86 is three standard deviation away from the singht side. It is called as influence

STANDARD NORMAL FORM AND QO-PLOTS:

* A Q-O plot, or quantile - quantile plot is a scatter plot created by plotting two sets of quintiles against one - another.

* It both seli of quantiles come from the some distribution, we see that - the points forming a Tine is noughly a straight line.

We mention here an example of a normal Q-Q plot when both seli of quantiles truely comes from normal dichribation.





Fig11 For Normal Dist. Example 1: + QQ-Plot

Department of CSE-Data Science | APSIT

Subject Incharge Prof. Sarala Mary



A.P. SHAH INSTITUTE OF TECHNOLOGY


Department of Computer Science and Engineering

Data Science



Subject Statistic for AIPDS Academic Year 2023 2024. Semester are often referred to as percentiles # Quantiles These are points in the dala set below which a Certain proportion of dalà-fall.

* For example, consider the bell-shaped slandard normal distribution with mean of 0. * The soth percentile is 0. * Half of the dala lie below o. * The 95% of dala lie below 1.64 from the * 8 ince the dala is normally distributed, the 00-plot generaler a straight line. 20 distribution) Straight 10 (9) maph 1 Sample Date Theoritical Quantiles Examplea. QQ-Plot - Graphs Subject Incharge Prof Sarata Mary Department of CSE-Data Science | APSIT



PARSHWANATH CHARITABLE TRUST'S

A.P. SHAH INSTITUTE OF TECHNOLOGY

Department of Computer Science and Engineering Data Science



Semester

subject slatistics for AIDS

Academic Year: 2023 20 24

* The enample 2 clearly describes how the


ad plat is generated.

* Y aru's takes the sample date and X-axis
lakes the normal Distribution Data

generalis a straight line then it is normal distribution.

LONG TAILED DISTRIBUTION!

of numbers is the portion of the distribution having many occurrence, and they are far from the "head" or cerdral part of the distribution.



Long tail.

In above graph shows long-tail distribution.

To the right (dotted part) is the long tail and to the left (lineal segment) are the few that dominate

Subject Incharge: Prof. Sarala Mary Page No. 6

Department of CSE-Data Science | APSIT