

W06-01

Slide 01: In this unit, we will introduce two fundamental subspaces associated with a matrix, namely the row space and the column space.

Slide 02: Given any $m \times n$ matrix \mathbf{A} ,

(#)

we will denote the rows of \mathbf{A} by \mathbf{r}_1 , \mathbf{r}_2 and so on till \mathbf{r}_m .

(#)

Each row of \mathbf{A} can be considered as a vector. Since the matrix \mathbf{A} has n columns, these vectors have n components and thus are vectors in \mathbb{R}^n .

(#)

The linear span of these row vectors of \mathbf{A} will thus be a subspace of \mathbb{R}^n .

(#)

We call this subspace the row space of the matrix \mathbf{A} .

Slide 03: We can also do the same for the columns of \mathbf{A} .

(#)

Let \mathbf{c}_1 , \mathbf{c}_2 until \mathbf{c}_n be the columns of \mathbf{A} .

(#)

We can consider these columns as vectors and since each of them has m components, these are vectors in \mathbb{R}^m .

(#)

The linear span of these column vectors of \mathbf{A} will be a subspace of \mathbb{R}^m .

(#)

We call this subspace the column space of the matrix \mathbf{A} .

Slide 04: Recall that when we transpose a matrix \mathbf{A} , the rows of \mathbf{A} becomes the columns of \mathbf{A}^T while the columns of \mathbf{A} becomes the rows of \mathbf{A}^T .

(#)

Thus, the row space of \mathbf{A} is the column space of \mathbf{A}^T .

(#)

While the column space of \mathbf{A} is the row space of \mathbf{A}^T .

Slide 05: Let us consider an example. The matrix \mathbf{A} shown here is a 4×3 matrix.

(#)

Thus the row space of \mathbf{A} is a subspace of \mathbb{R}^3

(#)

and the column space of \mathbf{A} is a subspace of \mathbb{R}^4 .

(#)

Note that if \mathbf{A} is not a square matrix, then the row space and column space of the matrix would contain totally different type of vectors, meaning that they would come from different Euclidean spaces.

Slide 06: Let us return to the matrix \mathbf{A} shown earlier, where the row space of \mathbf{A} is a subspace of \mathbb{R}^3 .

(#)

Let \mathbf{r}_1 to \mathbf{r}_4 denote the rows of \mathbf{A} , then the row space of \mathbf{A} is the linear span of \mathbf{r}_1 to \mathbf{r}_4 .

(#)

More precisely, the row space of \mathbf{A} contains all the possible linear combinations of $(2, 1, 0)$, $(1, -1, 3)$, $(-5, 1, 0)$ and $(1, 0, 1)$. Notice that when we write \mathbf{r}_1 as a vector, we use commas to indicate that this is a vector and not a row matrix.

(#)

We can rewrite the set to obtain an expression for an arbitrary vector in the row space of \mathbf{A} . Here, a, b, c, d can take on any real numbers.

Slide 07: For the same matrix \mathbf{A} , the column space is a subspace of \mathbb{R}^4 .

(#)

Once again, if we represent the columns of \mathbf{A} by \mathbf{c}_1 , \mathbf{c}_2 and \mathbf{c}_3 , then the column space of \mathbf{A} will be the linear span of \mathbf{c}_1 , \mathbf{c}_2 and \mathbf{c}_3 .

(#)

This is essentially the set of all linear combinations of $(2, 1, -5, 1)$, $(-1, -1, 1, 0)$ and $(0, 3, 0, 1)$.

(#)

The set can be rewritten to obtain an expression for an arbitrary vector in the column space of \mathbf{A} . Here, a, b, c can take on any real numbers.

Slide 08: Let us consider another matrix \mathbf{A} .

(#)

It is easy to see that the row space of \mathbf{A} is a subspace of \mathbb{R}^5 ,

(#)

and the column space of \mathbf{A} is a subspace of \mathbb{R}^3 .

(#)

We are now interested in finding a basis for the row space of \mathbf{A} .

Slide 09: The row space of \mathbf{A} is the linear span of the three row vectors of \mathbf{A} .

(#)

In other words, we already know that these three row vectors span the row space. If these three row vectors are also linearly independent, then they will form a basis for the row space of \mathbf{A} .

Slide 10: To see if they are indeed linearly independent, we set up the vector equation as shown.

(#)

By observation, looking at the first component leads to the conclusion that $a = 0$.

(#)

Looking at the second component tells us that $b = 0$.

(#)

And lastly looking at the third component tells us that $c = 0$.

(#)

Since the vector equation has only the trivial solution, the three rows of \mathbf{A} are linearly independent and thus form a basis for the row space of \mathbf{A} .

Slide 11: Observe that for the matrix \mathbf{A} which we have just seen, it was easy for us to conclude that the row vectors are linearly independent by making some quick observations. This was possible because \mathbf{A} is actually in row-echelon form. The position of the leading entry in each row is such that we can quickly conclude that the vector equation has only the trivial solution.

(#)

This is not a coincidence. For any matrix \mathbf{R} that is in row-echelon form, if we would like to find a basis for the row space of \mathbf{R} ,

Slide 12: The non zero rows of \mathbf{R} will always be linearly independent and thus forms a basis for the row space of \mathbf{R} . Notice that the zero rows found at the bottom of the matrix are zero vectors which can be disregarded.

Slide 13: To summarise the main points.

(#)

We defined what is the row space and column space of a $m \times n$ matrix \mathbf{A} .

(#)

We also saw that if the rows of \mathbf{A} are linearly independent, since they span the row space of \mathbf{A} , they would form a basis for the row space.

(#)

In the event that a matrix \mathbf{R} is in row-echelon form, then to find a basis for the row space of \mathbf{R} is easy. We simply take the non zero rows of \mathbf{R} in this case.