


Ayachi Zammel — hot and cold spots

Getis-Ord Gi* z-score, imada level

- Gi* z-score
- cold, $z < -2.58$ (1)
 - cold, -2.58 to -1.96 (67)
 - cold, -1.96 to -1.65 (185)
 - not notable, $|z| < 1.65$ (1453)
 - hot, 1.65 to 1.96 (55)
 - hot, 1.96 to 2.58 (77)
 - hot, $z > 2.58$ (204)
 - significant after FDR (52)

2042 mapped units



Gi* includes the unit itself, so it measures whether the local total is unusual rather than how the unit relates to its neighbours: it has no outlier classes and is the more legible of the two cluster statistics — but not a second one: under conditional permutation it selects the same units as the LISA map beside it, because both hold the unit's own value fixed and so test the same neighbourhood. What it adds is the continuous surface the quadrant map discards. Positive is a hot spot, negative a cold spot; the ramp carries the z-score in fixed bands rather than quantiles, so a shade means the same thing on every one of these maps; the observed range here is -2.6 to +12.8. The outline marks the 52 units that survive Benjamini-Hochberg at $q = 0.05$ — 52 hot, 0 cold. global Moran's $I = +0.591$ ($p < 0.0001$, 9,999 permutations, pseudo $z = +44.0$). Source: ISIE procès-verbaux, 9,419 certified stations · boundaries OCHA/HDX COD-AB (CC BY-IGO) · queen contiguity from shared boundary vertices