

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

ARICLAIM 40 mg hard gastro-resistant capsules.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

The active ingredient in ARICLAIM is duloxetine.
Each capsule contains 40 mg of duloxetine as duloxetine hydrochloride.

Excipients: sucrose.
For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard gastro-resistant capsule.

The capsule has an opaque orange body, imprinted with '40mg' and an opaque blue cap, imprinted with '9545'.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

ARICLAIM is indicated for women for the treatment of moderate to severe Stress Urinary Incontinence (SUI), (see section 5.1).

4.2 Posology and method of administration

The recommended dose of ARICLAIM is 40 mg twice daily without regard to meals. After 2-4 weeks of treatment, patients should be re-assessed in order to evaluate the benefit and tolerability of the therapy. Some patients may benefit from starting treatment at a dose of 20 mg twice daily for two weeks before increasing to the recommended dose of 40 mg twice daily. Dose escalation may decrease, though not eliminate, the risk of nausea and dizziness.
However, limited data are available to support the efficacy of ARICLAIM 20 mg twice daily.
A 20 mg capsule is also available.

The efficacy of ARICLAIM has not been evaluated for longer than 3 months in placebo-controlled studies. The benefit of treatment should be re-assessed at regular intervals.

Combining ARICLAIM with a pelvic floor muscle training (PFMT) program may be more effective than either treatment alone. It is recommended that consideration be given to concomitant PFMT.

Hepatic insufficiency:

ARICLAIM should not be used in women with liver disease resulting in hepatic impairment (see section 4.3).

Renal insufficiency:

No dosage adjustment is necessary for patients with mild or moderate renal dysfunction (creatinine clearance 30 to 80 ml/min).

Elderly:

Caution should be exercised when treating the elderly.

Children and adolescents:

The safety and efficacy of duloxetine in patients in these age groups have not been studied. Therefore, administration of ARICLAIM to children and adolescents is not recommended.

Discontinuation of treatment:

Abrupt discontinuation should be avoided. When stopping treatment with ARICLAIM the dose should be gradually reduced over a period of at least one to two weeks in order to reduce the risk of withdrawal reactions (see sections 4.4 and 4.8). If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the physician may continue decreasing the dose, but at a more gradual rate.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

Liver disease resulting in hepatic impairment (see section 5.2).

Pregnancy and lactation (see section 4.6).

ARICLAIM should not be used in combination with nonselective, irreversible Monoamine Oxidase Inhibitors - MAOIs (see section 4.5).

ARICLAIM should not be used in combination with CYP1A2 inhibitors, like fluvoxamine, ciprofloxacin or enoxacin since the combination results in elevated plasma concentrations of duloxetine (see section 4.5).

Severe renal impairment (creatinine clearance <30 ml/min) (see section 4.4).

The initiation of treatment with ARICLAIM is contraindicated in patients with uncontrolled hypertension that could expose patients to a potential risk of hypertensive crisis (see sections 4.4 and 4.8).

4.4 Special warnings and precautions for use

Mania and Seizures

ARICLAIM should be used with caution in patients with a history of mania or a diagnosis of bipolar disorder, and/or seizures.

Use with antidepressants

The use of ARICLAIM in combination with antidepressants (especially with SSRI, SNRI and reversible MAOIs) is not recommended (see below “Depression, suicidal ideation and behaviour” and Section 4.5).

Mydriasis

Mydriasis has been reported in association with duloxetine, therefore, caution should be used when prescribing duloxetine in patients with increased intraocular pressure, or those at risk of acute narrow-angle glaucoma.

Blood pressure and heart rate

Duloxetine has been associated with an increase in blood pressure and clinically significant hypertension in some patients. This may be due to the noradrenergic effect of duloxetine. Cases of hypertensive crisis have been reported with duloxetine, especially in patients with pre-existing hypertension. Therefore, in patients with known hypertension and/or other cardiac disease, blood pressure monitoring is recommended, especially during the first month of treatment. Duloxetine should be used with caution in patients whose conditions could be compromised by an increased heart rate or by an increase in blood pressure. Caution should also be exercised when duloxetine is used with drugs that may impair its metabolism (see section 4.5). For patients who experience a sustained

increase in blood pressure while receiving duloxetine either dose reduction or gradual discontinuation should be considered (see section 4.8). In patients with uncontrolled hypertension duloxetine should not be initiated (see section 4.3).

Renal impairment

Increased plasma concentrations of duloxetine occur in patients with severe renal impairment on haemodialysis (creatinine clearance <30 ml/min). For patients with severe renal impairment, see section 4.3. See section 4.2 for information on patients with mild or moderate renal dysfunction.

Sucrose

ARICLAIM hard gastro-resistant capsules contain sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine.

Haemorrhage

There have been reports of bleeding abnormalities, such as ecchymoses, purpura and gastrointestinal haemorrhage with selective serotonin reuptake inhibitors (SSRIs) and serotonin/norepinephrine reuptake inhibitors (SNRIs). Caution is advised in patients taking anticoagulants and/or medicinal products known to affect platelet function, and in patients with known bleeding tendencies.

Discontinuation of treatment

Withdrawal symptoms when treatment is discontinued are common, particularly if discontinuation is abrupt (see section 4.8). In a clinical trial adverse events seen on abrupt treatment discontinuation occurred in approximately 44% of patients treated with ARICLAIM and 24% of patients taking placebo.

The risk of withdrawal symptoms seen with SSRI's and SNRI's may be dependent on several factors including the duration and dose of therapy and the rate of dose reduction. The most commonly reported reactions are listed in section 4.8. Generally the symptoms are mild to moderate, however, in some patients they may be severe in intensity. They usually occur within the first few days of discontinuing treatment, but there have been very rare reports of such symptoms in patients who have inadvertently missed a dose. Generally these symptoms are self-limiting and usually resolve within 2 weeks, though in some individuals they may be prolonged (2-3 months or more). It is therefore advised that duloxetine should be gradually tapered when discontinuing treatment over a period of no less than 2 weeks, according to the patient's needs (see section 4.2).

Hyponatremia

Hyponatremia has been reported rarely, predominantly in the elderly, when administering ARICLAIM. Caution is required in patients at increased risk for hyponatremia; such as elderly, cirrhotic, or dehydrated patients or patients treated with diuretics. Hyponatremia may be due to a syndrome of inappropriate anti-diuretic hormone secretion (SIADH).

Depression, suicidal ideation and behaviour

Although ARICLAIM is not indicated for the treatment of depression, its active ingredient (duloxetine) also exists as an antidepressant medication. Isolated cases of suicidal ideation and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation. Physicians should encourage patients to report any distressing thoughts or feelings or depressive symptoms at any time. If while on ARICLAIM therapy, the patient develops agitation or depressive symptoms, specialised medical advice should be sought, as depression is a serious medical condition. If a decision to initiate antidepressant pharmacological therapy is taken, the gradual discontinuation of ARICLAIM is recommended (see Section 4.2).

Use in children and adolescents under 18 years of age

No clinical trials have been conducted with duloxetine in paediatric populations. ARICLAIM should not be used in the treatment of children and adolescents under the age of 18 years. Suicide-related behaviours (suicide attempts and suicidal thoughts), and hostility (predominantly aggression, oppositional behaviour and anger), were more frequently observed in clinical trials among children and adolescents treated with antidepressants compared to those treated with placebo. Long-term safety

data in children and adolescents concerning growth, maturation and cognitive and behavioural development are lacking.

Medicinal products containing duloxetine

Duloxetine is used under different trademarks in several indications (treatment of diabetic neuropathic pain, major depressive episodes as well as stress urinary incontinence). The use of more than one of these products concomitantly should be avoided.

Hepatitis/increased liver enzymes

Cases of liver injury, including severe elevations of liver enzymes (>10 times upper limit of normal), hepatitis and jaundice have been reported with duloxetine (see section 4.8). Most of them occurred during the first months of treatment. The pattern of liver damage was predominantly hepatocellular. Duloxetine should be used with caution in patients treated with other drugs associated with hepatic injury.

Akathisia/psychomotor restlessness

The use of duloxetine has been associated with the development of akathisia, characterised by a subjectively unpleasant or distressing restlessness and need to move often accompanied by an inability to sit or stand still. This is most likely to occur within the first few weeks of treatment. In patients who develop these symptoms, increasing the dose may be detrimental.

4.5 Interaction with other medicinal products and other forms of interaction

Monoamine oxidase inhibitors (MAOIs): due to the risk of serotonin syndrome, ARICLAIM should not be used in combination with nonselective, irreversible monoamine oxidase inhibitors (MAOIs), or within at least 14 days of discontinuing treatment with an MAOI. Based on the half-life of duloxetine, at least 5 days should be allowed after stopping ARICLAIM before starting an MAOI (see section 4.3).

Serotonin syndrome: in rare cases, serotonin syndrome has been reported in patients using SSRIs concomitantly with serotonergic medicinal products. The use of ARICLAIM in combination with serotonergic antidepressants like SSRIs, tricyclics like clomipramine or amitriptyline, venlafaxine, or triptans, tramadol and tryptophan is not recommended.

CNS drugs: caution is advised when ARICLAIM is taken in combination with other centrally acting drugs or substances, including alcohol and sedative drugs (benzodiazepines, morphinomimetics, antipsychotics, phenobarbital, sedative antihistamines).

Effect of duloxetine on other drugs

Medicinal products metabolised by CYP1A2: The pharmacokinetics of theophylline, a CYP1A2 substrate, were not significantly affected by co-administration with duloxetine (60 mg twice daily).

Medicinal products metabolised by CYP2D6: Duloxetine is a moderate inhibitor of CYP2D6. When duloxetine was administered at a dose of 60 mg twice daily with a single dose of desipramine, a CYP2D6 substrate, the AUC of desipramine increased 3-fold. The co-administration of duloxetine (40 mg twice daily) increases steady state AUC of tolterodine (2 mg twice daily) by 71 %, but does not affect the pharmacokinetics of its active 5-hydroxyl metabolite and no dosage adjustment is recommended. Caution is advised if ARICLAIM is co-administered with medicinal products that are predominantly metabolised by CYP2D6 (risperidone, tricyclic antidepressants [TCAs] such as nortriptyline, amitriptyline, and imipramine) particularly if they have a narrow therapeutic index (such as flecainide, propafenone and metoprolol).

Oral contraceptives and other steroidal agents: results of *in vitro* studies demonstrate that duloxetine does not induce the catalytic activity of CYP3A. Specific *in vivo* drug interaction studies have not been performed.

Anticoagulants and antiplatelet agents: Caution should be exercised when duloxetine is combined with oral anticoagulants or antiplatelet agents due to a potential increased risk of bleeding.

Furthermore, increases in INR values have been reported when duloxetine was co-administered with warfarin.

Effects of other drugs on duloxetine

Antacids and H₂ antagonists: co-administration of ARICLAIM with aluminium- and magnesium-containing antacids or with famotidine had no significant effect on the rate or extent of duloxetine absorption after administration of a 40 mg oral dose.

Inhibitors of CYP1A2: because CYP1A2 is involved in duloxetine metabolism, concomitant use of ARICLAIM with potent inhibitors of CYP1A2 is likely to result in higher concentrations of duloxetine. Fluvoxamine (100 mg once daily), a potent inhibitor of CYP1A2, decreased the apparent plasma clearance of duloxetine by about 77% and increased AUC₀₋₁ 6-fold. Therefore ARICLAIM should not be administered in combination with potent inhibitors of CYP1A2 like fluvoxamine (see section 4.3).

Inducers of CYP1A2: Population pharmacokinetic studies analyses have shown that smokers have almost 50% lower plasma concentrations of duloxetine compared with non-smokers.

4.6 Pregnancy and lactation

Pregnancy

There are no data on the use of duloxetine in pregnant women. Studies in animals have shown reproductive toxicity at systemic exposure levels (AUC) of duloxetine lower than the maximum clinical exposure (see section 5.3). The potential risk for humans is unknown. Withdrawal symptoms may occur in the neonate after maternal duloxetine use near term. ARICLAIM is contraindicated during pregnancy (see section 4.3).

Breast feeding

Duloxetine is very weakly excreted into human milk based on a study of 6 lactating patients, who did not breast feed their children. The estimated daily infant dose on a mg/kg basis is approximately 0.14% of the maternal dose (see section 5.2). As the safety of duloxetine in infants is not known, the use of ARICLAIM while breast-feeding is not recommended.

4.7 Effects on ability to drive and use machines

Although in controlled studies duloxetine has not been shown to impair psychomotor performance, cognitive function, or memory, it may be associated with sedation and dizziness. Therefore, patients should be cautioned about their ability to drive a car or operate hazardous machinery.

4.8 Undesirable effects

Table 1 gives the adverse reactions observed from spontaneous reporting and in placebo-controlled clinical trials (comprising a total of 7977 patients, 4371 on duloxetine and 3606 on placebo) in SUI and other lower urinary tract disorders.

The most commonly reported adverse events in patients treated with ARICLAIM in clinical trials in SUI and other lower urinary tract disorders were nausea, dry mouth and fatigue. The data analysis of four 12-week, placebo-controlled clinical trials in patients with SUI, including 958 duloxetine-treated and 955 placebo-treated patients, showed that the onset of the reported adverse events typically occurred in the first week of therapy. However, the majority of the most frequent adverse events were mild to moderate and resolved within 30 days of occurrence (e.g. nausea).

Table 1: Adverse reactions

Frequency estimate: Very common (≥10%), common (≥1% and <10%), uncommon (≥0.1% and <1%), rare (≥0.01% and <0.1%), very rare (<0.01%), frequency not known (data from spontaneous reports)

Very common	Common	Uncommon	Rare	Very Rare	Frequency not known
<i>Infections and infestations</i>					
		Laryngitis			
<i>Immune system disorders</i>					
		Hyper-sensitivity disorder			Anaphylactic reaction
<i>Endocrine disorders</i>					
		Hypo-thyroidism			
<i>Metabolism and Nutrition Disorders</i>					
	Appetite decreased	Dehydration			Hyponatremia, SIADH
<i>Psychiatric Disorders</i>					
	Insomnia Sleep disorder Anxiety Libido decreased Agitation	Bruxism Disorientation Orgasm abnormal Apathy Abnormal dreams			Hallucinations Mania
<i>Nervous System Disorders</i>					
	Headache Dizziness Tremor Nervousness Lethargy Somnolence Paraesthesia	Dysgeusia Disturbance in attention	Myoclonus Dyskinesia		Serotonin syndrome Extrapyramidal symptoms Convulsions Akathisia Psychomotor restlessness
<i>Eye Disorders</i>					
	Blurred vision	Mydriasis Visual disturbance	Glaucoma		
<i>Ear and Labyrinth Disorders</i>					
	Vertigo	Ear pain			
<i>Cardiac Disorders</i>					
	Palpitations	Tachycardia			Supra-ventricular arrhythmia, mainly atrial fibrillation
<i>Vascular Disorders</i>					
	Hot flush	Flushing Blood pressure increase Syncope ¹	Peripheral coldness Orthostatic hypotension ¹ Hypertensive crisis		Hypertension
<i>Respiratory, thoracic and mediastinal disorders</i>					
		Yawning	Throat tightness		

<i>Gastrointestinal Disorders</i>					
Nausea (22%) Dry mouth (11.2%)	Constipation Diarrhoea Vomiting Dyspepsia	Eructation Gastroenteritis Stomatitis Halitosis Gastritis Flatulence			
<i>Hepato-biliary disorders</i>					
		Elevated liver enzymes (ALT, AST, alkaline phosphatase) Hepatitis ² Acute liver injury			Jaundice Hepatic failure
<i>Skin and Subcutaneous Tissue Disorders</i>					
	Sweating increased	Night sweats Cold sweat Rash	Photo-sensitivity reactions		Angioneurotic oedema Stevens-Johnson Syndrome Urticaria
<i>Muscoskeletal and connective tissue disorders</i>					
		Muscle tightness Musculo-skeletal pain Trismus	Muscle twitching		
<i>Renals and Urinary Disorders</i>					
		Nocturia Urinary hesitation Urine odour abnormal Dysuria			Urinary retention
<i>Reproductive System and Breast Disorders</i>					
		Menopausal symptoms			
<i>General Disorders and Administration Site Conditions</i>					
Fatigue (10.3%)	Pruritus Weakness Abdominal pain Chills	Feeling hot Malaise Thirst Feeling abnormal	Feeling cold		Chest pain
<i>Investigations</i>					
		Weight increase Weight decrease Creatinine phosphokinase increase			

		Blood cholesterol increased			
--	--	-----------------------------	--	--	--

¹Cases of orthostatic hypotension and syncope have been reported especially at the initiation of treatment

²See section 4.4

Discontinuation of duloxetine (particularly when abrupt) commonly leads to withdrawal symptoms. Dizziness, sensory disturbances (including paraesthesia), sleep disturbances (including insomnia and intense dreams), agitation or anxiety, nausea and/or vomiting, tremor and headache are the most commonly reported reactions.

Generally, for SSRIs and SNRIs, these events are mild to moderate and self-limiting, however, in some patients they may be severe and/or prolonged. It is therefore advised that when duloxetine treatment is no longer required, gradual discontinuation by dose tapering should be carried out (see sections 4.2 and 4.4).

Electrocardiograms were obtained from 755 duloxetine-treated patients with SUI and 779 placebo-treated patients in 12-week clinical trials. The heart rate-corrected QT interval in duloxetine-treated patients did not differ from that seen in placebo-treated patients.

In the 12 week acute phase of three clinical trials of duloxetine in patients with diabetic neuropathic pain, small but statistically significant increases in fasting blood glucose were observed in duloxetine-treated patients. HbA_{1c} was stable in both duloxetine-treated and placebo-treated patients. In the extension phase of these studies, which lasted up to 52 weeks, there was an increase in HbA_{1c} in both the duloxetine and routine care groups, but the mean increase was 0.3% greater in the duloxetine-treated group. There was also a small increase in fasting blood glucose and in total cholesterol in duloxetine-treated patients while those laboratory tests showed a slight decrease in the routine care group.

4.9 Overdose

There is limited clinical experience with duloxetine overdose in humans. Cases of overdoses, alone or in combination with other drugs, with duloxetine doses of almost 2000 mg were reported. Some fatalities have occurred, primarily with mixed overdoses, but also with duloxetine alone at a dose of approximately 1000 mg. Signs and symptoms of overdose (mostly with mixed drugs) included serotonin syndrome, somnolence, vomiting and seizures.

No specific antidote for duloxetine is known but if serotonin syndrome ensues, specific treatment (such as with cyproheptadine and/or temperature control) may be considered.. A free airway should be established. Monitoring of cardiac and vital signs is recommended, along with appropriate symptomatic and supportive measures. Gastric lavage may be indicated if performed soon after ingestion or in symptomatic patients. Activated charcoal may be useful in limiting absorption. Duloxetine has a large volume of distribution and forced diuresis, haemoperfusion, and exchange perfusion are unlikely to be beneficial.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Other antidepressants. ATC code: N06AX21

Duloxetine is a combined serotonin (5-HT) and norepinephrine (NE) reuptake inhibitor. It weakly inhibits dopamine reuptake with no significant affinity for histaminergic, dopaminergic, cholinergic and adrenergic receptors.

In animal studies, increased levels of 5-HT and NE in the sacral spinal cord, lead to increased urethral tone via enhanced pudendal nerve stimulation to the urethral striated sphincter muscle only during the storage phase of the micturition cycle. A similar mechanism in women is believed to result in stronger urethral closure during urine storage with physical stress that could explain the efficacy of duloxetine in the treatment of women with SUI.

The efficacy of duloxetine 40 mg given twice daily in the treatment of SUI was established in four double-blind, placebo-controlled studies, that randomised 1913 women (22 to 83 years) with SUI; of these, 958 patients were randomised to duloxetine and 955 to placebo. The primary efficacy measures were Incontinence Episode Frequency (IEF) from diaries and an incontinence specific quality of life questionnaire score (I-QOL).

Incontinence Episode Frequency: in all four studies the duloxetine-treated group had a 50% or greater median decrease in IEF compared with 33% in the placebo-treated group. Differences were observed at each visit after 4 weeks (duloxetine 54%; and placebo 22%), 8 weeks (52% and 29%), and 12 weeks (52% and 33%) of medication.

In an additional study limited to patients with severe SUI, all responses with duloxetine were achieved within 2 weeks.

The efficacy of ARICLAIM has not been evaluated for longer than 3 months in placebo-controlled studies. The clinical benefit of ARICLAIM compared with placebo has not been demonstrated in women with mild SUI, defined in randomised trials as those with IEF < 14 per week. In these women, ARICLAIM may provide no benefit beyond that afforded by more conservative behavioural interventions.

Quality of Life: Incontinence Quality of Life (I-QOL) questionnaire scores were significantly improved in the duloxetine-treated patient group compared with the placebo-treated group (9.2 versus 5.9 score improvement, $p < .001$). Using a global improvement scale (PGI), significantly more women using duloxetine considered their symptoms of stress incontinence to be improved with treatment compared with women using placebo (64.6% versus 50.1%, $p < .001$).

ARICLAIM and Prior Continence Surgery: there are limited data that suggest that the benefits of ARICLAIM are not diminished in women with stress urinary incontinence who have previously undergone continence surgery.

ARICLAIM and Pelvic Floor Muscle Training (PFMT): during a 12-week blinded, randomised, controlled study, ARICLAIM demonstrated greater reductions in IEF compared with either placebo treatment or with PFMT alone. Combined therapy (duloxetine + PFMT) showed greater improvement in both pad use and condition-specific quality of life measures than ARICLAIM alone or PFMT alone.

5.2 Pharmacokinetic properties

Duloxetine is administered as a single enantiomer. Duloxetine is extensively metabolised by oxidative enzymes (CYP1A2 and the polymorphic CYP2D6), followed by conjugation. The pharmacokinetics of duloxetine demonstrate large intersubject variability (generally 50-60%), partly due to gender, age, smoking status and CYP2D6 metaboliser status.

Duloxetine is well absorbed after oral administration with a C_{max} occurring 6 hours post dose. The absolute oral bioavailability of duloxetine ranged from 32% to 80% (mean of 50%; N=8 subjects). Food delays the time to reach the peak concentration from 6 to 10 hours and it marginally decreases the extent of absorption (approximately 11%).

Duloxetine is approximately 96% bound to human plasma proteins. Duloxetine binds to both albumin and alpha-1 acid glycoprotein. Protein binding is not affected by renal or hepatic impairment.

Duloxetine is extensively metabolised and the metabolites are excreted principally in urine. Both CYP2D6 and CYP1A2 catalyse the formation of the two major metabolites glucuronide conjugate of 4-hydroxy duloxetine and sulphate conjugate of 5-hydroxy,6-methoxy duloxetine. Based upon *in vitro* studies, the circulating metabolites of duloxetine are considered pharmacologically inactive. The pharmacokinetics of duloxetine in patients who are poor metabolisers with respect to CYP2D6 has not been specifically investigated. Limited data suggest that the plasma levels of duloxetine are higher in these patients.

The elimination half-life of duloxetine after an oral dose ranges from 8 to 17 hours (mean of 12 hours). After an intravenous dose the plasma clearance of duloxetine ranges from 22 l/hr to 46 l/hr (mean of 36 l/hr) After an oral dose the apparent plasma clearance of duloxetine ranges from 33 to 261 l/hr (mean 101 l/hr).

Special populations:

Age: pharmacokinetic differences have been identified between younger and elderly females (≥ 65 years) (AUC increases by about 25% and half-life is about 25% longer in the elderly), although the magnitude of these changes is not sufficient to justify adjustments to the dose.

Renal impairment: end stage renal disease (ESRD) patients receiving dialysis had a 2-fold higher duloxetine C_{max} and AUC values compared to healthy subjects. Pharmacokinetic data on duloxetine is limited in patients with mild or moderate renal impairment.

Hepatic insufficiency: moderate liver disease (Child Pugh Class B) affected the pharmacokinetics of duloxetine. Compared with healthy subjects, the apparent plasma clearance of duloxetine was 79% lower, the apparent terminal half-life was 2.3 times longer, and the AUC was 3.7 times higher in patients with moderate liver disease. The pharmacokinetics of duloxetine and its metabolites have not been studied in patients with mild or severe hepatic insufficiency.

Nursing mothers: The disposition of duloxetine was studied in 6 lactating women who were at least 12-weeks postpartum. Duloxetine is detected in breast milk, and steady-state concentrations in breast milk are about one-fourth those in plasma. The amount of duloxetine in breast milk is approximately 7 µg/day while on 40 mg twice daily dosing. Lactation did not influence duloxetine pharmacokinetics.

5.3 Preclinical safety data

Duloxetine was not genotoxic in a standard battery of tests and was not carcinogenic in rats. Multinucleated cells were seen in the liver in the absence of other histopathological changes in the rat carcinogenicity study. The underlying mechanism and the clinical relevance are unknown.

Female mice receiving duloxetine for 2 years had an increased incidence of hepatocellular adenomas and carcinomas at the high dose only (144 mg/kg/day), but these were considered to be secondary to hepatic microsomal enzyme induction. The relevance of this mouse data to humans is unknown. Female rats receiving duloxetine before and during mating and early pregnancy had a decrease in maternal food consumption and body weight, oestrous cycle disruption, decreased live birth indices and progeny survival, and progeny growth retardation at systemic exposure levels estimated to be at the most at maximum clinical exposure (AUC). In an embryotoxicity study in the rabbit, a higher incidence of cardiovascular and skeletal malformations was observed at systemic exposure levels below the maximum clinical exposure (AUC). No malformations were observed in another study testing a higher dose of a different salt of duloxetine. In pre/postnatal toxicity study in the rat, duloxetine induced adverse behavioural effects in the offspring at systemic exposures levels below maximum clinical exposure (AUC).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content:

Hypromellose.
Hydroxypropyl methylcellulose acetate succinate
Sucrose
Sugar spheres
Talc
Titanium dioxide (E171)
Triethyl citrate.

Capsule Shell:

Gelatin
Sodium Lauryl Sulfate
Titanium Dioxide (E171)
Indigo Carmine (E132)
Red Iron oxide (E172)
Yellow Iron Oxide (E172)
Edible black ink.

Edible Ink:

Black Iron Oxide-Synthetic (E172)
Propylene glycol
Shellac.

Capsule Shell Cap colour:

Opaque Blue

Capsule Shell Body colour:

Opaque Orange

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Store in the original package. Do not store above 30°C.

6.5 Nature and contents of container

Polyvinylchloride (PVC), Polyethylene (PE), and Polychlorotrifluoroethylene (PCTFE) blister sealed with an aluminum foil.

Packs of 28, 56, 98, 140 and 196 (2x98) capsules.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements.

7. MARKETING AUTHORISATION HOLDER

Eli Lilly Nederland BV, Grootslag 1-5, NL-3991 RA Houten, The Netherlands.

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/04/283/002

EU/1/04/283/003

EU/1/04/283/004

EU/1/04/283/005

EU/1/04/283/006

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

11 August 2004

10. DATE OF REVISION OF THE TEXT

1. NAME OF THE MEDICINAL PRODUCT

ARICLAIM 20 mg hard gastro-resistant capsules.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

The active ingredient in ARICLAIM is duloxetine.

Each capsule contains 20 mg of duloxetine as duloxetine hydrochloride.

Excipients: sucrose.

For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard gastro-resistant capsule.

The capsule has an opaque blue body, imprinted with '20 mg' and an opaque blue cap, imprinted with '9544'.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

ARICLAIM is indicated for women for the treatment of moderate to severe Stress Urinary Incontinence (SUI), (see section 5.1).

4.2 Posology and method of administration

The recommended dose of ARICLAIM is 40 mg twice daily without regard to meals. After 2-4 weeks of treatment, patients should be re-assessed in order to evaluate the benefit and tolerability of the therapy. Some patients may benefit from starting treatment at a dose of 20 mg twice daily for two weeks before increasing to the recommended dose of 40 mg twice daily. Dose escalation may decrease, though not eliminate, the risk of nausea and dizziness. However, limited data are available to support the efficacy of ARICLAIM 20 mg twice daily.

The efficacy of ARICLAIM has not been evaluated for longer than 3 months in placebo-controlled studies. The benefit of treatment should be re-assessed at regular intervals.

Combining ARICLAIM with a pelvic floor muscle training (PFMT) program may be more effective than either treatment alone. It is recommended that consideration be given to concomitant PFMT.

Hepatic insufficiency:

ARICLAIM should not be used in women with liver disease resulting in hepatic impairment (see section 4.3).

Renal insufficiency:

No dosage adjustment is necessary for patients with mild or moderate renal dysfunction (creatinine clearance 30 to 80 ml/min).

Elderly:

Caution should be exercised when treating the elderly.

Children and adolescents:

The safety and efficacy of duloxetine in patients in these age groups have not been studied. Therefore, administration of ARICLAIM to children and adolescents is not recommended.

Discontinuation of treatment:

Abrupt discontinuation should be avoided. When stopping treatment with ARICLAIM the dose should be gradually reduced over a period of at least one to two weeks in order to reduce the risk of withdrawal reactions (see sections 4.4 and 4.8). If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the physician may continue decreasing the dose, but at a more gradual rate.).

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

Liver disease resulting in hepatic impairment (see section 5.2).

Pregnancy and lactation (see section 4.6).

ARICLAIM should not be used in combination with nonselective, irreversible Monoamine Oxidase Inhibitors - MAOIs (see section 4.5).

ARICLAIM should not be used in combination with CYP1A2 inhibitors, like fluvoxamine, ciprofloxacin or enoxacin since the combination results in elevated plasma concentrations of duloxetine (see section 4.5).

Severe renal impairment (creatinine clearance <30 ml/min) (see section 4.4).

The initiation of treatment with ARICLAIM is contraindicated in patients with uncontrolled hypertension that could expose patients to a potential risk of hypertensive crisis (see sections 4.4 and 4.8).

4.4 Special warnings and precautions for use

Mania and Seizures

ARICLAIM should be used with caution in patients with a history of mania or a diagnosis of bipolar disorder, and/or seizures.

Use with antidepressants

The use of ARICLAIM in combination with antidepressants (especially with SSRI, SNRI and reversible MAOIs) is not recommended (see below “Depression, suicidal ideation and behaviour” and Section 4.5).

Mydriasis

Mydriasis has been reported in association with duloxetine, therefore, caution should be used when prescribing duloxetine in patients with increased intraocular pressure, or those at risk of acute narrow-angle glaucoma.

Blood pressure and heart rate

Duloxetine has been associated with an increase in blood pressure and clinically significant hypertension in some patients. This may be due to the noradrenergic effect of duloxetine. Cases of hypertensive crisis have been reported with duloxetine, especially in patients with pre-existing hypertension. Therefore, in patients with known hypertension and/or other cardiac disease, blood pressure monitoring is recommended, especially during the first month of treatment. Duloxetine should be used with caution in patients whose conditions could be compromised by an increased heart rate or by an increase in blood pressure. Caution should also be exercised when duloxetine is used

with drugs that may impair its metabolism (see section 4.5). For patients who experience a sustained increase in blood pressure while receiving duloxetine either dose reduction or gradual discontinuation should be considered (see section 4.8). In patients with uncontrolled hypertension duloxetine should not be initiated (see section 4.3).

Renal impairment

Increased plasma concentrations of duloxetine occur in patients with severe renal impairment on haemodialysis (creatinine clearance <30 ml/min). For patients with severe renal impairment, see section 4.3. See section 4.2 for information on patients with mild or moderate renal dysfunction.

Sucrose

ARICLAIM hard gastro-resistant capsules contain sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine.

Haemorrhage

There have been reports of bleeding abnormalities, such as ecchymoses, purpura and gastrointestinal haemorrhage with selective serotonin reuptake inhibitors (SSRIs) and serotonin/norepinephrine reuptake inhibitors (SNRIs). Caution is advised in patients taking anticoagulants and/or medicinal products known to affect platelet function, and in patients with known bleeding tendencies.

Discontinuation of treatment

Withdrawal symptoms when treatment is discontinued are common, particularly if discontinuation is abrupt (see section 4.8). In a clinical trial, adverse events seen on abrupt treatment discontinuation occurred in approximately 44% of patients treated with ARICLAIM and 24% of patients taking placebo.

The risk of withdrawal symptoms seen with SSRI's and SNRI's may be dependent on several factors including the duration and dose of therapy and the rate of dose reduction. The most commonly reported reactions are listed in section 4.8. Generally the symptoms are mild to moderate, however, in some patients they may be severe in intensity. They usually occur within the first few days of discontinuing treatment, but there have been very rare reports of such symptoms in patients who have inadvertently missed a dose. Generally these symptoms are self-limiting and usually resolve within 2 weeks, though in some individuals they may be prolonged (2-3 months or more). It is therefore advised that duloxetine should be gradually tapered when discontinuing treatment over a period of no less than 2 weeks, according to the patient's needs (see section 4.2).

Hyponatremia

Hyponatremia has been reported rarely, predominantly in the elderly, when administering ARICLAIM. Caution is required in patients at increased risk for hyponatremia; such as elderly, cirrhotic, or dehydrated patients or patients treated with diuretics. Hyponatremia may be due to a syndrome of inappropriate anti-diuretic hormone secretion (SIADH).

Depression, suicidal ideation and behaviour

Although ARICLAIM is not indicated for the treatment of depression, its active ingredient (duloxetine) also exists as an antidepressant medication. Isolated cases of suicidal ideation and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation. Physicians should encourage patients to report any distressing thoughts or feelings or depressive symptoms at any time. If while on ARICLAIM therapy, the patient develops agitation or depressive symptoms, specialised medical advice should be sought, as depression is a serious medical condition. If a decision to initiate antidepressant pharmacological therapy is taken, the gradual discontinuation of ARICLAIM is recommended (see Section 4.2).

Use in children and adolescents under 18 years of age

No clinical trials have been conducted with duloxetine in paediatric populations. ARICLAIM should not be used in the treatment of children and adolescents under the age of 18 years. Suicide-related behaviours (suicide attempts and suicidal thoughts), and hostility (predominantly aggression, oppositional behaviour and anger), were more frequently observed in clinical trials among children

and adolescents treated with antidepressants compared to those treated with placebo. Long-term safety data in children and adolescents concerning growth, maturation and cognitive and behavioural development are lacking.

Medicinal products containing duloxetine

Duloxetine is used under different trademarks in several indications (treatment of diabetic neuropathic pain, major depressive episodes as well as stress urinary incontinence). The use of more than one of these products concomitantly should be avoided.

Hepatitis/increased liver enzymes

Cases of liver injury, including severe elevations of liver enzymes (>10 times upper limit of normal), hepatitis and jaundice have been reported with duloxetine (see section 4.8). Most of them occurred during the first months of treatment. The pattern of liver damage was predominantly hepatocellular. Duloxetine should be used with caution in patients treated with other drugs associated with hepatic injury.

Akathisia/psychomotor restlessness

The use of duloxetine has been associated with the development of akathisia, characterised by a subjectively unpleasant or distressing restlessness and need to move often accompanied by an inability to sit or stand still. This is most likely to occur within the first few weeks of treatment. In patients who develop these symptoms, increasing the dose may be detrimental.

4.5 Interaction with other medicinal products and other forms of interaction

Monoamine oxidase inhibitors (MAOIs): due to the risk of serotonin syndrome, ARICLAIM should not be used in combination with nonselective, irreversible monoamine oxidase inhibitors (MAOIs), or within at least 14 days of discontinuing treatment with an MAOI. Based on the half-life of duloxetine, at least 5 days should be allowed after stopping ARICLAIM before starting an MAOI (see section 4.3).

Serotonin syndrome: in rare cases, serotonin syndrome has been reported in patients using SSRIs concomitantly with serotonergic medicinal products. The use of ARICLAIM in combination with serotonergic antidepressants like SSRIs, tricyclics like clomipramine or amitriptyline, venlafaxine, or triptans, tramadol and tryptophan is not recommended.

CNS drugs: caution is advised when ARICLAIM is taken in combination with other centrally acting drugs or substances, including alcohol and sedative drugs (benzodiazepines, morphinomimetics, antipsychotics, phenobarbital, sedative antihistamines).

Effect of duloxetine on other drugs

Medicinal products metabolised by CYP1A2: The pharmacokinetics of theophylline, a CYP1A2 substrate, were not significantly affected by co-administration with duloxetine (60 mg twice daily).

Medicinal products metabolised by CYP2D6: Duloxetine is a moderate inhibitor of CYP2D6. When duloxetine was administered at a dose of 60 mg twice daily with a single dose of desipramine, a CYP2D6 substrate, the AUC of desipramine increased 3-fold. The co-administration of duloxetine (40 mg twice daily) increases steady state AUC of tolterodine (2 mg twice daily) by 71 %, but does not affect the pharmacokinetics of its active 5-hydroxyl metabolite and no dosage adjustment is recommended. Caution is advised if ARICLAIM is co-administered with medicinal products that are predominantly metabolised by CYP2D6 (risperidone, tricyclic antidepressants [TCAs] such as nortriptyline, amitriptyline, and imipramine) particularly if they have a narrow therapeutic index (such as flecainide, propafenone and metoprolol).

Oral contraceptives and other steroidal agents: results of *in vitro* studies demonstrate that duloxetine does not induce the catalytic activity of CYP3A. Specific *in vivo* drug interaction studies have not been performed.

Anticoagulants and antiplatelet agents: Caution should be exercised when duloxetine is combined

with oral anticoagulants or antiplatelet agents due to a potential increased risk of bleeding. Furthermore, increases in INR values have been reported when duloxetine was co-administered with warfarin.

Effects of other drugs on duloxetine

Antacids and H₂ antagonists: co-administration of ARICLAIM with aluminium- and magnesium-containing antacids or with famotidine had no significant effect on the rate or extent of duloxetine absorption after administration of a 40 mg oral dose.

Inhibitors of CYP1A2: because CYP1A2 is involved in duloxetine metabolism, concomitant use of ARICLAIM with potent inhibitors of CYP1A2 is likely to result in higher concentrations of duloxetine. Fluvoxamine (100 mg once daily), a potent inhibitor of CYP1A2, decreased the apparent plasma clearance of duloxetine by about 77% and increased AUC₀₋₁ 6-fold. Therefore ARICLAIM should not be administered in combination with potent inhibitors of CYP1A2 like fluvoxamine (see section 4.3).

Inducers of CYP1A2: Population pharmacokinetic studies analyses have shown that smokers have almost 50% lower plasma concentrations of duloxetine compared with non-smokers.

4.6 Pregnancy and lactation

Pregnancy

There are no data on the use of duloxetine in pregnant women. Studies in animals have shown reproductive toxicity at systemic exposure levels (AUC) of duloxetine lower than the maximum clinical exposure (see section 5.3). The potential risk for humans is unknown. Withdrawal symptoms may occur in the neonate after maternal duloxetine use near term. ARICLAIM is contraindicated during pregnancy (see section 4.3).

Breast feeding

Duloxetine is very weakly excreted into human milk based on a study of 6 lactating patients, who did not breast feed their children. The estimated daily infant dose on a mg/kg basis is approximately 0.14% of the maternal dose (see section 5.2). As the safety of duloxetine in infants is not known, the use of ARICLAIM while breast-feeding is not recommended.

4.7 Effects on ability to drive and use machines

Although in controlled studies duloxetine has not been shown to impair psychomotor performance, cognitive function, or memory, it may be associated with sedation and dizziness. Therefore, patients should be cautioned about their ability to drive a car or operate hazardous machinery.

4.8 Undesirable effects

Table 1 gives the adverse reactions observed from spontaneous reporting and in placebo-controlled clinical trials (comprising a total of 7977 patients, 4371 on duloxetine and 3606 on placebo) in SUI and other lower urinary tract disorders.

The most commonly reported adverse events in patients treated with ARICLAIM in clinical trials in SUI and other lower urinary tract disorders were nausea, dry mouth and fatigue. The data analysis of four 12-week, placebo-controlled clinical trials in patients with SUI, including 958 duloxetine-treated and 955 placebo-treated patients, showed that the onset of the reported adverse events typically occurred in the first week of therapy. However, the majority of the most frequent adverse events were mild to moderate and resolved within 30 days of occurrence (e.g. nausea).

Table 1: Adverse reactions

Frequency estimate: Very common (≥10%), common (≥1% and <10%), uncommon (≥0.1% and <1%), rare (≥0.01% and <0.1%), very rare (<0.01%), frequency not known (data from spontaneous reports)

Very common	Common	Uncommon	Rare	Very Rare	Frequency not known
<i>Infections and infestations</i>					
		Laryngitis			
<i>Immune system disorders</i>					
		Hyper-sensitivity disorder			Anaphylactic reaction
<i>Endocrine disorders</i>					
		Hypo-thyroidism			
<i>Metabolism and Nutrition Disorders</i>					
	Appetite decreased	Dehydration			Hyponatremia, SIADH
<i>Psychiatric Disorders</i>					
	Insomnia Sleep disorder Anxiety Libido decreased Agitation	Bruxism Disorientation Orgasm abnormal Apathy Abnormal dreams			Hallucinations Mania
<i>Nervous System Disorders</i>					
	Headache Dizziness Tremor Nervousness Lethargy Somnolence Paraesthesia	Dysgeusia Disturbance in attention	Myoclonus Dyskinesia		Serotonin syndrome Extrapyramidal symptoms Convulsions Akathisia Psychomotor restlessness
<i>Eye Disorders</i>					
	Blurred vision	Mydriasis Visual disturbance	Glaucoma		
<i>Ear and Labyrinth Disorders</i>					
	Vertigo	Ear pain			
<i>Cardiac Disorders</i>					
	Palpitations	Tachycardia			Supra-ventricular arrhythmia, mainly atrial fibrillation
<i>Vascular Disorders</i>					
	Hot flush	Flushing Blood pressure increase Syncope ¹	Peripheral coldness Orthostatic hypotension ¹ Hypertensive crisis		Hypertension
<i>Respiratory, thoracic and mediastinal disorders</i>					
		Yawning	Throat tightness		

<i>Gastrointestinal Disorders</i>					
Nausea (22%) Dry mouth (11.2%)	Constipation Diarrhoea Vomiting Dyspepsia	Eructation Gastroenteritis Stomatitis Halitosis Gastritis Flatulence			
<i>Hepato-biliary disorders</i>					
		Elevated liver enzymes (ALT, AST, alkaline phosphatase) Hepatitis ² Acute liver injury			Jaundice Hepatic failure
<i>Skin and Subcutaneous Tissue Disorders</i>					
	Sweating increased	Night sweats Cold sweat Rash	Photo-sensitivity reactions		Angioneurotic oedema Stevens-Johnson Syndrome Urticaria
<i>Muscoskeletal and connective tissue disorders</i>					
		Muscle tightness Musculo-skeletal pain Trismus	Muscle twitching		
<i>Renals and Urinary Disorders</i>					
		Nocturia Urinary hesitation Urine odour abnormal Dysuria			Urinary retention
<i>Reproductive System and Breast Disorders</i>					
		Menopausal symptoms			
<i>General Disorders and Administration Site Conditions</i>					
Fatigue (10.3%)	Pruritus Weakness Abdominal pain Chills	Feeling hot Malaise Thirst Feeling abnormal	Feeling cold		Chest pain
<i>Investigations</i>					
		Weight increase Weight decrease Creatinine phosphokinase increase			

		Blood cholesterol increased			
--	--	-----------------------------	--	--	--

¹Cases of orthostatic hypotension and syncope have been reported especially at the initiation of treatment

²See section 4.4

Discontinuation of duloxetine (particularly when abrupt) commonly leads to withdrawal symptoms. Dizziness, sensory disturbances (including paraesthesia), sleep disturbances (including insomnia and intense dreams), agitation or anxiety, nausea and/or vomiting, tremor and headache are the most commonly reported reactions.

Generally, for SSRIs and SNRIs, these events are mild to moderate and self-limiting, however, in some patients they may be severe and/or prolonged. It is therefore advised that when duloxetine treatment is no longer required, gradual discontinuation by dose tapering should be carried out (see sections 4.2 and 4.4).

Electrocardiograms were obtained from 755 duloxetine-treated patients with SUI and 779 placebo-treated patients in 12-week clinical trials. The heart rate-corrected QT interval in duloxetine-treated patients did not differ from that seen in placebo-treated patients.

In the 12 week acute phase of three clinical trials of duloxetine in patients with diabetic neuropathic pain, small but statistically significant increases in fasting blood glucose were observed in duloxetine-treated patients. HbA_{1c} was stable in both duloxetine-treated and placebo-treated patients. In the extension phase of these studies, which lasted up to 52 weeks, there was an increase in HbA_{1c} in both the duloxetine and routine care groups, but the mean increase was 0.3% greater in the duloxetine-treated group. There was also a small increase in fasting blood glucose and in total cholesterol in duloxetine-treated patients while those laboratory tests showed a slight decrease in the routine care group.

4.9 Overdose

There is limited clinical experience with duloxetine overdose in humans. Cases of overdoses, alone or in combination with other drugs, with duloxetine doses of almost 2000 mg were reported. Some fatalities have occurred, primarily with mixed overdoses, but also with duloxetine alone at a dose of approximately 1000 mg. Signs and symptoms of overdose (mostly with mixed drugs) included serotonin syndrome, somnolence, vomiting and seizures.

No specific antidote for duloxetine is known but if serotonin syndrome ensues, specific treatment (such as with cyproheptadine and/or temperature control) may be considered. A free airway should be established. Monitoring of cardiac and vital signs is recommended, along with appropriate symptomatic and supportive measures. Gastric lavage may be indicated if performed soon after ingestion or in symptomatic patients. Activated charcoal may be useful in limiting absorption. Duloxetine has a large volume of distribution and forced diuresis, haemoperfusion, and exchange perfusion are unlikely to be beneficial.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Other antidepressants. ATC code: N06AX21

Duloxetine is a combined serotonin (5-HT) and norepinephrine (NE) reuptake inhibitor. It weakly inhibits dopamine reuptake with no significant affinity for histaminergic, dopaminergic, cholinergic and adrenergic receptors.

In animal studies, increased levels of 5-HT and NE in the sacral spinal cord, lead to increased urethral tone via enhanced pudendal nerve stimulation to the urethral striated sphincter muscle only during the

storage phase of the micturition cycle. A similar mechanism in women is believed to result in stronger urethral closure during urine storage with physical stress that could explain the efficacy of duloxetine in the treatment of women with SUI.

The efficacy of duloxetine 40 mg given twice daily in the treatment of SUI was established in four double-blind, placebo-controlled studies, that randomised 1913 women (22 to 83 years) with SUI; of these, 958 patients were randomised to duloxetine and 955 to placebo. The primary efficacy measures were Incontinence Episode Frequency (IEF) from diaries and an incontinence specific quality of life questionnaire score (I-QOL).

Incontinence Episode Frequency: in all four studies the duloxetine-treated group had a 50% or greater median decrease in IEF compared with 33% in the placebo-treated group. Differences were observed at each visit after 4 weeks (duloxetine 54%; and placebo 22%), 8 weeks (52% and 29%), and 12 weeks (52% and 33%) of medication.

In an additional study limited to patients with severe SUI, all responses with duloxetine were achieved within 2 weeks.

The efficacy of ARICLAIM has not been evaluated for longer than 3 months in placebo-controlled studies. The clinical benefit of ARICLAIM compared with placebo has not been demonstrated in women with mild SUI, defined in randomised trials as those with IEF < 14 per week. In these women, ARICLAIM may provide no benefit beyond that afforded by more conservative behavioural interventions.

Quality of Life: Incontinence Quality of Life (I-QOL) questionnaire scores were significantly improved in the duloxetine-treated patient group compared with the placebo-treated group (9.2 versus 5.9 score improvement, $p < .001$). Using a global improvement scale (PGI), significantly more women using duloxetine considered their symptoms of stress incontinence to be improved with treatment compared with women using placebo (64.6% versus 50.1%, $p < .001$).

ARICLAIM and Prior Continence Surgery: there are limited data that suggest that the benefits of ARICLAIM are not diminished in women with stress urinary incontinence who have previously undergone continence surgery.

ARICLAIM and Pelvic Floor Muscle Training (PFMT): during a 12-week blinded, randomised, controlled study, ARICLAIM demonstrated greater reductions in IEF compared with either placebo treatment or with PFMT alone. Combined therapy (duloxetine + PFMT) showed greater improvement in both pad use and condition-specific quality of life measures than ARICLAIM alone or PFMT alone.

5.2 Pharmacokinetic properties

Duloxetine is administered as a single enantiomer. Duloxetine is extensively metabolised by oxidative enzymes (CYP1A2 and the polymorphic CYP2D6), followed by conjugation. The pharmacokinetics of duloxetine demonstrate large intersubject variability (generally 50-60%), partly due to gender, age, smoking status and CYP2D6 metaboliser status.

Duloxetine is well absorbed after oral administration with a C_{max} occurring 6 hours post dose. The absolute oral bioavailability of duloxetine ranged from 32% to 80% (mean of 50%; N=8 subjects). Food delays the time to reach the peak concentration from 6 to 10 hours and it marginally decreases the extent of absorption (approximately 11%).

Duloxetine is approximately 96% bound to human plasma proteins. Duloxetine binds to both albumin and alpha-1 acid glycoprotein. Protein binding is not affected by renal or hepatic impairment.

Duloxetine is extensively metabolised and the metabolites are excreted principally in urine. Both CYP2D6 and CYP1A2 catalyse the formation of the two major metabolites glucuronide conjugate of 4-hydroxy duloxetine and sulphate conjugate of 5-hydroxy,6-methoxy duloxetine. Based upon *in vitro*

studies, the circulating metabolites of duloxetine are considered pharmacologically inactive. The pharmacokinetics of duloxetine in patients who are poor metabolisers with respect to CYP2D6 has not been specifically investigated. Limited data suggest that the plasma levels of duloxetine are higher in these patients.

The elimination half-life of duloxetine after an oral dose ranges from 8 to 17 hours (mean of 12 hours). After an intravenous dose the plasma clearance of duloxetine ranges from 22 l/hr to 46 l/hr (mean of 36 l/hr). After an oral dose the apparent plasma clearance of duloxetine ranges from 33 to 261 l/hr (mean 101 l/hr).

Special populations:

Age: pharmacokinetic differences have been identified between younger and elderly females (≥ 65 years) (AUC increases by about 25% and half-life is about 25% longer in the elderly), although the magnitude of these changes is not sufficient to justify adjustments to the dose.

Renal impairment: end stage renal disease (ESRD) patients receiving dialysis had a 2-fold higher duloxetine C_{\max} and AUC values compared to healthy subjects. Pharmacokinetic data on duloxetine is limited in patients with mild or moderate renal impairment.

Hepatic insufficiency: moderate liver disease (Child Pugh Class B) affected the pharmacokinetics of duloxetine. Compared with healthy subjects, the apparent plasma clearance of duloxetine was 79% lower, the apparent terminal half-life was 2.3 times longer, and the AUC was 3.7 times higher in patients with moderate liver disease. The pharmacokinetics of duloxetine and its metabolites have not been studied in patients with mild or severe hepatic insufficiency.

Nursing mothers: The disposition of duloxetine was studied in 6 lactating women who were at least 12-weeks postpartum. Duloxetine is detected in breast milk, and steady-state concentrations in breast milk are about one-fourth those in plasma. The amount of duloxetine in breast milk is approximately 7 $\mu\text{g/day}$ while on 40 mg twice daily dosing. Lactation did not influence duloxetine pharmacokinetics.

5.3 Preclinical safety data

Duloxetine was not genotoxic in a standard battery of tests and was not carcinogenic in rats. Multinucleated cells were seen in the liver in the absence of other histopathological changes in the rat carcinogenicity study. The underlying mechanism and the clinical relevance are unknown.

Female mice receiving duloxetine for 2 years had an increased incidence of hepatocellular adenomas and carcinomas at the high dose only (144 mg/kg/day), but these were considered to be secondary to hepatic microsomal enzyme induction. The relevance of this mouse data to humans is unknown. Female rats receiving duloxetine before and during mating and early pregnancy had a decrease in maternal food consumption and body weight, oestrous cycle disruption, decreased live birth indices and progeny survival, and progeny growth retardation at systemic exposure levels estimated to be at the most at maximum clinical exposure (AUC). In an embryotoxicity study in the rabbit, a higher incidence of cardiovascular and skeletal malformations was observed at systemic exposure levels below the maximum clinical exposure (AUC). No malformations were observed in another study testing a higher dose of a different salt of duloxetine. In pre/postnatal toxicity study in the rat, duloxetine induced adverse behavioural effects in the offspring at systemic exposures levels below maximum clinical exposure (AUC).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content:

Hypromellose.

Hydroxypropyl methylcellulose acetate succinate

Sucrose
Sugar spheres
Talc
Titanium dioxide (E171)
Triethyl citrate.

Capsule Shell:

Gelatin
Sodium Lauryl Sulfate
Titanium Dioxide (E171)
Indigo Carmine (E132)
Edible Black Ink.

Edible Ink:
Black Iron Oxide-Synthetic (E172)
Propylene glycol
Shellac

Capsule Shell Cap colour:

Opaque Blue

Capsule Shell Body colour:

Opaque blue

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Store in the original package. Do not store above 30°C.

6.5 Nature and contents of container

Polyvinylchloride (PVC), Polyethylene (PE), and Polychlorotrifluoroethylene (PCTFE) blister sealed with an aluminum foil.

Packs of 28 and 56 capsules.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements.

7. MARKETING AUTHORISATION HOLDER

Eli Lilly Nederland BV, Grootslag 1-5, NL-3991 RA Houten, The Netherlands.

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/04/283/001
EU/1/04/283/007

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

11 August 2004

10. DATE OF REVISION OF THE TEXT

ANNEX II

- A. MANUFACTURING AUTHORISATION HOLDER(S)
RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OF THE MARKETING AUTHORISATION**

A MANUFACTURING AUTHORISATION HOLDER(S) RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer(s) responsible for batch release

Lilly SA
Avenida de la Industria No 30
28108 Alcobendas
Madrid
Spain

B CONDITIONS OF THE MARKETING AUTHORISATION

- **CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IMPOSED ON THE MARKETING AUTHORISATION HOLDER**

Medicinal product subject to medical prescription.

- **CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

Not applicable.

- **OTHER CONDITIONS**

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTONS FOR 40 MG HARD GASTRO-RESISTANT CAPSULES

1. NAME OF THE MEDICINAL PRODUCT

ARICLAIM 40 mg, hard gastro-resistant capsules.
Duloxetine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 40 mg duloxetine (as hydrochloride).

3. LIST OF EXCIPIENTS

Contains sucrose.

4. PHARMACEUTICAL FORM AND CONTENTS

packs of 28, 56, 98, 140 capsules.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP {MM/YYYY}.

9. SPECIAL STORAGE CONDITIONS

Store in the original package. Do not store above 30 °C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Eli Lilly Nederland BV, Grootslag 1-5, NL-3991 RA, Houten, The Netherlands.

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/04/283/002-005

13. BATCH NUMBER

Lot:

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

ARICLAIM 40 mg

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON FOR 98 CAPSULES (40 MG) AS INTERMEDIATE PACK / COMPONENT OF A MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

ARICLAIM 40 mg hard gastro-resistant capsules
Duloxetine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 40 mg duloxetine (as hydrochloride).

3. LIST OF EXCIPIENTS

Contains sucrose.

4. PHARMACEUTICAL FORM AND CONTENTS

98 capsules
Component of a multipack comprising 2 packs, each containing 98 capsules.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP {MM/YYYY}

9. SPECIAL STORAGE CONDITIONS

Store in the original package. Do not store above 30 °C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Eli Lilly Nederland BV, Grootslag 1-5, NL-3991 RA, Houten, The Netherlands.

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/04/283/004

13. BATCH NUMBER

Lot:

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

ARICLAIM 40 mg

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING
OUTER WRAPPER LABEL ON MULTIPACKS (2X98 CAPSULES, 40 MG) WRAPPED IN
FOIL (INCLUDING BLUE BOX)**

1. NAME OF THE MEDICINAL PRODUCT

ARICLAIM 40 mg hard gastro-resistant capsules
Duloxetine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 40 mg duloxetine (as hydrochloride).

3. LIST OF EXCIPIENTS

Contains sucrose.

4. PHARMACEUTICAL FORM AND CONTENTS

Multipack comprising 2 packs, each containing 98 capsules.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT
OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP {MM/YYYY}

9. SPECIAL STORAGE CONDITIONS

Store in the original package. Do not store above 30 °C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Eli Lilly Nederland BV, Grootslag 1-5, NL-3991 RA, Houten, The Netherlands.

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/04/283/006

13. BATCH NUMBER

Lot:

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

ARICLAIM 40 mg

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS (40 mg hard gastro-resistant capsules)

1. NAME OF THE MEDICINAL PRODUCT

ARICLAIM 40 mg hard gastro-resistant capsules
Duloxetine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Lilly

3. EXPIRY DATE

<EXP {MM/YYYY}.

4. BATCH NUMBER

Lot:

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTONS FOR 20 MG HARD GASTRO-RESISTANT CAPSULES

1. NAME OF THE MEDICINAL PRODUCT

ARICLAIM 20 mg, hard gastro-resistant capsules.
Duloxetine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 20 mg duloxetine (as hydrochloride).

3. LIST OF EXCIPIENTS

Contains sucrose.

4. PHARMACEUTICAL FORM AND CONTENTS

Packs of 28 capsules.
Packs of 56 capsules.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the leaflet before use
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP {MM/YYYY}.

9. SPECIAL STORAGE CONDITIONS

Store in the original package. Do not store above 30 °C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Eli Lilly Nederland BV, Grootslag 1-5, NL-3991 RA, Houten, The Netherlands.

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/04/283/001

EU/1/04/283/007

13. BATCH NUMBER

Lot:

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

ARICLAIM 20 mg

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS (20 mg hard gastro-resistant capsules)

1. NAME OF THE MEDICINAL PRODUCT

ARICLAIM 20 mg hard gastro-resistant capsules
Duloxetine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Lilly

3. EXPIRY DATE

<EXP {MM/YYYY}.

4. BATCH NUMBER

Lot:

5. OTHER

B. PACKAGE LEAFLET

PACKAGE LEAFLET : INFORMATION FOR THE USER

ARICLAIM 40 mg hard gastro resistant capsules ARICLAIM 20 mg hard gastro resistant capsules Duloxetine (as hydrochloride)

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What ARICLAIM is and what it is used for
2. Before you take ARICLAIM
3. How to take ARICLAIM
4. Possible side effects
5. How to store ARICLAIM
6. Further information

1. WHAT ARICLAIM IS AND WHAT IT IS USED FOR

ARICLAIM is a medicine to be taken by mouth to treat Stress Urinary Incontinence (SUI) in women.

Stress urinary incontinence is a medical condition in which patients have accidental loss or leakage of urine during physical exertion or activities such as laughing, coughing, sneezing, lifting, or exercise.

ARICLAIM is believed to work by increasing the strength of the muscle that holds back urine when you laugh, sneeze, or perform physical activities.

The efficacy of ARICLAIM is reinforced when combined with a training program called Pelvic Floor Muscle Training (PFMT).

2. BEFORE YOU TAKE ARICLAIM

ARICLAIM can only be prescribed by a doctor.

Do not take ARICLAIM

- If you are allergic (hypersensitive) to duloxetine or any of the inactive ingredients of ARICLAIM.
- If you are taking or have taken within the last 14 days, another medicine known as a monoamine oxidase inhibitor - MAOI (see section below 'Taking other medicines').
- If you have liver disease.
- If you are pregnant or breast-feeding.
- If you are taking a potent inhibitor of a liver enzyme called CYP1A2, like fluvoxamine, ciprofloxacin or enoxacin.
- If you have severe kidney disease.
- If you suffer from uncontrolled high blood pressure.

Take special care with ARICLAIM

The following are reasons why ARICLAIM may not be suitable for you. If any of them apply to

you, talk to your doctor before you take the medicine:

- If you are taking medicines to treat depression.
- You have kidney disease.
- You have a history of seizures (fits).
- You have a history of mania or bipolar disorder.
- You have eye problems such as certain kinds of glaucoma (increased pressure in the eye).
- You have a history of bleeding disorders (tendency to develop bruises).
- If you are younger than 18 years.
- If you have high blood pressure.

ARICLAIM may cause a sensation of restlessness or an inability to sit or stand still. You should tell your doctor if this happens to you.

Although ARICLAIM is not indicated for the treatment of depression, its active ingredient (duloxetine) also exists as an antidepressant medication. Isolated cases of suicidal thoughts and behaviours have been reported during duloxetine therapy or early after treatment discontinuation. Tell your doctor immediately if you have any distressing thoughts or feelings at any time or if you are feeling changes in mood like sadness, apathy or agitation, or if you start treatment for depression.

Use in children and adolescents under 18 years of age

ARICLAIM should not be used for children and adolescents under the age of 18 years. Also, you should know that patients under 18 have an increased risk of side-effects such as suicide attempt, suicidal thoughts and hostility (predominantly aggression, oppositional behaviour and anger) when they take this class of medicines. Also, the long-term safety effects concerning growth, maturation, and cognitive and behavioural development of ARICLAIM in this age group have not yet been demonstrated.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. The main ingredient of ARICLAIM, duloxetine, is used in other medicines for other conditions (diabetic neuropathic pain, depression and urinary incontinence). Using more than one of these medicines at the same time should be avoided. Check with your doctor if you are already taking other medicines containing duloxetine.

Your doctor should decide whether you can take ARICLAIM with other medicines. Do not start or stop taking any medicines, including those bought without a prescription and herbal remedies, before checking with your doctor.

Monoamine Oxidase Inhibitors (MAOI): you should not take ARICLAIM with an MAOI or within 14 days of stopping an MAOI. Taking an MAOI together with many prescription medicines, including ARICLAIM, can cause serious or even life-threatening side effects. You must wait at least 14 days after you have stopped taking an MAOI before you can take ARICLAIM. Also, you need to wait at least 5 days after you stop taking ARICLAIM before you take an MAOI.

CNS drugs: caution is advised when ARICLAIM is taken in combination with other centrally acting drugs or substances, including alcohol and sedative drugs (benzodiazepines, morphinomimetics, antipsychotics, phenobarbital, sedative antihistamines). Inform your doctor if you are taking any of these drugs.

Serotonin syndrome: you should tell your doctor if you are taking any of the medicines that act in a similar way to duloxetine. Examples of these medicines include: triptans, tramadol, tryptophan, certain antidepressants: SSRIs (such as paroxetine and fluoxetine), tricyclics (such as clomipramine, amitriptyline) and venlafaxine. These medicines increase the risk of side effects; if you get any unusual symptom taking any of these medicines together with ARICLAIM, you should see your doctor.

Oral -anticoagulants: You should tell your doctor if you are taking oral –anticoagulants (medicines which thin the blood). These medicines might increase the risk of bleeding.

Taking ARICLAIM with food and drink

ARICLAIM may be taken with or without food. You should take extra care if you drink alcohol while taking ARICLAIM.

Pregnancy and breast-feeding

ARICLAIM should not be used during pregnancy and if you are breast feeding. Tell your doctor if you become pregnant, or you are trying to become pregnant, while you are taking ARICLAIM.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Do not drive or use any tools or machines until you know how ARICLAIM affects you.

Important information about some of the ingredients of ARICLAIM

ARICLAIM contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. HOW TO TAKE ARICLAIM

Always take ARICLAIM exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

The recommended dose of ARICLAIM is one capsule of 40 mg twice a day (in the morning and late afternoon/evening). Your doctor may decide to start your treatment with one capsule of 20 mg twice a day for two weeks before increasing the dose to 40 mg twice a day.

ARICLAIM is for oral use. You should swallow your capsule whole with a drink of water.

To help you remember to take ARICLAIM, you may find it easier to take it at the same times every day.

Do not stop taking ARICLAIM without talking to your doctor.

If you take more ARICLAIM than you should

Call your doctor or pharmacist immediately if you take more than the amount of ARICLAIM prescribed by your doctor.

If you forget to take ARICLAIM

Do not take a double dose to make up for forgotten doses.

If you miss a dose, take it as soon as you remember. However, if it is time for your next dose, skip the missed dose and take only a single dose as usual. Do not take more than the daily amount of ARICLAIM that has been prescribed for you in one day.

If you stop taking ARICLAIM

Do not stop taking your capsules without the advice of your doctor even if you feel better. If your doctor thinks that you no longer need ARICLAIM he will ask you to reduce your dose over 2 weeks. Some patients, who suddenly stop taking ARICLAIM after more than 1 week of therapy, have felt dizzy, sick (nausea) or had a headache. These symptoms are usually not serious and disappear within a few days, but if you have symptoms that are troublesome you should ask your doctor for advice.

4. POSSIBLE SIDE EFFECTS

Like all medicines, ARICLAIM can cause side effects, although not everybody gets them. These effects are normally mild to moderate and often disappear after a short time.

The most common side effects with ARICLAIM are feeling sick (nausea), dry mouth. and tiredness.. Other common and less common side effects are listed below:

Infections: Throat inflammation.

Psychiatric disorders: Trouble sleeping, anxiety, nervousness or feeling agitated . Less common side effects affecting you mentally are feeling disorientated, having abnormal dreams , lack of motivation or experiencing hallucinations and mania (a disorder which symptoms are over activity, racing thoughts and decreased need for sleep). ARICLAIM may also make you feel sleepy or increase yawning.

Nervous system disorders: Dizziness, headache, tremor and numbness, including numbness or tingling of the skin, disturbance in attention. Less common effects could be tasting things differently than usual, fits, stiffness, spasms and involuntary movements of the muscles, a sensation of restlessness or an inability to sit or stand still or “Serotonin syndrome” (a rare reaction which may cause feelings of great happiness, drowsiness, clumsiness, restlessness, feeling of being drunk, fever, sweating or rigid muscles).

Digestive disorders: Constipation, diarrhoea, being sick (vomiting) or heartburn. You may also have a lack of appetite and weight change (loss or increase). Less common side effects on the digestive system are burping, indigestion, gastroenteritis, bad breath or breaking wind.

Liver disorders: Inflammation of the liver that may cause abdominal pain, tiredness and yellow coloration of the skin.

Ear disorders: Vertigo and ear pain.

Eye disorders: Blurred eyesight is common, but less common effects are large pupils (the dark centre of the eye), visual disturbance or increased pressure in the eye.

Heart or circulation disorders: The most common effects are feeling the heart pumping in the chest, hot flushes, shivering, weakness or increased sweating. Less common effects are flushing, increase in blood pressure, feeling cold in your fingers and/or toes, feeling dizzy (particularly when standing up too quickly), fast or irregular heart beat, night sweats, cold sweats or fainting.

Reproductive system disorders: sexual problems (less sex drive, changes in orgasm).

Skin disorders: Allergic reactions (itchy) rash, blisters or sensitivity to sunlight.

Bone and muscle disorders: Muscle pain, muscle tightness, twitching and contraction of the jaw muscle.

Urinary system disorders: Some patients need to pass urine during the night or may have difficulty or inability to pass urine. Some patients may have abnormal urine odour.

General disorders: Less common side effects include grinding of teeth, dehydration, feeling hot/cold, thirst or throat tightness, decrease of activity of the thyroid gland, chest or stomach pain or stiffness.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. HOW TO STORE ARICLAIM

Keep out of the reach and sight of children

Store in the original pack. Do not store above 30 °C.

Do not use ARICLAIM after the expiry date which is stated on the carton.

6. FURTHER INFORMATION

What ARICLAIM contains

ARICLAIM is available in 2 strengths: 20 and 40 mg. The active substance is duloxetine. Each capsule contains 20 or 40 mg of duloxetine (as duloxetine hydrochloride)

The other ingredients are:

Capsule content: hypromellose, hydroxypropyl methylcellulose acetate succinate, sucrose, sugar spheres, talc, titanium dioxide (E171), triethyl citrate.

Capsule shell: gelatin, sodium lauryl sulphate, titanium dioxide (E171), indigo carmine (E132), iron oxide red and iron oxide yellow, edible black ink.

Edible Ink: Black Iron Oxide-Synthetic (E172), Propylene glycol, Shellac.

What ARICLAIM looks like and contents of the pack

ARICLAIM is a hard gastro-resistant capsule.

Each capsule of ARICLAIM contains pellets of the active substance with a covering to protect them from stomach acid.

The 40 mg capsule has an opaque orange body imprinted with '40 mg' and an opaque blue cap, imprinted with '9545'.

The 20 mg capsule has an opaque blue body imprinted with '20 mg' and an opaque blue cap, imprinted with '9544'.

ARICLAIM 40 mg is available in blister packs of 28, 56, 98, 140 and 196 (2 x 98) capsules.

ARICLAIM 20 mg is available in blister packs of 28 and 56 capsules.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder: Marketing Authorisation Holder: Eli Lilly Nederland BV, Grootslag 1-5, NL-3991 RA Houten, The Netherlands.

Manufacturer: Lilly S.A., Avda. De la Industria, 30, 28108 Alcobendas, Madrid, Spain.

For any information about this medicine product, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien

S.A. Eli Lilly N.V.

Tél/Tel: + 32-(0)2 548 84 84

Česká republika

Eli Lilly ČR, s.r.o.

Tel: + 420 234 664 111

Danmark

Eli Lilly Danmark A/S

Tlf.: +45 45 26 60 00

Deutschland

Lilly Deutschland GmbH

Tel. + 49-(0) 6172 273 2222

Eesti

Eli Lilly Holdings Limited Eesti filiaal

Tel: +372 6817280

Ελλάδα

ΦΑΡΜΑΣΕΠΒ-ΛΙΑΛΥ Α.Ε.Β.Ε.

Τηλ: +30 210 629 4600

Luxembourg/Luxemburg

S.A. Eli Lilly N.V.

Tél/Tel: + 32-(0)2 548 84 84

Magyarország

Lilly Hungária Kft.

Tel: + 36 1 328 5100

Malta

Charles de Giorgio Ltd.

Tel: + 356 25600 500

Nederland

Eli Lilly Nederland B.V.

Tel: + 31(0)30 6025800

Norge

Eli Lilly Norge A.S

Tlf: +47 22 88 18 00

Österreich

Eli Lilly Ges.m.b.H.

Tel: + 43-(0) 1 711 780

España

Lilly S.A.

Tel: +34-91 623 17 32

France

Lilly France SAS.

Tel: +33-(0) 1 55 49 34 34

Ireland

Eli Lilly and Company (Ireland) Limited

Tel: +353 (0) 1 661 4377

Ísland

Eli Lilly Danmark A/S, Útibú á Íslandi

Tel: +354 520 34 00

Italia

Eli Lilly Italia S.p.A.

Tel: +39- 055 42571

Κύπρος

Phadisco Ltd

Τηλ: +357 22 715000

Latvija

Eli Lilly Holdings Limited pārstāvniecība Latvijā

Tel: +371 7364000

Lietuva

Eli Lilly Holdings Limited atstovybė

Tel. +370 (5) 2649600

Polska

Eli Lilly Polska Sp. z o.o.

Tel. +48 (0) 22 440 33 00

Portugal

Lilly Portugal - Produtos

Farmacêuticos, Lda

Tel: +351 21 412 66 32

Slovenija

Eli Lilly, Podružnica Ljubljana

Tel: +386 (0)1 580 00 10

Slovenská republika

Eli Lilly Slovakia, s.r.o.

Tel: +421 (2) 59224 111

Suomi/Finland

Oy Eli Lilly Finland Ab

Puh/Tel: +358 (0)9 8545 250

Sverige

Eli Lilly Sweden AB

Tel: +46-(0)8 7378800

United Kingdom

Eli Lilly and Company Limited

Tel: +44 (0) 1256 315999

This leaflet was last approved in